

RESULTANTS AND THE ALGEBRAICITY OF THE JOIN PAIRING ON CHOW VARIETIES

JUDITH PLÜMER

ABSTRACT. The Chow/Van der Waerden approach to algebraic cycles via resultants is used to give a purely algebraic proof for the algebraicity of the complex suspension. The algebraicity of the join pairing on Chow varieties then follows. The approach implies a more algebraic proof of Lawson's complex suspension theorem in characteristic 0. The continuity of the action of the linear isometries operad on the group completion of the stable Chow variety is a consequence.

INTRODUCTION

In [14] Lawson showed that a suitable stabilization \mathcal{C} of the Chow variety of effective algebraic cycles on projective spaces is homotopy equivalent to the product $\prod_{i>0} K(\mathbb{Z}, 2i) =: K(\mathbb{Z}, \text{even})$ of Eilenberg Mac Lane spaces $K(\mathbb{Z}, 2i)$. The proof of the equivalence is based on the complex suspension theorem, which is fundamental for Lawson homology [8]. The original proof of this theorem uses geometric measure theory. Here we provide an algebraic and, as we think, more elementary proof (Thm. 3.1) which goes back to the construction of Chow varieties via resultants [5]. Once the basic continuity statements are proved, we can stay close to the geometric intuition [14], which is behind the measure theoretic arguments, by the Chow/Van der Waerden approach to Chow varieties. The main observation is that one can compute the Chow form of a suspended cycle (Prop. 2.2) from the original Chow form, showing that the suspension is an algebraic map.

In order to prove that the join pairing is algebraic (Thm. 2.3), hence continuous, one just has to combine the algebraicity of the complex suspension with well known results (Barlet [1]). (Independently Barlet [2] obtained a proof of Thm. 2.3 using different methods.)

In [4] the join pairing was used to define an infinite loop space structure on $K(\mathbb{Z}, \text{even})$ which is compatible with the infinite loop space structure on BU induced by Whitney sum. The required continuity statements (Lemma 4.1) for the operation of the linear isometries operad [4] also follow easily in the Chow/Van der Waerden picture. So there are three sets of results:

Received by the editors May 26, 1995.

1991 *Mathematics Subject Classification.* Primary 14C25; Secondary 55N20.

Key words and phrases. Resultants, Chow varieties, join pairing, infinite loop spaces.

This paper is an outgrowth of my diploma thesis [20]. I am indebted to R. Schwänzl for suggesting the problem to me, to P. Lima-Filho for calling my attention to [2], and to the DFG for support.

- The explicit description of Chow varieties and the complex suspension via Chow coordinates (sections 1, 2), which makes the join pairing an algebraic map.
- Another, more algebraic, proof of the complex suspension theorem in characteristic 0 (section 3). (For a quite different algebraization see [8].)
- The continuity of the action of the linear isometries operad on the group completion of a stabilized Chow variety (section 4), which makes the total Chern class map an infinite loop map.

Throughout this paper we work in the category \mathbf{kTop} of compactly generated, weak Hausdorff spaces [18]. All proofs are given for algebraic cycles with support in $\mathbb{P}^n(\mathbb{C})$ but they directly work for cycles with support on an algebraic subset $X \subset \mathbb{P}^n(\mathbb{C})$ (see also [14]), except that one has to take care of π_0 -phenomena (compare [8], [14, p. 265]).

As a counterpart to this paper, one may view Dalbec [6], who describes an algorithm for computing Chow coordinates of joins.

1. REVIEW OF THE CHOW/VAN DER WAERDEN APPROACH TO CHOW VARIETIES

There are many different ways to define the Chow form of a variety $V \subset \mathbb{P}^n(\mathbb{C})$ ([5], [12], [9]). The definition of Chow and Van der Waerden is constructive and therefore very useful for proofs of algebraicity. We first introduce some notations and basic definitions:

Definition 1.1. Let $\mathbb{C} \subset K \subset \Omega$ be fields.

- A projective algebraic set $V_K \subset \mathbb{P}^n(\Omega)$ over K is the set of common zeros of a homogeneous ideal $I \subset K[x_0, \dots, x_n]$.
- The *vanishing ideal* $I_K(V) \subset K[x_0, \dots, x_n]$ of a projective algebraic set $V_K \subset \mathbb{P}^n(\Omega)$ is the homogeneous ideal generated by

$$\{f \in K[x_0, \dots, x_n] \mid f \text{ homogeneous and } f(\underline{v}) = 0 \forall \underline{v} \in V\}.$$

An algebraic set $V_K \subset \mathbb{P}^n(\Omega)$ is called *irreducible* or *a variety* iff $I_K(V) \subset K[x_0, \dots, x_n]$ is prime. The dimension of a variety V_K is its dimension as a topological subspace of $\mathbb{P}^n(\Omega)$ in the Zariski topology. It follows from the Noetherian Normalization Theorem [13, Chap. II, Thm. 3.1] that the dimension of $V_K \subset \mathbb{P}^n(\Omega)$ is independent of the extension field Ω of K , which means

$$\dim V_K = \dim V_K \cap \mathbb{P}^n(K).$$

From now on, Ω will denote a fixed *universal field* in the sense of Van der Waerden [22], given as the algebraic closure of $\mathbb{C}(U)$, where U is a countably infinite set of algebraically independent elements over \mathbb{C} . K is always a subfield of Ω with finite transcendence degree over \mathbb{C} .

Given $\underline{\xi} = (\xi_0, \dots, \xi_n) \in \Omega^{n+1}$, one can define $K[\underline{\xi}] := K[\xi_0, \dots, \xi_n]$. If $\underline{\xi} \in \mathbb{P}^n(\Omega)$ then $K[\underline{\xi}]$ depends on the choice of a representative of $\underline{\xi}$ in Ω^{n+1} . The definition of a special representative of points in $\mathbb{P}^n(\Omega)$ makes this ring adjunction unique up to canonical isomorphisms. In the following the homogeneous coordinates of a point $\underline{\xi}$ will also be denoted (ξ_0, \dots, ξ_n) .

Definition 1.2. Let $(\xi_0, \dots, \xi_n) = \underline{\xi} \in \mathbb{P}^n(\Omega)$ and i minimal such that $\xi_i \neq 0$. Then

$$\tau \underline{\xi} := \left(\tau \frac{\xi_0}{\xi_i}, \dots, \tau \frac{\xi_i}{\xi_i}, \dots, \tau \frac{\xi_n}{\xi_i} \right) \in \Omega^{n+1}, \quad \tau \in \Omega - \overline{K(\xi_0, \dots, \xi_n)}.$$

For every point $\underline{\xi} \in \mathbb{P}^n(\Omega)$ the ring $K[\tau \underline{\xi}]$ depends on choices of τ up to canonical isomorphisms, which extend to K -automorphisms of Ω . From now on the condition $f(\underline{\xi}) = 0$, $f \in K[x_0, \dots, x_n]$, is understood as $f(\tau \underline{\xi}) = 0$ for $\underline{\xi} \in \mathbb{P}^n(\Omega)$. Also one defines $K(\tau \underline{\xi})$ to be the quotient field of $K[\tau \underline{\xi}]$.

The advantage of viewing varieties in $\mathbb{P}^n(K)$ as varieties in $\mathbb{P}^n(\Omega)$ over K is that varieties in $\mathbb{P}^n(\Omega)$ over K are completely described by single points in $\mathbb{P}^n(\Omega)$.

Definition 1.3. Let $V_K \subset \mathbb{P}^n(\Omega)$ be an algebraic set over K . A point $\underline{\xi}$ is a *general point* of V_K if it satisfies the following conditions:

- (i) $\underline{\xi} \in V_K$
- (ii) whenever $f \in K[x_0, \dots, x_n]$ satisfies $f(\underline{\xi}) = 0$, then it follows that $f \in I_K(V) \subset K[x_0, \dots, x_n]$.

As with the definition of $K[\tau \underline{\xi}]$ general points of a variety V_K are only determined up to K -automorphisms of Ω , that is, if $\underline{\xi}$ is a general point of V_K and σ a K -automorphism of Ω , then $(\sigma(\xi_0), \dots, \sigma(\xi_n))$ is a general point of V_K , too. We also say the general points are conjugate to each other. It is easy to show that an algebraic set in $\mathbb{P}^n(\Omega)$ has a general point iff it is irreducible. One can also prove that the projective coordinate ring of a variety V_K is naturally isomorphic to $K[\tau \underline{\xi}]$ for a general point $\underline{\xi}$ of V_K [22]. The isomorphism is given by

$$\begin{array}{ccc} E_{\tau \underline{\xi}} : \frac{K[x_0, \dots, x_n]}{I_K(V)} & \longrightarrow & K[\tau \underline{\xi}] \\ z & \longmapsto & z, \quad z \in K, \\ x_i & \longmapsto & (\tau \underline{\xi})_i, \quad i = 0, \dots, n. \end{array}$$

Therefore a variety V has infinitely many general points iff $\dim V > 0$.

For the definition of the Chow form of a variety over \mathbb{C} one looks at the intersections of varieties over K with K -general hyperplanes.

Definition and Notation 1.4. Let $u_i^j \in \Omega$, $i = 0, \dots, n$, $j = 1, \dots, r$, be algebraically independent over K . Denote

$$\mathcal{U}(n, r) := \begin{pmatrix} u_0^1 & \cdots & u_n^1 \\ \vdots & & \vdots \\ u_0^r & \cdots & u_n^r \end{pmatrix}.$$

The K -general linear subspace H_{n-r} in $\mathbb{P}^n(\Omega)$ of dimension $(n-r)$ is the zero set of $\mathcal{U}(n, r) \cdot \underline{x}$ viewed as r forms in $\Omega[x_0, \dots, x_n]$. H_{n-1} is also called a K -general hyperplane.

Observe that algebraic independence of the u_i^j implies linear independence of the $\underline{u}^j = (u_0^j, \dots, u_n^j)$, because linear dependence can be expressed in terms of determinants of submatrices of $\mathcal{U}(n, r)$.

Lemma 1.5. Let $V_K \subset \mathbb{P}^n(\Omega)$ be a variety of dimension r and H a K -general hyperplane given by $\langle \underline{u}, \underline{x} \rangle = 0$. Then the intersection of V_K and H is a variety over $K(\underline{u})$ of dimension $r-1$.

Proof. [23, p. 140]. □

Lemma 1.5 proves the following theorem:

Theorem 1.6. *Let $V_{\mathbb{C}} \subset \mathbb{P}^n(\Omega)$ be a variety of dimension r over \mathbb{C} and H_{n-r} a \mathbb{C} -general $(n-r)$ -dimensional linear subspace of $\mathbb{P}^n(\Omega)$. The intersection $V_{\mathbb{C}} \cap H_{n-r}$ is a 0-dimensional variety over $\mathbb{C}(\underline{u}^1, \dots, \underline{u}^r)$.*

0-dimensional varieties over a field which is not algebraically closed consist of finitely many points. All of them are general points [22, §129].

Construction and Definition 1.7. Let $V_{\mathbb{C}}$ be as in Theorem 1.6. Further let $V_{\mathbb{C}} \cap H_{n-r} = \{\underline{q}^1, \dots, \underline{q}^d\}$. As mentioned above \underline{q}^k , $k = 1, \dots, d$, are general points of $V_{\mathbb{C}} \cap H_{n-r}$ viewed as a variety over $\mathbb{C}(\underline{u}^1, \dots, \underline{u}^r)$. Therefore they are conjugate to each other over $\mathbb{C}(\underline{u}^1, \dots, \underline{u}^r)$. On the other hand it is obvious that \underline{q}^k , $k = 1, \dots, d$, are general points of $V_{\mathbb{C}}$ viewed as a variety over \mathbb{C} .

Without loss of generality, one can fix $m \in \{0, \dots, n\}$ such that $q_m^k \neq 0$ for all $k = 1, \dots, d$. Now choose

$$\underline{p}^k := \left(\frac{q_0^k}{q_m^k}, \frac{q_1^k}{q_m^k}, \dots, \frac{q_m^k}{q_m^k}, \dots, \frac{q_n^k}{q_m^k} \right) \in \Omega^{n+1}$$

as a representation of \underline{q}^k , $k = 1, \dots, d$. The p_i^k are now algebraic over $\mathbb{C}(\underline{u}^1, \dots, \underline{u}^r)$, because $V_{\mathbb{C}} \cap H_{n-r}$ has dimension 0.

Let u_0^0, \dots, u_n^0 be algebraically independent over $\mathbb{C}(\underline{u}^1, \dots, \underline{u}^r)$ and define

$$L_k := \langle \underline{u}^0, \underline{p}^k \rangle, \quad k = 1, \dots, d,$$

$$G(\underline{u}^0) := \prod_{k=1}^d L_k = \prod_{k=1}^d \sum_{i=0}^n u_i^0 p_i^k \in \overline{\mathbb{C}(\underline{u}^1, \dots, \underline{u}^r)}[\underline{u}^0].$$

Now let Z be the splitting field of the minimal polynomials of the p_i^k , $i = 0, \dots, n$, $k = 1, \dots, d$, and notice that Z is a Galois extension of $\mathbb{C}(\underline{u}^1, \dots, \underline{u}^r)$, because the characteristic of \mathbb{C} is 0. Suppose φ is an element of the Galois group of Z over $\mathbb{C}(\underline{u}^1, \dots, \underline{u}^r)$. The general points of a variety are conjugated to each other, therefore $\varphi(\underline{p}^k)$ must be equal to \underline{p}^l for some $l \in \{1, \dots, d\}$. Hence one obtains $\varphi(G(\underline{u}^0)) = G(\underline{u}^0)$, that is, the coefficients of G remain invariant under the Galois operations and are therefore elements of $\mathbb{C}(\underline{u}^1, \dots, \underline{u}^r)$.

Let $c \in \mathbb{C}[\underline{u}^1, \dots, \underline{u}^r]$ be the common denominator of the coefficients of $G(\underline{u}^0)$. Then

$$F(\underline{u}^0, \dots, \underline{u}^r) := cG(\underline{u}^0)$$

is an element of $\mathbb{C}[\underline{u}^0, \dots, \underline{u}^r]$.

The form $F(\underline{u}^0, \dots, \underline{u}^r)$ is a *Chow form* of $V_{\mathbb{C}}$. The *degree* of $V_{\mathbb{C}}$ is defined to be the degree d of F as a polynomial in $\mathbb{C}[\underline{u}^1, \dots, \underline{u}^r][\underline{u}^0]$. In fact F is *multihomogeneous* of *multidegree* d , which means it is homogeneous of degree d as a form in $\mathbb{C}[\underline{u}^0, \dots, \underline{u}^{j-1}, \underline{u}^{j+1}, \dots, \underline{u}^r][\underline{u}^j]$ for all $j = 0, \dots, r$ [5, p. 694].

It follows directly from the construction that the degree of a variety is the number of points in which a general linear $(n-r)$ -dimensional linear hyperplane meets the variety. This definition of the degree of a variety coincides with the definition of the degree by the Hilbert polynomial [11, p. 48].

The Chow form of a variety is uniquely determined up to multiplication by elements in \mathbb{C}^* . Viewed as an element of $\overline{\mathbb{C}(\underline{u}^1, \dots, \underline{u}^r)}[\underline{u}^0]$ the Chow form F of a

variety V splits into linear factors $\langle \underline{u}^0, \underline{p}^k \rangle$, where every \underline{p}^k is a general point of V . Therefore two varieties are equal iff their Chow forms coincide up to a factor in \mathbb{C}^* .

In the following some properties of Chow forms are described, especially their relations to resultants.

Lemma 1.8. *The Chow form $F(\underline{u}^0, \dots, \underline{u}^r)$ of a r -dimensional variety $V_{\mathbb{C}}$ is irreducible in $\mathbb{C}[\underline{u}^0, \dots, \underline{u}^r]$.*

Proof. [5, p. 693]. □

Reminder 1.9. Let S be a subring of Ω , and let $f_1, \dots, f_k \in S[x_0, \dots, x_n]$ be homogeneous polynomials. The *resultant of the homogeneous equations*

$$f_1 = \dots = f_k = 0$$

is an ideal $J(f) \subset S$, which is generated by determinants of matrices, where one fixed row of such a matrix consists of coefficients of one f_j [22, § 130]. Therefore there is a set of elements in the polynomial ring in the coefficients of f_1, \dots, f_k over \mathbb{Z} , whose image in S generates the ideal $J(f)$. Because the ideal $J(f)$ is generated by determinants, it is homogeneous in the coefficients of each f_j , $j = 1, \dots, k$.

We also say $J(f)$ is the resultant of the homogeneous polynomials f_1, \dots, f_k . The ideal $J(f)$ has the following property [22, §130]:

$$\begin{aligned} f_1 = \dots = f_k = 0 \text{ has a solution in } \mathbb{P}^n(\Omega) \\ \iff J(f) \text{ is the zero ideal in } S. \end{aligned}$$

Let $I(f) \subset S[x_0, \dots, x_n]$ denote the ideal which is generated by f_1, \dots, f_k . The zero set of $I(f)$ in $\mathbb{P}^n(\Omega)$ is empty iff $J(f) \neq 0$, but the ideal $J(f)$ depends on the choice of generators of $I(f)$. The radical of $J(f)$ only depends on the radical of $I(f)$:

To see this let $g_1, \dots, g_l \in S[x_0, \dots, x_n]$ be homogeneous polynomials and $I(g)$ be the ideal which is generated by g_1, \dots, g_l . Furthermore, let $J(g)$ be the resultant of $g_1 = \dots = g_l = 0$. Suppose that $\sqrt{I(g)} = \sqrt{I(f)}$. Then $g_1 = \dots = g_l = 0$ has a solution in $\mathbb{P}^n(\Omega)$ iff $f_1 = \dots = f_k = 0$ has a solution in $\mathbb{P}^n(\Omega)$. Hence $J(g) = 0$ if and only if $J(f) = 0$, which in turn implies that $\sqrt{J(g)} = \sqrt{J(f)}$. Therefore we denote by $\sqrt{J(f)}$ the *resultant of the ideal $I(f)$* .

The Chow form of a variety $V_{\mathbb{C}}$ arises from the intersection with H_{n-r} . If one cuts this 0-dimensional set with another sufficiently general hyperplane, the intersection is empty. Therefore one gets a set of homogeneous polynomials, whose resultant is not the zero ideal. In fact the radical of this resultant is generated by the Chow form:

Theorem 1.10. *Let $V_{\mathbb{C}} \subset \mathbb{P}^n(\Omega)$ be a r -dimensional variety, and let f_1, \dots, f_k be homogeneous generators of $I_{\mathbb{C}}(V) \subset \mathbb{C}[x_0, \dots, x_n]$. The Chow form $F(\underline{u}^0, \dots, \underline{u}^r)$ generates the radical $\sqrt{R} \subset \mathbb{C}[\underline{u}^0, \dots, \underline{u}^r]$ of the resultant R of the following equations:*

$$\begin{aligned} (1) \quad & f_1(x_0, \dots, x_n) = \dots = f_k(x_0, \dots, x_n) = 0, \\ & u_0^0 x_0 + \dots + u_n^0 x_n = 0, \\ & \mathcal{U}(n, r) \cdot \underline{x} = 0 \end{aligned}$$

viewed as polynomials in $\mathbb{C}[\underline{u}^0, \dots, \underline{u}^r][x_0, \dots, x_n]$.

Proof. The first step is to show that \sqrt{R} is a prime ideal in $\mathbb{C}[\underline{u}^0, \dots, \underline{u}^r]$: The proof of this statement uses the same techniques as the proof of Lemma 1.5 in [23, p. 140]. Let

$$A := \{(\underline{p}, \underline{y}^0, \dots, \underline{y}^r) \in \mathbb{P}^n(\Omega) \times (\mathbb{P}^n(\Omega))^{r+1} \mid \underline{p} \in V_{\mathbb{C}}, \langle \underline{p}, \underline{y}^0 \rangle = \dots = \langle \underline{p}, \underline{y}^r \rangle = 0\}.$$

The vanishing ideal of A is generated by the polynomials in (1). The projection pr_1 onto the first factor is again the set $V_{\mathbb{C}} = pr_1(A)$ and the projection pr_2 of A onto the second factor $(\mathbb{P}^n(\Omega))^{r+1}$ gives us the zero set of R .

Now we can apply [21, I.6.3 Th. 8] to pr_1 and obtain that A is irreducible. Hence $pr_2(A)$ is irreducible, which proves that \sqrt{R} is prime.

Now we can prove the theorem: The Chow form F of $V_{\mathbb{C}}$ is $F(\underline{u}^0, \dots, \underline{u}^r) = c \prod_{i=1}^d \langle \underline{p}^i, \underline{u}^0 \rangle$, where $\{\underline{p}^1, \dots, \underline{p}^d\}$ is the intersection of the zero set of $\mathcal{U}(n, r) \cdot \underline{x}$ and $V_{\mathbb{C}}$, $c \in \mathbb{C}[\underline{u}^1, \dots, \underline{u}^r]$. Denote by $Z \subset \overline{\mathbb{C}(\underline{u}^1, \dots, \underline{u}^r)}$ the splitting field of the p_j^i , $i = 1, \dots, d$, $j = 0, \dots, n$.

Now view the polynomials in (1) as polynomials in $Z[\underline{u}^0][x_0, \dots, x_n]$ and determine the resultant $R \cdot Z[\underline{u}^0]$. The zero set of $R \cdot Z[\underline{u}^0]$ equals the zero set of $\prod_{i=1}^d \langle \underline{p}^i, \underline{u}^0 \rangle$. This means that the radical of the ideal which is generated by R in $Z[\underline{u}^0]$ is the ideal which is generated by F :

$$(F)_{Z[\underline{u}^0]} = \sqrt{R \cdot Z[\underline{u}^0]} \quad \text{in } Z[\underline{u}^0].$$

Cutting down with $\mathbb{C}[\underline{u}^0, \dots, \underline{u}^r]$ we first obtain

$$(F)_{\mathbb{C}[\underline{u}^0, \dots, \underline{u}^r]} \subset (F)_{Z[\underline{u}^0]} \cap \mathbb{C}[\underline{u}^0, \dots, \underline{u}^r].$$

Now let $h \in Z[\underline{u}^0]$ such that $F \cdot h \in (F)_{Z[\underline{u}^0]} \cap \mathbb{C}[\underline{u}^0, \dots, \underline{u}^r]$. And let σ be an element of the Galois group of Z over $\mathbb{C}(\underline{u}^1, \dots, \underline{u}^r)$. Then $F \cdot h = \sigma(F \cdot h) = F \cdot \sigma(h)$ and this means that $h \in \mathbb{C}(\underline{u}^1, \dots, \underline{u}^r)[\underline{u}^0]$, because Z is a Galois extension. Therefore $h = \frac{h_1}{h_2}$, $h_1 \in \mathbb{C}[\underline{u}^0, \dots, \underline{u}^r]$, $h_2 \in \mathbb{C}[\underline{u}^1, \dots, \underline{u}^r]$, and the common divisor of h_1 and h_2 is 1. Now h_2 must be a divisor of F , and since F is irreducible and \underline{u}^0 does not appear in h_2 , then $h_2 \in \mathbb{C}$. Therefore

$$(F)_{Z[\underline{u}^0]} \cap \mathbb{C}[\underline{u}^0, \dots, \underline{u}^r] = (F)_{\mathbb{C}[\underline{u}^0, \dots, \underline{u}^r]}.$$

Now it remains to show that \sqrt{R} equals $\sqrt{R \cdot Z[\underline{u}^0]} \cap \mathbb{C}[\underline{u}^0, \dots, \underline{u}^r]$. Obviously $\sqrt{R} \subset \sqrt{R \cdot Z[\underline{u}^0]} \cap \mathbb{C}[\underline{u}^0, \dots, \underline{u}^r] = (F)_{Z[\underline{u}^0]} \cap \mathbb{C}[\underline{u}^0, \dots, \underline{u}^r] = (F)_{\mathbb{C}[\underline{u}^0, \dots, \underline{u}^r]}$. For the inverse inclusion we have to show that $F \in \sqrt{R}$.

We have already seen that $F \in \sqrt{R \cdot Z[\underline{u}^0]} \cap \mathbb{C}[\underline{u}^0, \dots, \underline{u}^r]$. This means that

$$F^k = \sum_{i=1}^l g_i z_i, \quad g_i \in R, z_i \in Z[\underline{u}^0], k \in \mathbb{N}^*.$$

Z is a Galois extension of $\mathbb{C}(\underline{u}^1, \dots, \underline{u}^r)$ and the Galois group $G(Z : \mathbb{C}(\underline{u}^1, \dots, \underline{u}^r))$ is finite. Furthermore, F and g_i , $i = 1, \dots, l$ are elements in $\mathbb{C}[\underline{u}^0, \dots, \underline{u}^r]$ and therefore they stay invariant under the action of the group $G(Z : \mathbb{C}(\underline{u}^1, \dots, \underline{u}^r))$.

Now one obtains:

$$\begin{aligned} |G(Z : \mathbb{C}(\underline{u}^1, \dots, \underline{u}^r))| \cdot F^k &= \sum_{\sigma \in G(Z : \mathbb{C}(\underline{u}^1, \dots, \underline{u}^r))} \sigma\left(\sum_{i=1}^l g_i z_i\right) \\ &= \sum_{i=1}^l g_i \underbrace{\sum_{\sigma \in G(Z : \mathbb{C}(\underline{u}^1, \dots, \underline{u}^r))} \sigma(z_i)}_{:= \tilde{z}_i} \end{aligned}$$

and the \tilde{z}_i are invariant under Galois action, hence elements in $\mathbb{C}(\underline{u}^1, \dots, \underline{u}^r)[\underline{u}^0]$. But then there are $s_1, \dots, s_l \in \mathbb{C}[\underline{u}^0, \dots, \underline{u}^r]$ and $t \in \mathbb{C}[\underline{u}^1, \dots, \underline{u}^r]$ such that

$$t \cdot F^k = \sum_{i=1}^l g_i \cdot s_i.$$

Because g_i , $i = 1, \dots, l$, are elements of R we have $tF^k \in R$. But that means $tF \in \sqrt{R}$. We have proved that \sqrt{R} is a prime ideal, therefore $t \in \sqrt{R}$ or $F \in \sqrt{R}$. Suppose $t \in \sqrt{R}$. We know that the polynomials (1) get common zeros $\underline{p}^1, \dots, \underline{p}^d$ if we set $\underline{u}^0 = (0, \dots, 0)$. That means the resultant R is mapped onto the zero ideal if \underline{u}^0 is mapped to $(0, \dots, 0)$. But $t \in \mathbb{C}[\underline{u}^1, \dots, \underline{u}^r]$ and therefore it is not mapped onto zero and hence not an element of \sqrt{R} . That finishes the proof of the theorem. \square

If the multihomogeneous monomials in $\mathbb{C}[\underline{u}^0, \dots, \underline{u}^r]$ are lexicographically ordered, then the coefficients of a Chow form F of degree d define a point in $\mathbb{P}^N(\mathbb{C})$, $N = \binom{n+d}{d}^{r+1} - 1$, because Chow forms are only determined up to a factor in \mathbb{C}^* . In the following the coefficients of a Chow form are also denoted *Chow coordinates*. Therefore the space of r -dimensional varieties in $\mathbb{P}^n(\mathbb{C})$ of degree d constitutes a subspace of $\mathbb{P}^N(\mathbb{C})$ and hence it is canonically topologized.

Definition 1.11. The *Chow variety* $\mathcal{C}_{r,\bullet}(\mathbb{P}^n(\mathbb{C}))$ as a set is the free abelian semi-group generated by the r -dimensional varieties $V \subset \mathbb{P}^n(\Omega)$ over \mathbb{C} . Let V_i be r -dimensional varieties and $F_i(\underline{u}^0, \dots, \underline{u}^r)$ the Chow forms of V_i . The *Chow form* of the cycle $\sum_{i=1}^m n_i V_i$ is defined to be

$$F(\underline{u}^0, \dots, \underline{u}^r) = \prod_{i=1}^m F_i(\underline{u}^0, \dots, \underline{u}^r)^{n_i}.$$

The *degree* of $\sum_{i=1}^m n_i V_i$ is again the multidegree of the multihomogeneous form $F(\underline{u}^0, \dots, \underline{u}^r)$ and hence equals $\sum_{i=1}^m n_i \cdot \deg V_i$.

The degree of cycles defines a natural filtration of the Chow variety $\mathcal{C}_{r,\bullet}(\mathbb{P}^n(\mathbb{C}))$.

Definition 1.12. The *Chow variety* $\mathcal{C}_{r,d}(\mathbb{P}^n(\mathbb{C}))$ is defined as follows:

$$\mathcal{C}_{r,d}(\mathbb{P}^n(\mathbb{C})) := \{V \in \mathcal{C}_{r,\bullet}(\mathbb{P}^n(\mathbb{C})) \mid \deg V = d\}.$$

The Chow variety $\mathcal{C}_{r,d}(\mathbb{P}^n(\mathbb{C}))$ again can be topologized as a subset of $\mathbb{P}^N(\mathbb{C})$, by the definition of Chow forms.

Example 1.13. There are three natural homeomorphisms:

- (i) $\mathcal{C}_{0,d}(\mathbb{P}^n(\mathbb{C}))$ is homeomorphic to the d -fold symmetric product $SP^d(\mathbb{P}^n(\mathbb{C}))$ of $\mathbb{P}^n(\mathbb{C})$.
- (ii) $\mathcal{C}_{r,1}(\mathbb{P}^n(\mathbb{C}))$ coincides with \mathcal{G}_{r+1}^{n-r} the Grassmannian of $(r+1)$ -dimensional linear subspaces of \mathbb{C}^{n+1} . The Chow coordinates of a linear subspace coincide with its Plücker coordinates.
- (iii) $\mathcal{C}_{n-1,d}(\mathbb{P}^n(\mathbb{C})) \cong \mathbb{P}^N(\mathbb{C})$, $N = \binom{n+d}{d} - 1$, because the Chow forms of cycles of codimension 1 and degree d coincide naturally with homogeneous forms of degree d in $\mathbb{C}[x_0, \dots, x_n]$.

$\mathcal{C}_{r,d}(\mathbb{P}^n(\mathbb{C}))$ is an algebraic set [5, Thm. 2] and hence compact. It consists of algebraic cycles whose Chow forms are multihomogeneous forms of multidegree d . The following theorem of Chow and Van der Waerden demonstrates which multihomogeneous forms of multidegree d are Chow forms of algebraic cycles.

Theorem 1.14. *Let $F \in \mathbb{C}[\underline{u}^0, \dots, \underline{u}^r]$ be multihomogeneous of multidegree d . F is a Chow form of an algebraic cycle in $\mathcal{C}_{r,d}(\mathbb{P}^n(\mathbb{C}))$ iff the following conditions hold:*

There are d points $\underline{p}^i \in \overline{\mathbb{C}(\underline{u}^1, \dots, \underline{u}^r)}^{n+1} - \{0\}$ and $c \in \mathbb{C}(\underline{u}^1, \dots, \underline{u}^r)$ satisfying:

- (i) $F(\underline{u}^0, \dots, \underline{u}^r) = c \prod_{i=1}^d \langle \underline{p}^i, \underline{u}^0 \rangle$, in $\overline{\mathbb{C}(\underline{u}^1, \dots, \underline{u}^r)}[\underline{u}^0]$.
- (ii) $L_i(\underline{u}^j) := \langle \underline{p}^i, \underline{u}^j \rangle = 0$ for $i = 1, \dots, d$, $j = 1, \dots, r$.
- (iii) *For every $i \in \{1, \dots, d\}$ one has: Let S^j be skew symmetric $(n+1) \times (n+1)$ -matrices, $j = 1, \dots, r$, whose entries $s_{k,l}^j \in \Omega$ are algebraically independent over $\mathbb{C}(\underline{p}^i)$, $k, l = 0, \dots, n$. In this situation $L_i(\underline{u}^0)$ divides the form $F(\underline{u}^0, S^1 \underline{p}^i, \dots, S^r \underline{p}^i)$ in $\mathbb{C}(\underline{p}^i)[s_{0,0}^1, \dots, s_{n,n}^r][\underline{u}^0]$.*

Proof. [5, Thm. 1]. □

Remark 1.15. Let $\underline{p} \in \Omega^{n+1} - \{0\}$ and $L(\underline{x}) := \langle \underline{p}, \underline{x} \rangle$. Then it is easy to see that $\underline{v} \in \Omega^{n+1}$ is an element of the zero set of L iff there is a skew symmetric matrix S satisfying $\underline{v} = S\underline{p}$. Therefore the third condition can be reformulated as: Let $v_k^j \in \Omega$, $k = 0, \dots, n$, $j = 1, \dots, r$ and $i \in \{1, \dots, d\}$, such that

$$L_i(\underline{v}^j) = 0 \quad \forall j = 1, \dots, r.$$

Then $L_i(\underline{u}^0)$ divides $F(\underline{u}^0, \underline{v}^1, \dots, \underline{v}^r)$ in $\mathbb{C}(\underline{p}^i)[\underline{v}^1, \dots, \underline{v}^r][\underline{u}^0]$.

Let $V_{\mathbb{C}} = \sum_{i=1}^m n_i V_i \in \mathcal{C}_{r,d}(\mathbb{P}^n(\mathbb{C}))$. Then the algebraic set $\text{supp}(V_{\mathbb{C}}) := \bigcup_{i=1}^m V_i$ is called the *support* of $V_{\mathbb{C}}$. From a given Chow form F the support of the cycle, whose Chow form is F , can be determined:

Lemma 1.16. *Let $V_{\mathbb{C}} \in \mathcal{C}_{r,d}(\mathbb{P}^n(\mathbb{C}))$ and $F(\underline{u}^0, \dots, \underline{u}^r)$ be the Chow form of $V_{\mathbb{C}}$, and let S^i be skew symmetric $(n+1) \times (n+1)$ -matrices whose entries $s_{j,k}^i$ are algebraically independent over \mathbb{C} . The coefficients of*

$$F(S^0 \underline{x}, \dots, S^r \underline{x}) \in \mathbb{C}[x_0, \dots, x_n][s_{0,1}^0, \dots, s_{n-1,n}^r],$$

are denoted $f_l(x_0, \dots, x_n) \in \mathbb{C}[x_0, \dots, x_n]$. Then the zero set of the f_l is $\text{supp}(V_{\mathbb{C}})$.

Proof. [5, Thm. 3]. □

From Theorem 1.10 one concludes directly the following corollary, which makes the connection to Šafarevič's [21, p. 65] definition of Chow varieties transparent.

The equivalence of the definitions of Chow/Van der Waerden and Šafarevič is implicitly shown in the proof of 1.10.

Corollary 1.17. *Let $F(\underline{u}^0, \dots, \underline{u}^r)$ be the Chow form of $V \in \mathcal{C}_{r,d}(\mathbb{P}^n(\mathbb{C}))$, then the following statements are equivalent.*

- (i) \underline{p} is contained in $\text{supp}(V)$
- (ii) $F(\underline{v}^0, \dots, \underline{v}^r) = 0$ for all $\underline{v}^0, \dots, \underline{v}^r \in \Omega^{n+1}$ such that $\langle \underline{v}^0, \underline{p} \rangle = \dots = \langle \underline{v}^r, \underline{p} \rangle = 0$.

2. STABILIZATIONS AND PAIRINGS OF CHOW VARIETIES

The additive structure on the free abelian semi group $\mathcal{C}_{r,\bullet}(\mathbb{P}^n(\mathbb{C}))$ is given by formal addition of cycles, i. e. multiplication of the Chow forms of cycles. Therefore $\mathcal{C}_{r,\bullet}(\mathbb{P}^n(\mathbb{C}))$ is a topological semi group. In particular there is a continuous pairing

$$+ : \mathcal{C}_{r,d_1}(\mathbb{P}^n(\mathbb{C})) \times \mathcal{C}_{r,d_2}(\mathbb{P}^n(\mathbb{C})) \longrightarrow \mathcal{C}_{r,d_1+d_2}(\mathbb{P}^n(\mathbb{C})) .$$

There is a second pairing on Chow varieties: Let $V \subset \mathbb{P}^n(\Omega)$ and $W \subset \mathbb{P}^m(\Omega)$ be varieties and $I_{\mathbb{C}}(V) \subset \mathbb{C}[x_0, \dots, x_n]$ and $I_{\mathbb{C}}(W) \subset \mathbb{C}[x_{n+1}, \dots, x_{n+m+1}]$ be the vanishing ideals of V and W . Define $V \# W \subset \mathbb{P}^{n+m+1}(\Omega)$ to be the zero set of the ideal $J \subset \mathbb{C}[x_0, \dots, x_{n+m+1}]$ which is generated by the elements of $I_{\mathbb{C}}(V)$ and $I_{\mathbb{C}}(W)$ viewed as elements in $\mathbb{C}[x_0, \dots, x_{n+m+1}]$.

Definition 2.1. The algebraic set $V \# W$ is called the *complex join* of V and W and we call the join with a point $\mathbb{P}^0(\mathbb{C}) \# V$ the *complex suspension* of V .

There are other equivalent ways to define the join of varieties. For example let $C(V)$ and $C(W)$ be the affine cones of V and W , i. e. $C(V)$ is the zero set of $I_{\mathbb{C}}(V)$ in \mathbb{C}^{n+1} . Then the join of V and W is uniquely determined by

$$C(V \# W) := C(V) \times C(W) .$$

From the second definition we deduce that the dimension of $V \# W$ is $\dim V + \dim W + 1$, and one also obtains $V \# W = (V \# \mathbb{P}^m(\mathbb{C})) \cap (\mathbb{P}^n(\mathbb{C}) \# W)$.

The join of two varieties is again a variety, and one wants to calculate the degree of the join of two varieties:

The varieties $V \# \mathbb{P}^m(\mathbb{C})$ and $\mathbb{P}^n(\mathbb{C}) \# W$ intersect transversally, hence by Bezout [19, Chap. 5, Thm. 5.16]:

$$\begin{aligned} \deg V \# W &= \deg(V \# \mathbb{P}^m(\mathbb{C})) \cdot \deg(\mathbb{P}^n(\mathbb{C}) \# W) \\ &= \deg V \cdot \deg W . \end{aligned}$$

By linear extension to algebraic cycles the join defines an associative pairing on Chow varieties:

$$\mathcal{C}_{r_1,d_1}(\mathbb{P}^{n_1}(\mathbb{C})) \times \mathcal{C}_{r_2,d_2}(\mathbb{P}^{n_2}(\mathbb{C})) \longrightarrow \mathcal{C}_{r_1+r_2+1,d_1d_2}(\mathbb{P}^{n_1+n_2+1}(\mathbb{C})).$$

Proposition 2.2. *The complex suspension $\mathbb{Z} : \mathcal{C}_{r-1,d}(\mathbb{P}^{n-1}(\mathbb{C})) \rightarrow \mathcal{C}_{r,d}(\mathbb{P}^n(\mathbb{C}))$ is algebraic and hence continuous.*

Proof. Let $F(\underline{u}^1, \dots, \underline{u}^r)$ be the Chow form of $V = \sum_{i=1}^k n_i V_i \in \mathcal{C}_{r-1,d}(\mathbb{P}^{n-1}(\mathbb{C}))$.

Let $\underline{u}^i = (u_1^i, \dots, u_n^i)$, $i = 1, \dots, r$, and let $w_k^j \in \Omega$ be algebraically independent

over $\mathbb{C}(\underline{u}^1, \dots, \underline{u}^r)$, $k = 0, \dots, n$, $j = 0, \dots, r$.

For suitable $\underline{p}^l \in \left(\overline{\mathbb{C}(\underline{u}^2, \dots, \underline{u}^r)}\right)^n$, $l = 1, \dots, d$, the Chow form of V splits:

$$\begin{aligned} F_V(\underline{u}^1, \dots, \underline{u}^r) &= \prod_{l=1}^d \sum_{k=1}^n p_k^l u_k^1 \in \overline{\mathbb{C}(\underline{u}^2, \dots, \underline{u}^r)}[\underline{u}^1] \\ &= \sum_{1 \leq i_1 \leq \dots \leq i_d \leq n} u_{i_1}^1 \cdots u_{i_d}^1 \underbrace{\sum_{\sigma \in \Sigma(i_1, \dots, i_d)} p_{\sigma(i_1)}^1 \cdots p_{\sigma(i_d)}^d}_{=: G_{i_1, \dots, i_d}(\underline{u}^2, \dots, \underline{u}^r)}, \end{aligned}$$

such that $G_{i_1, \dots, i_d} \in \mathbb{C}[\underline{u}^2, \dots, \underline{u}^r]$. The group $\Sigma(i_1, \dots, i_d)$ is that subgroup of Σ_n which only permutes the numbers $i_1, \dots, i_d \in \{1, \dots, n\}$. Each \underline{p}^l is contained in one of the V_j and is a solution of the following equations:

$$\langle \underline{u}^i, \underline{x} \rangle = 0, \quad i = 2, \dots, r.$$

On the other hand the Chow form of $\mathbb{A}^1 V$ is obtained from points \underline{q}^l of one of the $\mathbb{A}^1 V_j$ which are solutions of the equation:

$$\begin{pmatrix} v_0^1 & v_1^1 & \cdots & v_n^1 \\ \vdots & \vdots & & \vdots \\ v_0^r & v_1^r & \cdots & v_n^r \end{pmatrix} \begin{pmatrix} x_0 \\ \vdots \\ x_n \end{pmatrix} = 0.$$

This equation is equivalent to

$$\underbrace{\begin{pmatrix} v_0^1 & v_1^1 & \cdots & v_n^1 \\ 0 & v_1^2 v_0^1 - v_1^1 v_0^2 & \cdots & v_n^2 v_0^1 - v_n^1 v_0^2 \\ \vdots & \vdots & & \vdots \\ 0 & v_1^r v_0^1 - v_1^1 v_0^r & \cdots & v_n^r v_0^1 - v_n^1 v_0^r \end{pmatrix}}_{=: A} \begin{pmatrix} x_0 \\ x_1 \\ \vdots \\ x_n \end{pmatrix} = 0,$$

and the entries of the matrix A are algebraically independent over $\mathbb{C}(\underline{u}^1, \dots, \underline{u}^r)$.

Step 1: Observe the following injective ring homomorphism:

$$\begin{array}{ccc} \tilde{\varphi}: \mathbb{C}[\underline{u}^2, \dots, \underline{u}^r, \underline{v}^0, \underline{v}^1] & \longrightarrow & \mathbb{C}[\underline{v}^0, \dots, \underline{v}^r], \\ u_k^i & \longmapsto & v_k^i v_0^1 - v_k^1 v_0^i, \quad i = 2, \dots, r, k = 1, \dots, n, \\ c & \longmapsto & c \quad \forall c \in \mathbb{C}[\underline{v}^0, \underline{v}^1]. \end{array}$$

It is extendable to an automorphism φ of Ω . Then one obtains:

$$\varphi^{-1}(A) = \begin{pmatrix} v_0^1 & v_1^1 & \cdots & v_n^1 \\ 0 & u_1^2 & \cdots & u_n^2 \\ \vdots & \vdots & & \vdots \\ 0 & u_1^r & \cdots & u_n^r \end{pmatrix}.$$

Now one can calculate the points \underline{q}^l from which the Chow form of $\mathbb{A}^1 V$ arises:

$$\begin{aligned} \underline{q}^i &= (t_i, \varphi(p_1^i), \dots, \varphi(p_n^i)), \quad i = 1, \dots, d, \\ t_i &= -\frac{v_1^1 \varphi(p_1^i) + \cdots + v_n^1 \varphi(p_n^i)}{v_0^1}. \end{aligned}$$

With this one determines the Chow form as in Construction 1.7. We start with the form G using the notations of 1.7 multiplied by $(v_0^1)^d$. Let I_k be the index set

$I_k := \{(i_1, \dots, i_d) \in \mathbb{N}^d \mid 1 \leq i_1 \leq \dots \leq i_k \leq n, 1 \leq i_{k+1} \leq \dots \leq i_d \leq n\}$ for $0 \leq k \leq d$.

$$\begin{aligned}
 G(\underline{v}^0, \dots, \underline{v}^r) &:= (v_0^1)^d \prod_{i=1}^d \sum_{j=0}^n q_j^i v_j^0 = \prod_{i=1}^d \left(v_0^1 t_i v_0^0 + \sum_{j=1}^n v_0^1 \varphi(p_j^i) v_j^0 \right) \\
 &= \prod_{i=1}^d \left(-(v_1^1 \varphi(p_1^i) + \dots + v_n^1 \varphi(p_n^i)) v_0^0 + \sum_{j=1}^n v_0^1 \varphi(p_j^i) v_j^0 \right) \\
 &= \prod_{i=1}^d \left(-v_0^0 \sum_{j=1}^n v_j^1 \varphi(p_j^i) + v_0^1 \sum_{j=1}^n \varphi(p_j^i) v_j^0 \right) \\
 &= \sum_{k=0}^d \sum_{I_k} (-v_0^0)^k v_{i_1}^1 \dots v_{i_k}^1 (v_0^1)^{d-k} v_{i_{k+1}}^0 \dots v_{i_d}^0 \sum_{\sigma \in \Sigma(i_1, \dots, i_d)} \varphi(p_{\sigma(i_1)}^1) \dots \varphi(p_{\sigma(i_d)}^d) \\
 &= \sum_{k=0}^d \sum_{I_k} (-v_0^0)^k v_{i_1}^1 \dots v_{i_k}^1 (v_0^1)^{d-k} v_{i_{k+1}}^0 \dots v_{i_d}^0 \cdot \varphi \left(\sum_{\sigma \in \Sigma(i_1, \dots, i_d)} p_{\sigma(i_1)}^1 \dots p_{\sigma(i_d)}^d \right) \\
 &= \sum_{k=0}^d \sum_{I_k} (-v_0^0)^k v_{i_1}^1 \dots v_{i_k}^1 (v_0^1)^{d-k} v_{i_{k+1}}^0 \dots v_{i_d}^0 \cdot \varphi(G_{i_1, \dots, i_d}(\underline{u}^2, \dots, \underline{u}^r)) .
 \end{aligned}$$

This calculation shows that the coefficients of the form G are polynomials in the coefficients of F_V . These polynomials depend only on the dimension, the codimension and the degree of V .

Step 2: The form $G \in \mathbb{C}[\underline{v}^0, \dots, \underline{v}^r]$ is homogeneous of degree d in $\underline{v}^0, \underline{v}^2, \dots, \underline{v}^r$ and homogeneous of degree rd in \underline{v}^1 . Therefore G is not the Chow form of $\mathbb{Z}V$, but as in Construction 1.7 there is a $T \in \mathbb{C}(\underline{v}^1, \dots, \underline{v}^r)$ such that

$$G(\underline{v}^0, \dots, \underline{v}^r) = F_{\mathbb{Z}V}(\underline{v}^0, \dots, \underline{v}^r) T ,$$

where $F_{\mathbb{Z}V}$ denotes the Chow form of $\mathbb{Z}V$. Let $T_1 \in \mathbb{C}[\underline{v}^1, \dots, \underline{v}^r]$ be the common denominator of T . Then T_1 has to be a divisor of $F_{\mathbb{Z}V}$ in $\mathbb{C}[\underline{v}^0, \dots, \underline{v}^r]$. But this can only happen if $T_1 \in \mathbb{C}^*$, because the irreducible factors of $F_{\mathbb{Z}V}$ are polynomials of positive degree in \underline{v}^0 . This proves that $T \in \mathbb{C}[\underline{v}^1]$ and T is homogeneous of degree $(r-1)d$.

For the case $r = 1$ the theorem is proved. For the case $r > 1$ one has to show that $T(\underline{v}^1) = (v_0^1)^{(r-1)d}$. Without loss of generality assume that V is a variety:

Step 3: Determine the zero set of $T(\underline{v}^1)$ as a polynomial in $\mathbb{C}[\underline{v}^1]$:

Let $\underline{u} = (0, v_1^1, \dots, v_n^1)$ and $\tilde{\underline{u}} = (v_1^1, \dots, v_n^1)$. Then one obtains:

$$\begin{aligned}
 &G(\underline{v}^0, \underline{u}, \underline{v}^2, \dots, \underline{v}^r) \\
 &= \sum_{k=0}^d \sum_{\substack{1 \leq i_1 \leq \dots \leq i_k \leq n \\ 1 \leq i_{k+1} \leq \dots \leq i_d \leq n}} (-v_0^0)^k v_{i_1}^1 \dots v_{i_k}^1 0^{d-k} v_{i_{k+1}}^0 \dots v_{i_d}^0 \cdot \varphi(G_{i_1, \dots, i_d}(\underline{u}^2, \dots, \underline{u}^r)) \\
 &= \sum_{1 \leq i_1 \leq \dots \leq i_d \leq n} (-v_0^0)^d v_{i_1}^1 \dots v_{i_d}^1 \cdot G_{i_1, \dots, i_d}(v_0^2 \cdot \tilde{\underline{u}}, v_0^3 \cdot \tilde{\underline{u}}, \dots, v_0^r \cdot \tilde{\underline{u}}) \\
 &= (-v_0^0)^d \sum_{1 \leq i_1 \leq \dots \leq i_d \leq n} v_{i_1}^1 \dots v_{i_d}^1 \cdot (v_0^2)^d \dots (v_0^r)^d \cdot G_{i_1, \dots, i_d}(\tilde{\underline{u}}, \dots, \tilde{\underline{u}}) \\
 &= (-v_0^0)^d \cdot (v_0^2)^d \dots (v_0^r)^d \sum_{1 \leq i_1 \leq \dots \leq i_d \leq n} v_{i_1}^1 \dots v_{i_d}^1 \cdot G_{i_1, \dots, i_d}(\tilde{\underline{u}}, \dots, \tilde{\underline{u}}) \\
 &= (-v_0^0)^d \cdot (v_0^2)^d \dots (v_0^r)^d \cdot F_V(\tilde{\underline{u}}, \dots, \tilde{\underline{u}}) .
 \end{aligned}$$

But $F_V(\underline{w}, \dots, \underline{w})$ must be zero, because the dimension of V is greater than 0, and $F_V(\underline{w}, \dots, \underline{w})$ is a divisor of a resultant which has to be 0. It is easy to show that the Chow form $F_{\mathbb{P}^n}(\underline{v}^0, \underline{w}, \underline{v}^2, \dots, \underline{v}^r) \neq 0$ using the property that $F_{\mathbb{P}^n}$ generates the radical of a resultant. Therefore $T(0, v_1^1, \dots, v_n^1) = 0$ for all $v_1^1, \dots, v_n^1 \in \Omega$ and hence v_0^1 divides T .

Now we claim that v_0^1 is the only irreducible factor of T : Suppose $T(\underline{w}) = 0$ such that $w_0 \neq 0$, $\underline{w} \in \mathbb{P}^n(\mathbb{C})$. We may assume $w_0 = 1$. For the Chow form of \mathbb{P}^n one has

$$F_{\mathbb{P}^n}(\underline{v}^0, \underline{w}, \underline{v}^2, \dots, \underline{v}^r) = 0 \text{ iff } \langle \underline{p}, \underline{w} \rangle = 0 \text{ for all } \underline{p} \in \mathbb{P}^n \cap H_{n-(r-1)}.$$

But this would mean $\langle \underline{x}, \underline{w} \rangle \in \mathbb{C}[\underline{x}]$ is an element of $I_{\mathbb{C}}(\mathbb{P}^n)$ because $\mathbb{P}^n \cap H_{n-(r-1)}$ contains general points of \mathbb{P}^n . This implies that $w_0 = 0$, because $\langle \underline{x}, \underline{w} \rangle$ is irreducible and $I_{\mathbb{C}}(\mathbb{P}^n)$ is generated by elements in $\mathbb{C}[x_1, \dots, x_n]$. This is a contradiction to the assumption $w_0 = 1$.

On the other hand we have $G(\underline{v}^0, \underline{w}, \underline{v}^2, \dots, \underline{v}^r) = 0$, because $T(\underline{w}) = 0$. That means there is a point $\underline{p} \in \mathbb{P}^n$ satisfying the following equation:

$$\begin{pmatrix} v_0^0 & v_1^0 & \dots & v_n^0 \\ 1 & w_1^1 & \dots & w_n^1 \\ 0 & v_1^2 - w_1^1 v_0^2 & \dots & v_n^2 - w_n^1 v_0^2 \\ \vdots & \vdots & & \vdots \\ 0 & v_1^r - w_1^1 v_0^r & \dots & v_n^r - w_n^1 v_0^r \end{pmatrix} \begin{pmatrix} x_0 \\ \vdots \\ x_n \end{pmatrix} = 0,$$

which is equivalent to

$$\begin{pmatrix} v_0^0 & v_1^0 & \dots & v_n^0 \\ 1 & w_1^1 & \dots & w_n^1 \\ v_0^2 & v_1^2 & \dots & v_n^2 \\ \vdots & \vdots & & \vdots \\ v_0^r & v_1^r & \dots & v_n^r \end{pmatrix} \begin{pmatrix} x_0 \\ \vdots \\ x_n \end{pmatrix} = 0,$$

because in the case where V is irreducible, the form G equals $\varphi(F_{\mathbb{P}^n})$ up to a factor in $\mathbb{C}[v_0^1]$. But this means that $F_{\mathbb{P}^n}(\underline{v}^0, \underline{w}, \underline{v}^2, \dots, \underline{v}^r) = 0$, which leads to a contradiction as above. \square

As a corollary we obtain that the complex join is an algebraic map:

Theorem 2.3. *The join map*

$$\mathcal{C}_{r_1, d_1}(\mathbb{P}^{n_1}(\mathbb{C})) \times \mathcal{C}_{r_2, d_2}(\mathbb{P}^{n_2}(\mathbb{C})) \longrightarrow \mathcal{C}_{r_1+r_2+1, d_1 d_2}(\mathbb{P}^{n_1+n_2+1}(\mathbb{C}))$$

is algebraic. In particular, it is a continuous map.

Proof. The image of the continuous map $(\# \mathbb{P}^{n_2}, \mathbb{P}^{n_1} \#)$:

$$\mathcal{C}_{r_1, d_1}(\mathbb{P}^{n_1}) \times \mathcal{C}_{r_2, d_2}(\mathbb{P}^{n_2}) \longrightarrow \mathcal{C}_{r_1+n_2+1, d_1}(\mathbb{P}^{n_1+n_2+1}) \times \mathcal{C}_{r_2+n_1+1, d_2}(\mathbb{P}^{n_1+n_2+1})$$

consists only on pairs of cycles whose irreducible components intersect transversally. Therefore the intersection pairing is well defined and analytic on the image of $(\# \mathbb{P}^{n_2}, \mathbb{P}^{n_1} \#)$ [1, Thm. VI.2.10 and p.142]. The Chow varieties are compact and hence the join is algebraic by GAGA (e.g. see [10, Chap. IV.1]). \square

Now we want to define stabilizations of Chow varieties. For this purpose let $L \in \mathcal{C}_{r,1}(\mathbb{P}^n(\mathbb{C}))$ be a fixed linear subspace. Further let inc be the map of Chow varieties, which is induced by the inclusion

$$\begin{aligned} inc : \quad \mathbb{P}^n(\Omega) &\longrightarrow \mathbb{P}^{n+1}(\Omega) \\ (p_0, \dots, p_n) &\longmapsto (p_0, \dots, p_n, 0) \end{aligned} .$$

The following diagram is commutative:

$$\begin{array}{ccccc} & \mathcal{C}_{r,d+1}(\mathbb{P}^n(\mathbb{C})) & \xrightarrow{\quad inc \quad} & \mathcal{C}_{r,d+1}(\mathbb{P}^{n+1}(\mathbb{C})) & \\ & \nearrow +L & & \nearrow +L & \\ \mathcal{C}_{r,d}(\mathbb{P}^n(\mathbb{C})) & \xrightarrow{\quad inc \quad} & \mathcal{C}_{r,d}(\mathbb{P}^{n+1}(\mathbb{C})) & & \\ \downarrow \mathbb{P} & & \downarrow \mathbb{P} & & \downarrow \mathbb{P} \\ & \mathcal{C}_{r+1,d+1}(\mathbb{P}^{n+1}(\mathbb{C})) & \xrightarrow{\quad inc \quad} & \mathcal{C}_{r+1,d+1}(\mathbb{P}^{n+2}(\mathbb{C})) & \\ & \nearrow +\mathbb{P}L & & \nearrow +\mathbb{P}L & \\ \mathcal{C}_{r+1,d}(\mathbb{P}^{n+1}(\mathbb{C})) & \xrightarrow{\quad inc \quad} & \mathcal{C}_{r+1,d}(\mathbb{P}^{n+2}(\mathbb{C})) & & \end{array}$$

The map $+L$ is obviously continuous. The continuity of the map inc follows directly from the definition of Chow forms. Hence we can define several stabilizations of Chow varieties:

Notation 2.4.

$$\mathcal{C}_r(\mathbb{P}^n(\mathbb{C})) := \lim_d \mathcal{C}_{r,d}(\mathbb{P}^n(\mathbb{C})), \quad \mathcal{D}(d) := \lim_{r,n} \mathcal{C}_{r,d}(\mathbb{P}^n(\mathbb{C})),$$

$$\mathcal{C} := \lim_{r,d,n} \mathcal{C}_{r,d}(\mathbb{P}^n(\mathbb{C})), \quad \mathfrak{C} := \coprod_d \mathcal{D}(d).$$

From the examples 1.13 one deduces that $\mathcal{C}_0(\mathbb{P}^n(\mathbb{C}))$ is homeomorphic to the reduced infinite symmetric product $\widetilde{\mathrm{SP}}(\mathbb{P}^n(\mathbb{C}))$, and that $\mathcal{D}(1)$ is homeomorphic to BU , the classifying space of U -bundles.

The spaces $\mathcal{C}_r(\mathbb{P}^n(\mathbb{C}))$ and \mathcal{C} are commutative topological monoids by formal addition of cycles with unit L , and \mathfrak{C} is a commutative topological monoid by formal addition of cycles with unit \emptyset the empty cycle.

Intertwining coordinates, the join pairing extends to a multiplicative pairing on \mathcal{C} , \mathfrak{C} , and $\mathcal{D}(1)$, which on the latter space coincides with the Whitney sum on BU . In the next section the homotopy type of \mathcal{C} will be determined.

3. THE COMPLEX SUSPENSION THEOREM

The following theorem is due to H. B. Lawson [14]. We are providing a more algebraic proof of the theorem, as well as giving a few more details.

The Complex Suspension Theorem 3.1. *The complex suspension*

$$\mathbb{P} : \mathcal{C}_r(\mathbb{P}^n(\mathbb{C})) \longrightarrow \mathcal{C}_{r+1}(\mathbb{P}^{n+1}(\mathbb{C}))$$

is a homotopy equivalence.

As in [14] the theorem will be proved in two steps. First one can show that $\mathbb{Z}\mathcal{C}_{r,d}(\mathbb{P}^n(\mathbb{C}))$ is a strong deformation retract of a space $\mathcal{T}_d \subset \mathcal{C}_{r+1,d}(\mathbb{P}^{n+1}(\mathbb{C}))$. The second step is to show that the inclusion $\mathcal{T} \hookrightarrow \mathcal{C}_{r+1}(\mathbb{P}^{n+1}(\mathbb{C}))$ induces isomorphisms on homotopy groups.

Definition 3.2. Let $\mathbb{P}^n(\mathbb{C}) \subset \mathbb{P}^{n+1}(\mathbb{C})$ denote the hyperplane given by the equation $x_0 = 0$. Then define

$$\mathcal{T}_d := \left\{ \sum_{i=1}^m n_i V_i \in \mathcal{C}_{r+1,d}(\mathbb{P}^{n+1}(\mathbb{C})) \mid V_i \not\subset \mathbb{P}^n(\mathbb{C}), i = 1, \dots, m \right\}.$$

Fixing an element of \mathcal{T}_1 , whose intersection with the hyperplane $\mathbb{P}^n(\mathbb{C})$ is the base point of $\mathcal{C}_{r,d}(\mathbb{P}^n(\mathbb{C}))$, one defines $\mathcal{T} := \lim_{d \rightarrow \infty} \mathcal{T}_d$.

Let $V \in \mathcal{C}_{r+1,d}(\mathbb{P}^{n+1}(\mathbb{C}))$ and F_V be the Chow form of V , and note that V is an element of \mathcal{T}_d iff $F_V(\underline{u}^0, \dots, \underline{u}^r, \underline{e}) \neq 0$, $\underline{e} = (1, 0, \dots, 0)$. This is obviously true for varieties, since F_V generates the radical of a resultant by 1.10. Hence it is also true for cycles. This condition is polynomial in the coefficients of F_V ; hence \mathcal{T}_d is a Zariski open subset of $\mathcal{C}_{r+1,d}(\mathbb{P}^{n+1}(\mathbb{C}))$.

We first construct a homotopy ϕ on Chow varieties, induced by

$$\begin{aligned} \mathbb{P}^{n+1}(\mathbb{C}) \times \mathbb{R}_+^* &\longrightarrow \mathbb{P}^{n+1}(\mathbb{C}), \\ ((p_0, \dots, p_{n+1}), t) &\longmapsto (tp_0, p_1, \dots, p_{n+1}). \end{aligned}$$

The map ϕ retracts \mathcal{T}_d to $\mathbb{Z}\mathcal{C}_{r,d}(\mathbb{P}^n(\mathbb{C}))$. But first one has to check how this induced map is defined on Chow forms.

Lemma 3.3. Let $V \in \mathcal{C}_{r+1,d}(\mathbb{P}^{n+1}(\mathbb{C}))$ and $F_V(\underline{u}^0, \dots, \underline{u}^{r+1})$ its Chow form. Then $\phi(V, t)$ has the Chow form $F_V({}^t\underline{u}^0, \dots, {}^t\underline{u}^{r+1})$, where ${}^t\underline{u}^i = (tu_0^i, u_1^i, \dots, u_{n+1}^i)$.

Proof. Let V be a variety and f_1, \dots, f_k homogeneous generators of $I_{\mathbb{C}}(V)$. Then F_V generates the radical of the resultant of

$$f_1(x_0, \dots, x_{n+1}), \dots, f_k(x_0, \dots, x_{n+1}), \begin{pmatrix} u_0^0 & \cdots & u_{n+1}^0 \\ \vdots & & \vdots \\ u_0^{r+1} & \cdots & u_{n+1}^{r+1} \end{pmatrix} \cdot \begin{pmatrix} x_0 \\ \vdots \\ x_{n+1} \end{pmatrix}.$$

The Chow form of $\phi(V, t)$ is the generator of the radical of the resultant of

$$f_1\left(\frac{1}{t}x_0, \dots, x_{n+1}\right), \dots, f_k\left(\frac{1}{t}x_0, \dots, x_{n+1}\right), \begin{pmatrix} u_0^0 & \cdots & u_{n+1}^0 \\ \vdots & & \vdots \\ u_0^{r+1} & \cdots & u_{n+1}^{r+1} \end{pmatrix} \cdot \begin{pmatrix} x_0 \\ \vdots \\ x_{n+1} \end{pmatrix}$$

which is obviously the same as the radical of the resultant of

$$f_1(x_0, \dots, x_{n+1}), \dots, f_k(x_0, \dots, x_{n+1}), \begin{pmatrix} tu_0^0 & u_1^0 & \cdots & u_{n+1}^0 \\ \vdots & \vdots & & \vdots \\ tu_0^{r+1} & u_1^{r+1} & \cdots & u_{n+1}^{r+1} \end{pmatrix} \begin{pmatrix} x_0 \\ \vdots \\ x_{n+1} \end{pmatrix}.$$

The algebraic independence of the u_i^j remains invariant under multiplication with t ; hence the lemma is proved. \square

Suppose V lies in the image of \mathbb{Z} . Then the vanishing ideal of each irreducible component is generated by polynomials in $\mathbb{C}[x_1, \dots, x_{n+1}]$, which means x_0 does

not appear in these generators. Hence one obtains directly from the proof above that

$$\phi(\cdot, t) |_{\mathbb{Y}\mathcal{C}_{r,d}(\mathbb{P}^n(\mathbb{C}))} = id_{\mathbb{Y}\mathcal{C}_{r,d}(\mathbb{P}^n(\mathbb{C}))} \quad \forall t \in \mathbb{R}_+^*.$$

On the other hand the algebraic independence of the u_i^j remains invariant under the multiplication with t which implies

$$F_V(\underline{u}^0, \dots, \underline{u}^r, \underline{e}) \neq 0 \iff F_V(t\underline{u}^0, \dots, t\underline{u}^r, t\underline{e}) \neq 0.$$

Hence $\phi(\mathcal{T}_d, t) \subset \mathcal{T}_d$. We claim that ϕ extends continuously to $\mathcal{T}_d \times]0, \infty]$. The following lemma is very useful for this purpose:

Lemma 3.4. *Let $V \in \mathcal{C}_{r+1,d}(\mathbb{P}^{n+1}(\mathbb{C}))$ be a variety and F_V be a fixed representative of the Chow form of V . Suppose*

$$F_V(t\underline{u}^0, \dots, t\underline{u}^{r+1}) = t^k F_V(\underline{u}^0, \dots, \underline{u}^{r+1}) \quad \forall t \in \mathbb{C}^*,$$

then V lies in the image of \mathbb{Y} and $k = d$.

Proof. Let $\{\underline{p}^1, \dots, \underline{p}^d\}$ be the intersection of V with a \mathbb{C} -general $(n-r)$ -dimensional linear subspace H_{n-r} . Then the Chow form F_V splits in $\overline{\mathbb{C}(\underline{u}^1, \dots, \underline{u}^{r+1})}[\underline{u}^0]$:

$$\begin{aligned} F_V(t\underline{u}^0, \dots, t\underline{u}^{r+1}) &= t^k F_V(\underline{u}^0, \dots, \underline{u}^{r+1}) \\ &= \frac{t^k}{t^d} \prod_{i=1}^d t \langle \underline{p}^i, \underline{u}^0 \rangle \\ &= \frac{t^k}{t^d} \prod_{i=1}^d \left(p_0^i(tu_0^0) + \sum_{j=1}^{n+1} (tp_j^i)u_j^0 \right) \\ &= \left(\frac{t^k}{t^d} p_0^1(tu_0^0) + \sum_{j=1}^{n+1} \left(\frac{t^k}{t^d} tp_j^1 \right) u_j^0 \right) \prod_{i=2}^d \left(p_0^i(tu_0^0) + \sum_{j=1}^{n+1} (tp_j^i)u_j^0 \right). \end{aligned}$$

$(p_0^i, tp_1^i, \dots, tp_{n+1}^i)$ is a general point of V for every $t \in \mathbb{C}^*$. Hence $(tp_0^i, p_1^i, \dots, p_{n+1}^i)$ is a general point for V , too. One has $p_0^i \neq 0$ unless $k = 0$. Now let $f \in I_{\mathbb{C}}(V)$ and observe that tp_0^i is a zero of $f(x_0, p_1^i, \dots, p_{n+1}^i)$ for every $t \in \mathbb{C}^*$. But this means that the polynomial has infinitely many zeros and hence there must be a set of generators of $I_{\mathbb{C}}(V)$ which are constant as polynomials in x_0 . Therefore V lies in the image of \mathbb{Y} . Hence $k = d$ because $(u_0^0)^d t^d$ is a monomial of the Chow form of V whose coefficient is constant in t . \square

Now we want to extend ϕ : Let $V \in \mathcal{T}_d$ and F_V be a fixed representative of the Chow form of V . Let k be the degree of $F_V(t\underline{u}^0, \dots, t\underline{u}^{r+1})$ as a polynomial in t . For $\{t_n \in]0, \infty[\}_{n \in \mathbb{N}}$ one obtains a sequence $\{\phi(V, t_n)\}_{n \in \mathbb{N}}$ in the Chow variety. If one divides the n -th element of this sequence by $(t_n)^k$, the sequence remains invariant, because Chow forms are only determined up to a factor in \mathbb{C}^* . If $\lim_{n \rightarrow \infty} t_n = \infty$ we obtain that the original coefficients of $(t_n)^k$ remain invariant and the original coefficients of $(t_n)^l$, $l < k$, tend to zero, showing that the sequence is convergent. Therefore $\phi(V, \infty) := \lim_{t \rightarrow \infty} \phi(V, t)$ is an element of the Chow variety $\mathcal{C}_{r+1,d}(\mathbb{P}^{n+1}(\mathbb{C}))$. Furthermore we have already seen that the Chow form $F_{\phi(V, \infty)}(t\underline{u}^0, \dots, t\underline{u}^{r+1})$ is homogeneous of degree k as a polynomial in t .

If one could check that this degree k in t equals d , then by 3.4 the map ϕ retracts \mathcal{T}_d to $\mathbb{Y}\mathcal{C}_{r,d}(\mathbb{P}^n(\mathbb{C}))$: The degree k in t cannot be lower than the degree d

of V , because the monomial $(u_0^0)^d t^d$ appears in the Chow form of $V \in \mathcal{T}_d$. Hence $k \geq d$.

On the other hand from Lemma 3.4 for each irreducible component V_i one knows that $F_{\phi(V_i, \infty)}(t\underline{u}^0, \dots, t\underline{u}^{r+1})$ is homogeneous of degree $\deg V_i$ or of degree zero as polynomial in t . But in the case where the degree is zero for one irreducible component one obtains $k < d$. This contradicts $k \geq d$. Hence k has to be d . This proves the following lemma:

Lemma 3.5. *The map $\phi : \mathcal{T}_d \times [1, \infty] \rightarrow \mathcal{T}_d$ is a homotopy, which retracts \mathcal{T}_d to $\mathbb{Z}\mathcal{C}_{r,d}(\mathbb{P}^n(\mathbb{C}))$.*

The first part of the complex suspension theorem is proved. We need some preparations for the second part of the proof, where we have to check that the inclusion $\mathcal{T} \hookrightarrow \mathcal{C}_{r+1}(\mathbb{P}^{n+1}(\mathbb{C}))$ induces isomorphisms on homotopy groups:

Notation 3.6. In the following let $\underline{z}_1 := (1, 1, 0, \dots, 0) \in \mathbb{P}^{n+2}(\mathbb{C})$ and

$$\mathcal{C}'_{r+1,d}(\mathbb{P}^{n+2}(\mathbb{C})) := \{V \in \mathcal{C}_{r+1,d}(\mathbb{P}^{n+2}(\mathbb{C})) \mid \underline{z}_1 \notin \text{supp}(V)\}.$$

Further let $pr_1 : \mathbb{P}^{n+2}(\mathbb{C}) - \{\underline{z}_1\} \rightarrow \mathbb{P}^{n+1}(\mathbb{C})$ be the projection from \underline{z}_1 onto the hyperplane given by the equation $x_0 = 0$.

Suppose $V \in \mathcal{C}'_{r+1,d}(\mathbb{P}^{n+2}(\mathbb{C}))$, then the algebraic set $pr_1(\text{supp}(V))$ has dimension $r+1$ because the projection is an algebraic map, and the restriction $pr_1|_V$ has finite fibers (otherwise $\underline{z}_1 \in V$). Now we want to define a map π_1 for cycles $V \in \mathcal{C}'_{r+1,d}(\mathbb{P}^{n+2}(\mathbb{C}))$ such that $\text{supp}(\pi_1(V)) = pr_1(\text{supp}(V))$.

Let θ be the following ring homomorphism:

$$\begin{array}{ccc} \theta : \mathbb{C}[\underline{u}^0, \dots, \underline{u}^r] & \longrightarrow & \mathbb{C}[\underline{\tilde{u}}^0, \dots, \underline{\tilde{u}}^r], \\ u_0^j & \longmapsto & -u_1^j, & j = 0, \dots, r, \\ z & \longmapsto & z & \forall z \in \mathbb{C}[\underline{\tilde{u}}^0, \dots, \underline{\tilde{u}}^r], \end{array}$$

where $\underline{\tilde{u}}^j := (u_1^j, \dots, u_{n+2}^j)$ and $\underline{u}^j = (u_0^j, \dots, u_{n+2}^j)$.

Lemma 3.7. *θ maps the Chow form of a cycle $V \in \mathcal{C}'_{r+1,d}(\mathbb{P}^{n+2}(\mathbb{C}))$ to the Chow form of a cycle, which is denoted $\pi_1(V)$, in $\mathcal{C}_{r+1,d}(\mathbb{P}^{n+1}(\mathbb{C}))$, such that*

$$\text{supp}(\pi_1(V)) = pr_1(\text{supp}(V)).$$

Proof. Let $V \in \mathcal{C}'_{r+1,d}(\mathbb{P}^{n+2}(\mathbb{C}))$ be a variety and F_V be the Chow form of V . Further let $f_1, \dots, f_k \in \mathbb{C}[x_0, \dots, x_{n+2}]$ be homogeneous generators of the vanishing ideal of V , and let $g_1, \dots, g_l \in \mathbb{C}[x_1, \dots, x_{n+2}]$ be homogeneous generators of the vanishing ideal of $pr_1(V)$. F_V is the generator of the radical of the resultant of the following equations:

$$(2) \quad f_1(\underline{x}) = \dots = f_k(\underline{x}) = \begin{pmatrix} u_0^0 & u_1^0 & \cdots & u_{n+2}^0 \\ \vdots & \vdots & & \vdots \\ u_0^r & u_1^r & \cdots & u_{n+2}^r \end{pmatrix} \cdot \underline{x} = 0.$$

But the resultant of (2) is surjectively (R. 1.9) mapped under θ onto the resultant of

$$(3) \quad f_1(\underline{x}) = \dots = f_k(\underline{x}) = \begin{pmatrix} -u_1^0 & u_1^0 & \cdots & u_{n+2}^0 \\ \vdots & \vdots & & \vdots \\ -u_1^r & u_1^r & \cdots & u_{n+2}^r \end{pmatrix} \cdot \underline{x} = 0.$$

Therefore the radical of the ideal which is generated by $\theta(F_V)$ in $\mathbb{C}[\underline{u}^0, \dots, \underline{u}^r]$ is the radical of the resultant of (3). But the resultant of (3) is the same as the resultant of

$$(4) \quad g_1(\underline{x}) = \dots = g_l(\underline{x}) = \begin{pmatrix} u_1^0 & \dots & u_{n+2}^0 \\ \vdots & & \vdots \\ u_1^r & \dots & u_{n+2}^r \end{pmatrix} \cdot \begin{pmatrix} x_1 \\ \vdots \\ x_{n+2} \end{pmatrix} = 0.$$

To check this let $\underline{v}^0, \dots, \underline{v}^r \in \Omega^{n+2}$ be a zero of the resultant of (3). That means the equations

$$f_1(\underline{x}) = \dots = f_k(\underline{x}) = \begin{pmatrix} -v_1^0 & v_1^0 & \dots & v_{n+2}^0 \\ \vdots & \vdots & & \vdots \\ -v_1^r & v_1^r & \dots & v_{n+2}^r \end{pmatrix} \cdot \underline{x} = 0$$

have a solution $(q_0, \dots, q_{n+2}) \in V$. But $(q_1 - q_0, q_2, \dots, q_{n+2})$ is contained in $pr_1(V)$ and therefore a solution of the equations

$$g_1(\underline{x}) = \dots = g_l(\underline{x}) = \begin{pmatrix} v_1^0 & \dots & v_{n+2}^r \\ \vdots & & \vdots \\ v_1^r & \dots & v_{n+2}^r \end{pmatrix} \cdot \begin{pmatrix} x_1 \\ \vdots \\ x_{n+2} \end{pmatrix} = 0.$$

Therefore the radical of the resultant of (4) is a subideal of the radical of the resultant of (3). In the analogous way one checks the inverse inclusion.

This shows that $\theta(F_V) = F_{pr_1(V)}^m$ for some $m \in \mathbb{N} - \{0\}$. Since θ is a homomorphism of rings, we obtain for an arbitrary cycle $V = \sum_{i=1}^s n_i V_i$:

$$\theta(F_V) = (F_{V_1}^{n_1})^{m_1} \dots (F_{V_s}^{n_s})^{m_s}$$

for some $m_1, \dots, m_s \in \mathbb{N} - \{0\}$, where F_{V_i} denotes the Chow form of V_i .

This proves that $\theta(F_V)$ is a Chow form of multidegree d and the support of the cycle $\pi_1(V)$, whose Chow form is $\theta(F_V)$, is $pr_1(V)$. \square

Notation 3.8. The Chow variety $\mathcal{C}_{n+1,d}(\mathbb{P}^{n+2}(\mathbb{C}))$ of effective divisors of degree d can be identified with the equivalence classes of homogeneous polynomials in $\mathbb{C}[x_0, \dots, x_{n+2}]$ of degree d (Lemma 1.13), where $f \sim g$ iff $f = zg$ for some $z \in \mathbb{C}^*$. Let $Div'_d \subset \mathcal{C}_{n+1,d}(\mathbb{P}^{n+2}(\mathbb{C}))$ denote the set of cycles

$$\sum_{i_0, \dots, i_{n+2}=d} a_{i_0, \dots, i_{n+2}} x_0^{i_0} \dots x_{n+2}^{i_{n+2}},$$

such that $a_{d,0,\dots,0} \neq 0$ and $\sum_{i_0+i_1=d} a_{i_0,i_1,0,\dots,0} = 0$.

Div'_d is a quasi projective algebraic subset of $\mathcal{C}_{n+1,d}(\mathbb{P}^{n+2}(\mathbb{C}))$ of codimension 1. Hence it has dimension $\binom{n+2+d}{d} - 2$. Because elements in Div'_d are only determined up to a factor in \mathbb{C}^* , we can suppose $a_{d,0,\dots,0} = 1$. The space Div'_d is an affine subspace of $\mathcal{C}_{n+1,d}(\mathbb{P}^{n+2}(\mathbb{C}))$, hence it is contractible for instance by the following homotopy: Let I be the index set $I := \{(i_0, \dots, i_{n+2}) \in \mathbb{N}_0 \mid i_0 + \dots + i_{n+2} = d, i_0 \neq d\}$.

$$\begin{aligned} \Phi : \quad & Div'_d \times [0, 1] \longrightarrow Div'_d, \\ & \left(x_0^d + \sum_I a_{i_0, \dots, i_{n+2}} x_0^{i_0} \dots x_{n+2}^{i_{n+2}}, t \right) \longmapsto x_0^d + \sum_I t \cdot a_{i_0, \dots, i_{n+2}} x_0^{i_0} \dots x_{n+2}^{i_{n+2}}. \end{aligned}$$

$\Phi(Div'_d, 0)$ is the hyperplane $x_0 = 0$ with multiplicity d . Simple calculations show that $\underline{z}_0 := (1, 0, \dots, 0)$ and $\underline{z}_1 := (1, 1, 0, \dots, 0)$ are not contained in the support of any cycle in Div'_d .

Now we are able to define a map which will prove the second part of the complex suspension theorem:

Definition 3.9.

$$\Psi := \pi_1 \circ \cap \circ (\mathbb{Y}, \Phi) : \mathcal{C}_{r+1,d}(\mathbb{P}^{n+1}(\mathbb{C})) \times Div'_k \times [0, 1] \longrightarrow \mathcal{C}_{r+1,dk}(\mathbb{P}^{n+1}(\mathbb{C})) ,$$

where \cap denotes the intersection of cycles in the sense of Barlet [1].

We want to check that Ψ is well defined: First we have $\Phi(Div'_k, [0, 1]) \subset Div'_k$. Each irreducible component of a cycle in $\mathbb{Y}\mathcal{C}_{r+1,d}(\mathbb{P}^{n+1}(\mathbb{C}))$ meets the point \underline{z}_0 , and \underline{z}_0 is not contained in the support of any element in Div'_k . Therefore a pair of cycles in $\mathbb{Y}\mathcal{C}_{r+1,d}(\mathbb{P}^{n+1}(\mathbb{C})) \times Div'_k$ is in general position and the intersection is well defined. The support of the resulting cycle is just the geometric intersection of the supports of the original cycles. Hence the support of the resulting cycle cannot contain the point \underline{z}_1 , because \underline{z}_1 is not contained in the support of any cycle in Div'_k . That means the map π_1 is well defined in this situation.

The maps \mathbb{Y} , Φ , and π_1 are algebraic and \cap is analytic; hence Ψ is analytic. Ψ has nice geometric properties. First the map $\Psi(_, _, 0)$ is multiplication by k . Moreover it moves subsets of $\mathcal{C}_{r+1,d}(\mathbb{P}^{n+1}(\mathbb{C})) - \mathcal{T}_d$ to \mathcal{T}_{dk} for fixed $D \in Div'_k$. On the other hand Ψ maps cycles in $\mathbb{Y}\mathcal{C}_{r,d}(\mathbb{P}^n(\mathbb{C}))$ to cycles in $\mathbb{Y}\mathcal{C}_{r,dk}(\mathbb{P}^n(\mathbb{C}))$.

Lemma 3.10. *The map $\Psi|_{\mathbb{Y}\mathcal{C}_{r,d}(\mathbb{P}^n(\mathbb{C})) \times Div'_k \times [0, 1]}$ is the map which sends a cycle $V \in \mathbb{Y}\mathcal{C}_{r,d}(\mathbb{P}^n(\mathbb{C}))$ to the cycle $k \cdot V$.*

Proof. The support of an element in $\mathbb{Y}\mathcal{C}_{r,d}(\mathbb{P}^n(\mathbb{C}))$ can be viewed as a union of points lying on lines through points $(0, p_1, \dots, p_{n+1})$ and $(1, 0, \dots, 0)$ in $\mathbb{P}^{n+1}(\mathbb{C})$. Therefore it suffices to check the case $r = 0$ and $d = 1$. Ψ first takes the complex suspension of this line l . The complex suspension is cut with a cycle $D \in Div'_k$, and the common points of D and l are projected from \underline{z}_1 to the hyperplane $x_0 = 0$ in $\mathbb{P}^{n+2}(\mathbb{C})$. See Figure 1.

Therefore the support of $\Psi(l, D, t)$ is a closed subset of l which has dimension 1. But a closed 1-dimensional subset of a line is the line itself. Hence $\Psi(l, D, t) = k \cdot l$. \square

For a fixed $D \in Div'_k$ the map Ψ is a homotopy. We now determine which divisors $D \in Div'_k$ move $V \in \mathcal{C}_{r+1,d}(\mathbb{P}^{n+1}(\mathbb{C}))$ to an element in \mathcal{T}_{dk} .

Lemma 3.11. *Let π_0 denote the projection from \underline{z}_0 onto the hyperplane $x_0 = 0$. There is an isomorphism $F := (\pi_0|_{\overline{\pi_1^{-1}(\mathbb{P}^n(\mathbb{C}))}})^{-1} : \mathbb{P}^{n+1}(\mathbb{C}) \rightarrow \pi_1^{-1}(\mathbb{P}^n(\mathbb{C}))$ such that*

$$\Psi(V, D, 1) \notin \mathcal{T}_{dk} \implies F(\text{supp}(V)) \subset \text{supp}(D) .$$

A proof of this lemma is given in [14, pp. 286, 287], where no arguments of continuity are needed, one only has to check this pointwise.

For a fixed cycle $V \in \mathcal{C}_{r+1,d}(\mathbb{P}^{n+1}(\mathbb{C}))$ let $\mathcal{B}(V)$ denote the set of those cycles $D \in Div'_k$ such that $\Psi(V, D, 1) \notin \mathcal{T}_{dk}$. As one has already seen $\mathcal{B}(V) = \{D \in Div'_k \mid F(\text{supp}(V)) \subset \text{supp}(D)\}$. In general $\mathcal{B}(V)$ is a proper subset of Div'_k .




FIGURE 1

Lemma 3.12. *Given $V = \sum_{i=1}^m n_i V_i \in \mathcal{C}_{r+1,d}(\mathbb{P}^{n+1}(\mathbb{C}))$, the complex codimension of $\mathcal{B}(V)$ in Div'_k satisfies:*

$$\text{codim}_{\mathbb{C}} \mathcal{B}(V) = \max_i \text{codim}_{\mathbb{C}} \mathcal{B}(V_i) \geq \binom{r+k+1}{k} - 1.$$

Proof. We expand the ideas of [14, p. 288] (see also [8, p. 83]). It suffices to check the case $m = 1$, therefore let V be a variety. The embedding of V in \mathbb{P}^{n+2} can be viewed as an affine variety in $\mathbb{C}^{n+2} \cong \mathbb{P}^{n+2}(\mathbb{C}) - H$, where $V \not\subset H$ and $z_0 \in H$. So H is the zero set of $\sum_{i=1}^{n+2} a_i x_i$ for some $a_1, \dots, a_{n+2} \in \mathbb{C}$. First we want to check that if $V \not\subset H$ then $F(V)$ is not contained in H :

Let $\underline{p} := (0, p_1, \dots, p_{n+2}) \in V \subset \mathbb{P}^{n+2}(\mathbb{C})$. It follows directly from the definition of F (Lemma 3.11) that $F(\underline{p}) = (-p_1, p_1, \dots, p_{n+2})$. Hence if $\underline{p} \notin H$, which means $\sum_{i=1}^{n+2} a_i p_i \neq 0$, then $F(\underline{p}) \notin H$.

For simplicity we assume H to be the zero set of $x_{n+2} = 0$. In this situation the set Div'_k can be viewed as a subset of the polynomials of degree $\leq k$ in x_0, \dots, x_{n+1} . And the set $\mathcal{B}(V) \subset \text{Div}'_k$ is the set of those polynomials, which vanish on $F(V) \cap \mathbb{C}^{n+2}$.

Since $F(V)$ is $r+1$ -dimensional, if one projects $F(V) \cap \mathbb{C}^{n+2}$ onto a linear subspace of dimension $r+1$, then without loss of generality the image of $F(V) \cap \mathbb{C}^{n+2}$ has dimension $r+1$, too, because $F(V) \cap \mathbb{C}^{n+2}$ contains regular points. Up to linear isomorphisms one can suppose the linear subspace to be the zero set of $x_{r+1} = \dots = x_{n+1} = 0$.

In this situation a polynomial $f \in \mathbb{C}[x_0, \dots, x_r]$ can only vanish on the image of $F(V) \cap \mathbb{C}^{n+2}$ under the projection, if $f = 0$. This is obvious, because the zero set of f is a r -dimensional subset of the linear subspace.

Therefore the complex codimension of $\mathcal{B}(V)$ in $\mathcal{C}_{n+1,k}(\mathbb{P}^{n+1}(\mathbb{C}))$ is greater or equal to the dimension of the space of polynomials in $\mathbb{C}[x_0, \dots, x_r]$ of degree $\leq k$. This is $\binom{r+k+1}{k}$. Because $\text{Div}'_k \subset \mathcal{C}_{n+1,k}(\mathbb{P}^{n+2}(\mathbb{C}))$ has codimension 1 the lemma is proved. \square

Now let \mathcal{B} denote the following set:

$$\begin{aligned} \mathcal{B} &:= (\Psi|_{\mathcal{C}_{r+1,d}(\mathbb{P}^{n+1}(\mathbb{C})) \times \text{Div}'_k \times \{1\}})^{-1}(\mathcal{C}_{r+1,dk}(\mathbb{P}^{n+1}(\mathbb{C})) - \mathcal{T}_{dk}) \\ &= \{(V, D) \in \mathcal{C}_{r+1,d}(\mathbb{P}^{n+1}(\mathbb{C})) \times \text{Div}'_k \mid \Psi(V, D, 1) \notin \mathcal{T}_{dk}\}. \end{aligned}$$

The set $\mathcal{C}_{r+1,dk}(\mathbb{P}^{n+1}(\mathbb{C})) - \mathcal{T}_{dk}$ is an algebraic set and Ψ is an analytic map, hence \mathcal{B} is an analytic set. Let $V \in \mathcal{C}_{r+1,d}(\mathbb{P}^{n+1}(\mathbb{C}))$ then $(V \times \text{Div}'_k) \cap \mathcal{B} \subset \mathcal{B}(V)$. Therefore the codimension of \mathcal{B} is greater or equal $\binom{r+k+1}{k} - 1$. Now choose $E(r, d) \in \mathbb{N}$ such that $\binom{r+k+1}{k} - 1 > \dim \mathcal{C}_{r+1,d}(\mathbb{P}^{n+1}(\mathbb{C}))$, $\forall k \geq E(r, d)$. Further denote by pr_2 the projection from $\mathcal{C}_{r+1,d}(\mathbb{P}^{n+1}(\mathbb{C})) \times \text{Div}'_k$ onto the second factor. Then one obtains

$$\begin{aligned} \dim pr_2(\mathcal{B}) &\leq \dim \mathcal{B} \\ &\leq \dim \mathcal{C}_{r+1,d}(\mathbb{P}^{n+1}(\mathbb{C})) + \dim \text{Div}'_k - \binom{r+k+1}{k} + 1 \\ &< \dim \text{Div}'_k. \end{aligned}$$

Hence the projection from \mathcal{B} on Div'_k cannot be surjective $\forall k > E(r, d)$. Therefore one obtains a cycle $\mathcal{D} \in \text{Div}'_k$ for each $k \geq E(r, d)$ such that the map $\Psi : \mathcal{C}_{r+1,d}(\mathbb{P}^{n+1}(\mathbb{C})) \times \{\mathcal{D}\} \times [0, 1] \rightarrow \mathcal{C}_{r+1,dk}(\mathbb{P}^{n+1}(\mathbb{C}))$ has the following properties:

$$\Psi(V, \mathcal{D}, 0) = k \cdot V \quad \text{and} \quad \Psi(\mathcal{C}_{r+1,d}(\mathbb{P}^{n+1}(\mathbb{C})), \mathcal{D}, 1) \subset \mathcal{T}_{dk}.$$

To finish the proof of the complex suspension theorem one has to check that the inclusion $i : \mathcal{T} \hookrightarrow \mathcal{C}_{r+1}(\mathbb{P}^{n+1}(\mathbb{C}))$ induces isomorphisms on homotopy groups. For this purpose let $\mathbb{Z}\mathbb{P}^r(\mathbb{C})$ be the basepoint $*$ of \mathcal{T} and $\mathcal{C}_{r+1}(\mathbb{P}^{n+1}(\mathbb{C}))$. First of all the monoid $\mathcal{C}_{r+1}(\mathbb{P}^{n+1}(\mathbb{C}))$ is connected. This is obvious from the properties of Ψ . Furthermore, the addition on homotopy groups, which is induced by the H-cogroup structure of the spheres, coincides with the addition which is induced by the topological monoid structure of \mathcal{T} and $\mathcal{C}_{r+1}(\mathbb{P}^{n+1}(\mathbb{C}))$.

Surjectivity: Let $\alpha \in \pi_j(\mathcal{C}_{r+1}(\mathbb{P}^{n+1}(\mathbb{C})))$. There is $d \in \mathbb{N}$ such that $\alpha : S^j \rightarrow \mathcal{C}_{r+1,d}(\mathbb{P}^{n+1}(\mathbb{C}))$. For $k \geq E(d, r)$ there exists $\alpha_k \in \pi_j(\mathcal{T}_{dk})$ such that $i_*[\alpha_k] \stackrel{\Psi}{=} [k \cdot \alpha]$. Therefore $\alpha_{k+1} - \alpha_k \in \pi_j(\mathcal{T})$ and $i_*[\alpha_{k+1} - \alpha_k] = \alpha$.

Injectivity: Let $\beta \in \pi_j(\mathcal{T})$ such that $i_*[\beta] = *$. Without loss of generality suppose $\beta : S^j \rightarrow \mathbb{Z}\mathcal{C}_r(\mathbb{P}^n(\mathbb{C}))$, because $\mathbb{Z}\mathcal{C}_r(\mathbb{P}^n(\mathbb{C}))$ is a deformation retract of \mathcal{T} . Now there is a map $h : S^j \times [0, 1] \rightarrow \mathcal{C}_{r+1}(\mathbb{P}^{n+1}(\mathbb{C}))$ such that $h(\cdot, 0) = i_*(\beta)$ and $h(\cdot, 1) = *$. There exists $d \in \mathbb{N}$ such that $h(S^j \times [0, 1]) \subset \mathcal{C}_{r+1,d}(\mathbb{P}^{n+1}(\mathbb{C}))$. For each $k \geq E(d, r)$ there exists $h_k : S^j \times [0, 1] \rightarrow \mathcal{T}_{dk}$ such that $i_*[h_k] \stackrel{\Psi}{=} [k \cdot h]$. Because Ψ restricted to $\mathbb{Z}\mathcal{C}_{r,d}(\mathbb{P}^n(\mathbb{C}))$ is multiplication by d one obtains $h_k(\cdot, 1) = *$ and $h_k(\cdot, 0) = k \cdot \beta$. In this situation one obtains $(h_{k+1} - h_k) : S^j \times [0, 1] \rightarrow \mathcal{T}$ such that $(h_{k+1} - h_k)(\cdot, 1) = *$ and $(h_{k+1} - h_k)(\cdot, 0) = \beta$.

The complex suspension theorem enables one to determine the homotopy type of the monoids $\mathcal{C}_r(\mathbb{P}^n(\mathbb{C}))$.

Corollary 3.13. *The topological monoid $\mathcal{C}_r(\mathbb{P}^n(\mathbb{C}))$ is homotopy equivalent to a product of Eilenberg–Mac Lane spaces:*

$$\mathcal{C}_r(\mathbb{P}^n(\mathbb{C})) \simeq \prod_{i=1}^{n-r} K(\mathbb{Z}, 2i) .$$

Proof. Let H be the unreduced singular homology theory. By Dold/Thom [7, Satz 7.1] we have

$$\widetilde{\mathrm{SP}}(\mathbb{P}^{n-r}(\mathbb{C})) \simeq \prod_{i>0} K(H_k(\mathbb{P}^{n-r}(\mathbb{C}), \mathbb{Z}), i)$$

$$\text{Hence } \mathcal{C}_r(\mathbb{P}^n(\mathbb{C})) \xrightarrow{\simeq} \dots \xrightarrow{\simeq} \mathcal{C}_0(\mathbb{P}^{n-r}(\mathbb{C})) \cong \widetilde{\mathrm{SP}}(\mathbb{P}^{n-r}(\mathbb{C})) \simeq \prod_{i=1}^{n-r} K(\mathbb{Z}, 2i). \quad \square$$

It follows that \mathcal{C} is homotopy equivalent to the weak product $\prod_{i>0} K(\mathbb{Z}, 2i)$. The naive group completion of $\mathfrak{C} = \coprod_d \mathcal{D}(d)$ is then homotopy equivalent to $\mathcal{C} \times \mathbb{Z}$ [15, Thm. 5.4] using the complex suspension theorem and Dold/Thom [7, Thm. 6.10.III]. The naive group completion of \mathfrak{C} is homotopy equivalent to the homotopy theoretic group completion $\Omega B(\mathfrak{C})$ by [16, Thm. 4.4']. Hence we obtain $\Omega B(\mathfrak{C}) \simeq \prod_{i>0} K(\mathbb{Z}, 2i)$. The inclusion $BU \cong \mathcal{D}(1) \hookrightarrow \mathcal{C}$ represents the total Chern class [15, Thm. 3.6]. Assuming continuity Boyer, Lawson, Lima–Filho, Mann and Michelsohn [4] proved that there is an \mathcal{I}_* -functor T [17] which induces an infinite loop space structure on $\Omega B\mathfrak{C}$ compatible with the infinite loop space structure on $\mathcal{D}(1) \cong BU$ induced by Whitney sum of vector bundles. Hence the total Chern class map is an infinite loop map.

4. THE CONTINUITY OF THE \mathcal{I}_* -FUNCTOR DEFINED BY CHOW VARIETIES

The authors of [4] provided a framework for producing an \mathcal{I}_* -functor (cp. [17, Def. I.1.8]) from Chow varieties. Here we show, how continuity assertions in [4] can be derived in the language of resultants. The continuity of the \mathcal{I}_* -functor T follows from the following lemma.

Lemma 4.1. *Let $V \subset \mathbb{C}^m$ be a n -dimensional vector space and $W \subset \mathbb{C}^m$ a p -dimensional vector space. If $f : V \rightarrow W$ is a linear, isometric embedding, then the map*

$$\begin{aligned} Tf : \coprod_d \mathcal{C}_{n-1,d}(\mathbb{P}(V \oplus V)) &\longrightarrow \coprod_d \mathcal{C}_{p-1,d}(\mathbb{P}(W \oplus W)) \\ c &\longmapsto \mathbb{P}(f(V)^\perp \oplus \{0_W\}) \# (f \oplus f)_*(c) \end{aligned}$$

is algebraic.

Proof. The complex join is algebraic and building orthogonal complements is an algebraic map on the Grassmannian. Hence it suffices to check that

$$(f \oplus f)_* : \coprod_d \mathcal{C}_{n-1,d}(\mathbb{P}(V \oplus V)) \longrightarrow \coprod_d \mathcal{C}_{n-1,d}(\mathbb{P}(W \oplus W))$$

is algebraic. We shall calculate the Chow form $F_{(f \oplus f)_*(c)}$ of the cycle $(f \oplus f)_*(c)$. We can extend the map $(f \oplus f)$ to an automorphism of the vector space \mathbb{C}^{2m} by choice of the orthogonal complement of $V \oplus V$ in \mathbb{C}^{2m} . This automorphism is given by an invertible $(2m \times 2m)$ -matrix A over \mathbb{C} .

Without loss of generality suppose c to consist only on one irreducible component. Let $g_1, \dots, g_k \in \mathbb{C}[x_0, \dots, x_{2m-1}]$ be homogeneous generators of the vanishing ideal of c . Then $(f \oplus f)_*(c)$ is the irreducible zero set of

$$g_1(A^{-1}\underline{x}), \dots, g_k(A^{-1}\underline{x}) .$$

The Chow form F_c of c generates the radical of the resultant of

$$g_1(\underline{x}) = \dots = g_k(\underline{x}) = \begin{pmatrix} u_0^0 & \dots & u_{2m-1}^0 \\ \vdots & & \vdots \\ u_0^{n-1} & \dots & u_{2m-1}^{n-1} \end{pmatrix} \cdot \underline{x} = 0 .$$

The solutions of this equations are the same as the solutions of

$$(5) \quad g_1(A^{-1}\underline{x}) = \dots = g_k(A^{-1}\underline{x}) = \begin{pmatrix} u_0^0 & \dots & u_{2m-1}^0 \\ \vdots & & \vdots \\ u_0^{n-1} & \dots & u_{2m-1}^{n-1} \end{pmatrix} A^{-1}\underline{x} = 0 .$$

Therefore F_c generates the radical of the resultant of (5), too. The following map is a ring automorphism:

$$\begin{array}{ccc} \varphi_f : \mathbb{C}[\underline{u}^0, \dots, \underline{u}^{n-1}] & \rightarrow & \mathbb{C}[\underline{u}^0, \dots, \underline{u}^{n-1}], \\ \begin{matrix} u_j^i \\ z \end{matrix} & \mapsto & \begin{matrix} (\underline{u}^i A)_j \\ z \end{matrix}, & i = 0, \dots, n-1, j = 0, \dots, 2m-1, \\ & & \forall z \in \mathbb{C}, \end{array}$$

because the $(\underline{u}^i A)_j$ are algebraically independent over \mathbb{C} , $i = 0, \dots, n-1$, $j = 0, \dots, 2m-1$, for the entries of A are elements of \mathbb{C} . The equations (5) are mapped onto

$$(6) \quad g_1(A^{-1}\underline{x}) = \dots = g_k(A^{-1}\underline{x}) = \begin{pmatrix} u_0^0 & \dots & u_{2m-1}^0 \\ \vdots & & \vdots \\ u_0^{n-1} & \dots & u_{2m-1}^{n-1} \end{pmatrix} A A^{-1}\underline{x} = 0 .$$

Hence $\varphi_f(F_c) = F_c(\underline{u}^0 A, \dots, \underline{u}^{n-1} A)$ generates the radical of the resultant of (6). But $F_{(f \oplus f)_*(c)}$ is a generator of the radical of the resultant of (6), too. Hence $\varphi_f(F_c)$ and $F_{(f \oplus f)_*(c)}$ coincide up to a factor in \mathbb{C}^* . But this means that $\varphi_f(F_c)$ is the Chow form of $(f \oplus f)_*$. φ_f depends on an extension of the linear isometry $(f \oplus f)$ to a matrix A . But $\varphi_f(F_c)$ coincides with $F_{(f \oplus f)_*}$ and is therefore independent of a chosen extension.

We obtain, that $(f \oplus f)_*$ is given on Chow forms by the map φ_f , which is polynomial in the coefficients of Chow forms. Hence $(f \oplus f)_*$ and Tf are algebraic. \square

REFERENCES

- [1] D. Barlet, Espace analytique réduit des cycles analytiques complexes compacts d' un espace analytique complexe de dimension finie, Fonctions de Plusieurs Variables Complexes II (Séminaire F. Norguet), LNM 482. Springer-Verlag, Berlin, Heidelberg, New York (1975), 1 - 158. MR **53**:3347
- [2] D. Barlet, Preprint (1993).
- [3] J. M. Boardman and R. M. Vogt, Homotopy Invariant Algebraic Structures on Topological Spaces, LNM 347. Springer-Verlag, Berlin, Heidelberg, New York (1973). MR **54**:8623a
- [4] C. P. Boyer, H. B. Lawson, P. Lima-Filho, B. M. Mann, and M.-L. Michelsohn, Algebraic Cycles and Infinite Loop Spaces. *Inventiones Math.* 113 (1993), 373-388. MR **95a**:55021
- [5] W.-L. Chow and B. L. van der Waerden, ZAG 9 Über zugeordnete Formen und algebraische Systeme von algebraischen Mannigfaltigkeiten. *Math. Annalen* 113 (1937), 692 - 704.

- [6] J. Dalbec, The Join of Two Varieties. Preprint.
- [7] A. Dold and R. Thom, Quasifaserungen und unendliche symmetrische Produkte. Ann. of Math. 67 (1958), 230 - 281. MR **20**:3542
- [8] E. M. Friedlander, Algebraic cycles, Chow varieties, and Lawson homology. Comp. Math. 77 (1991), 55–93. MR **92a**:14005
- [9] I. M. Gelfand, M. M. Kapranov, and A. V. Zelevinsky, Discriminants, Resultants, and Multidimensional Determinants. Birkhäuser, Boston, Basel, Berlin (1994). MR **95e**:14045
- [10] P. Griffiths and J. Adams, Topics in algebraic and analytic geometry. Princeton University Press and University of Tokyo Press, Princeton, New Jersey (1974). MR **50**:7596
- [11] R. Hartshorne, Algebraic Geometry. Springer–Verlag, New York Inc. (1977). MR **57**:3116
- [12] W. V. D. Hodge and D. Pedoe, Methods of Algebraic Geometry, Vol. II. Cambridge University Press (1952). MR **13**:972c
- [13] E. Kunz, Introduction to Commutative Algebra and Algebraic Geometry. Birkhäuser, Boston (1985). MR **86e**:14001
- [14] H. B. Lawson, Algebraic Cycles and Homotopy Theory. Annals of Math. 129 (1989), 253 - 291. MR **90h**:14008
- [15] H. B. Lawson and M.–L. Michelsohn, Algebraic Cycles, Bott Periodicity, and the Chern Characteristic Map, “The Mathematical Heritage of Herman Weyl,” AMS (1988), 241 - 264. MR **90d**:14010
- [16] P. Lima–Filho, Completions and Fibrations for Topological Monoids. Trans. Amer. Math. Soc. 340, No. 1 (1993), 127–147. MR **94a**:55009
- [17] J. P. May, E_∞ Ring Spaces and E_∞ Ring Spectra, LMN 577. Springer–Verlag, Berlin, Heidelberg, New York (1977). MR **58**:13008
- [18] M. C. McCord, Classifying Spaces and Infinite Symmetric Products. Trans. Amer. Math. Soc. 146 (1969), 273 - 298. MR **40**:4946
- [19] D. Mumford, Algebraic Geometry I, Complex Projective Varieties. Springer–Verlag, Berlin, Heidelberg (1976). MR **56**:11992
- [20] J. Plümer, Stabilisierte Chow–Varietäten und die Chernklassenabbildung. Diplomarbeit, Universität Osnabrück (1993).
- [21] I. R. Šafarevič, Basic Algebraic Geometry. Springer–Verlag, Berlin, Heidelberg, New York (1974). MR **51**:3163
- [22] B. L. van der Waerden, Algebra 2. Springer–Verlag, Berlin, Heidelberg (1967). MR **38**:1968
- [23] B. L. van der Waerden, ZAG 6 Algebraische Korrespondenzen und rationale Abbildungen. Math. Annalen 110 (1935), 134 - 160.

UNIVERSITÄT OSNABRÜCK, FACHBEREICH MATHEMATIK/INFORMATIK, 49069 OSNABRÜCK, GERMANY

E-mail address: `judith@mathematik.uni-osnabrueck.de`