

APPROXIMATION BY HARMONIC FUNCTIONS

EVGENY A. POLETSKY

ABSTRACT. For a compact set $X \subset \mathbb{R}^n$ we construct a restoring covering for the space $h(X)$ of real-valued functions on X which can be uniformly approximated by harmonic functions. Functions from $h(X)$ restricted to an element Y of this covering possess some analytic properties. In particular, every nonnegative function $f \in h(Y)$, equal to 0 on an open non-void set, is equal to 0 on Y . Moreover, when $n = 2$, the algebra $H(Y)$ of complex-valued functions on Y which can be uniformly approximated by holomorphic functions is analytic. These theorems allow us to prove that if a compact set $X \subset \mathbb{C}$ has a nontrivial Jensen measure, then X contains a nontrivial compact set Y with analytic algebra $H(Y)$.

1. INTRODUCTION

The Shilov–Bishop theorem (see [Gam], Th. 13.1) asserts that any uniform algebra $A(X)$ on a compact set X has an antisymmetric restoring covering, i.e., a covering of X by disjoint compacta $\{X_\alpha\}$ such that: 1) this partition is restoring, i.e. if $f \in A|_{X_\alpha}$ for all α , then $f \in A(X)$; 2) the algebras $A|_{X_\alpha}$ are antisymmetric, i.e., if $\mathbf{Im} f = 0$ for $f \in A(X_\alpha)$, then f is a constant.

For $X \subset \mathbb{C}$ and the *complex* uniform algebra $H(X)$ of functions on X which can be uniformly approximated by holomorphic functions, for several years there were attempts to find restoring coverings of X by compacta X_α with analytic algebras $H(X_\alpha)$. Recall that a uniform algebra is *analytic* if every function from this algebra, which is equal to 0 on an open non-void set, is equal to 0 everywhere. However, McKissick ([M]) gave an example of a compact set X in \mathbb{C} such that the algebra $H(X)$ is a normal proper subset of the algebra $C(X)$ of continuous functions on X (see also [S], §27). (A uniform algebra A on X is normal if for every pair of disjoint sets K_1 and K_2 in X there exists a function $f \in A$ such that $f(K_1) = 0$ and $f(K_2) = 1$.) This example disproved the existence of analytic restoring coverings for $H(X)$, because the normality of $H(X)$ implies that for a compact set $Y \subset X$ the algebra $H(Y)$ is analytic if and only if Y consists of one point.

In this paper for the *real* space $h(X)$ of functions which can be uniformly approximated by functions harmonic on neighborhoods of X , we prove

Theorem 1.1. *For any compact set $X \subset \mathbb{R}^n$, $n \geq 2$, there exists a covering of X by disjoint subsets Q_α such that*

1. *$f \in h(X)$ if and only if $f|_{\overline{Q}_\alpha} \in h(\overline{Q}_\alpha)$ and*
2. *every nonconstant $f \in h(\overline{Q}_\alpha)$ cannot attain its maximum on Q_α .*

Received by the editors September 10, 1995.

1991 *Mathematics Subject Classification.* Primary 32F05; Secondary 32E25, 32E20.

Key words and phrases. Harmonic functions, potential theory, uniform algebras.

Our approach to this theorem is very geometric, and it allows us to get many properties of the sets Q_α and the spaces $h(\overline{Q}_\alpha)$. In particular, criteria for the identity $h(X) = C_{\mathbb{R}}(X)$, due to Keldysh [Ga] and Khavinson [Kh], and the Hartogs–Rosenthal theorem [Ga] follow from our results.

Also we are able to provide additional information on compacta X with analytic algebras $H(X)$. The author's interest in this topic was raised by Wermer's example of a compact set X in \mathbb{C}^2 without analytic structure [W]. Goldmann proved in [G] that $H(X)$ is analytic. This example is the cluster of some perfect sequence in \mathbb{C}^2 (see [P]), and in this paper we show that the algebra $H(X)$ is analytic whenever X is the cluster of a perfect sequence in \mathbb{C} .

I am grateful to T. Bagby, N. Levenberg, T. McConnell and J. Wermer for useful discussions during my work on this paper. I also wish to thank P. M. Gauthier and the referee, whose comments and corrections significantly improved the paper.

2. PROPERTIES OF HARMONIC MEASURES

We will denote by $B^n(a, r)$ and $S^n(a, r)$ the ball and the sphere of radius r with center a lying in the real n -dimensional space \mathbb{R}^n . We will omit a and r when $a = 0$ and $r = 1$. We also will omit n when it does not lead to misunderstanding. Let $R(z, s, t)$ be the spherical shell $\{x \in \mathbb{R}^n : s < \|x - z\| < t\}$ and $R(s, t) = R(0, s, t)$. A standard Lebesgue measure on sets in \mathbb{R}^n will be denoted by $m(dz)$, and $\sigma_n = m(B^n)$. The surface area measure on spheres will be denoted by $s(dz)$.

The Green function $G_D(x, y)$ on a domain D in this paper is positive and superharmonic (sph). For the unit ball B in \mathbb{R}^n let $\Gamma(t) = G_B(x, 0)$, where $\|x\| = t$. Then $\Gamma(t) = -\ln t$ when $n = 2$, and $\Gamma(t) = t^{2-n} - 1$ when $n > 2$.

Let D be a bounded open set in \mathbb{R}^n . Then for every point $x \in D$ there is the harmonic measure μ_x supported by ∂D such that for every bounded Borel function f on ∂D the function

$$u(x) = \int_{\partial D} f(y) \mu_x(dy)$$

is the harmonic extension of f into D (see [H] for details). For a closed set $E \subset \mathbb{R}^n$ we define $\omega(x, E, D) = \mu_x(E)$, where μ is the harmonic measure relative to $D \setminus E$ and x . It follows from the general theory (see [H]) that $\omega(x, E, D) = \inf u(x)$, where the infimum is taken over all positive sph functions on D with lower limits greater than 1 at points of E . It is easy to see that $\omega(z, \overline{B}(0, t), B) = \omega(z, \overline{S}(0, t), B) = \Gamma(\|z\|)/\Gamma(t)$, when $\|z\| \geq t$.

The following lemma provides connections between Green functions and harmonic measures.

Lemma 2.1. *Let D be an open set in $B \subset \mathbb{R}^n$ and $0 \in D$. If*

$$u_t(x) = \Gamma(t)\omega(x, \overline{B}(0, t), D),$$

then $u_t(x) \leq u_s(x)$ when $t < s$ and $\|x\| > s$, and

$$u_0(x) = \lim_{t \rightarrow 0} u_t(x) = G_D(x, 0).$$

Proof. Since $u_t(x) \leq \Gamma(\|x\|)$, we see that $u_t(x) \leq u_s(x)$ when $t < s$ and $\|x\| = s$. Thus $u_t(x) \leq u_s(x)$ when $\|x\| > s$.

Since $G_D(x, 0) \leq \Gamma(t)$ when $\|x\| = t$, we see that $u_t(x) \geq G_D(x, 0)$ when $\|x\| > t$ and, therefore, $u_0(x) \geq G_D(x, 0)$ on D . The function $u_0(x)$ is harmonic on $D \setminus \{0\}$

and sph on D . Thus $u_0 = cG_D$ for some $c > 0$. But $u_0(x) \leq \Gamma(\|x\|)$, so $u_0(x) \equiv G_D(x, 0)$. \square

The following theorem gives us an important estimate for Green functions.

Theorem 2.2. *Let $D \subset B \subset \mathbb{R}^n$ be an open set and $0 \in D$. Suppose that*

$$\omega(0, \overline{B}(z, r), D) = a > 0,$$

$k > 1$, and $kr \leq \|z\|/2$. Then there is a constant $b = b(k, n) > 0$ such that the measure of points $y \in R(z, r, kr)$, where

$$G_D(y, 0) \geq d = \frac{ab}{2\sigma_n(k^n - 1)r^{n-2}},$$

is greater than or equal to $\frac{abr^2}{2\Gamma(\|z\|/2)}$.

Proof. Let us find the continuous sph function u on $\mathbb{R}^n \setminus B$ such that $\Delta u = -m(dx)$ on the spherical shell $R(1, k)$, $u \equiv b(k, n)$ on $\mathbb{R}^n \setminus B(0, k)$, and $u \equiv 0$ on S . Clearly on $R(1, k)$

$$u(x) = -\frac{1}{4}(\|x\|^2 - 1) + c \ln \|x\|$$

when $n = 2$ and

$$u(x) = -\frac{1}{2n}(\|x\|^2 - 1) - \frac{c}{\|x\|^{n-2}} + c$$

when $n > 2$. Since, as a measure, Δu is equal to 0 on $S(0, k)$ and u is a constant outside $B(0, k)$, we conclude that the normal derivative of u on $S(0, k)$ is 0. This allows us to find c . Direct calculations show that

$$u(x) = -\frac{1}{4}(\|x\|^2 - 1) + \frac{k^2}{2} \ln \|x\|$$

on $R(1, k)$ when $n = 2$, and

$$b(k, 2) = \frac{1 - k^2}{4} + \frac{k^2 \ln k}{2}.$$

If $n > 2$ then

$$b(k, n) = \frac{1}{n-2} \left(\frac{k^n - 1}{n} - \frac{k^2 - 1}{2} \right)$$

and

$$u(x) = -\frac{1}{2n}(\|x\|^2 - 1) - \frac{k^n}{n(n-2)}(\|x\|^{2-n} - 1)$$

on $R(1, k)$.

Let

$$v(x) = r^2 u \left(\frac{x - z}{r} \right)$$

and $R = R(z, r, kr)$.

Let $D_1 = D \setminus \overline{B}(z, r)$, $g(y, x) = G_{D_1}(x, y)$, and let h be the greatest harmonic minorant of v on D_1 . Then by Theorem 6.18 of [H] the function

$$f(x) = v(x) - h(x) = \int_{R \cap D} g(x, y) m(dy).$$

The symmetry of the Green function implies that

$$f(0) = \int_{R \cap D} g(y, 0) m(dy).$$

On the other hand, by Theorems 8.11 and 8.15 of [H],

$$h(x) = \int_{\partial D_1} v(y) \mu_x(dy) \leq br^2 \omega(x, \partial D, D_1) = br^2(1 - \omega(x, \overline{B}(z, r), D)),$$

where in the first equality $\mu_x(E) = \omega(x, E, D_1)$ for $E \subset \partial D_1$. Therefore

$$\int_{R \cap D} g(y, 0) m(dy) = f(0) = v(0) - h(0) \geq abr^2.$$

Let $E = \{y \in R : g(y, 0) \geq d\}$. Since $g(y, 0) \leq \gamma$, where $\gamma = \Gamma(\|z\|/2)$ and $y \in R$, and $m(R) = \sigma_n(k^n - 1)r^n$, we see that

$$m(E) \geq \frac{abr^2 - dm(R)}{\gamma - d} \geq \frac{abr^2}{2\gamma}$$

and the theorem is proved because $G_D(x, y) \geq g(x, y)$. \square

We also will need the following trivial remark.

Lemma 2.3. *Let D be a bounded open set in \mathbb{R}^n and $z_0 \in D$. If $A \subset D$ is a closed set, $\omega(z_0, A, D) = b$, u is a positive sph function on D and $u \geq a$ on A , then $u(z_0) \geq ab$.*

Proof. Indeed

$$u(z_0) \geq a\omega(z_0, A, D) = ab.$$

\square

The following lemma provides lower estimates for the area $s(E)$ of a set E in $S(0, t)$ through the harmonic measure of E .

Lemma 2.4. *Let $n > 2$, $z_0 \in B$, and $\|z_0\| \geq k > 0$. If a closed set $E \subset S(0, t)$, $0 < t < k/2$, and $\omega(z_0, E, R(t, 1)) \geq \frac{b}{\Gamma(t)}$, $b > 0$, then there is a constant $d = d(k, b, n)$ such that $s(E) \geq dt^{n-1}$.*

Proof. The inversion

$$I(x) = \frac{tx}{\|x\|^2}$$

maps $R = R(t, 1)$ onto itself and the point z_0 into the point z_1 with $\|z_1\| = t\|z_0\|^{-1}$. If a set $E \subset S(0, t)$, then

$$\omega(z, I(E), R) = \frac{1}{\|z\|^{n-2}} \omega(I(z), E, R).$$

Let P_R be the Poisson kernel on the shell R . The function

$$\phi(t, m) = \inf_{x, y \in S} \frac{P_R(z, x)}{P_R(z, y)}, \quad \|z\| = \frac{t}{m},$$

is positive and continuous on the set of positive numbers m and t such that $t < m < 1$. Let us find the limit of $\phi(t, m)$ when t converges to 0 and m to m_0 , $0 < m_0 \leq l < 1$.

If $x \in S$ then

$$P_R(z, x) = P(z, x) - \int_{S(0, t)} P(y, x) \mu_z(dy),$$

where

$$P(z, x) = \frac{1}{n\sigma_n} \frac{1 - \|z\|^2}{\|z - x\|^n}$$

is the Poisson kernel on the unit ball and $\mu_z(F) = \omega(z, F, R)$ for $F \subset \partial R$. Recalling that

$$\omega(z, S(0, t), R(t, 1)) = \frac{\|z\|^{2-n} - 1}{t^{2-n} - 1} = \frac{m^{n-2} - t^{n-2}}{1 - t^{n-2}}$$

and the maximal value of $P(y, x)$ on $S(0, t)$ is equal to

$$\frac{1}{n\sigma_n} \frac{1 - t^2}{(1 - t)^n},$$

we see that

$$P_R(z, x) \geq P(z, x) \left(1 - \frac{1}{P(z, x)} \frac{1 - t^2}{n\sigma_n(1 - t)^n} \frac{m^{n-2} - t^{n-2}}{1 - t^{n-2}} \right).$$

Since $P(z, x) \rightarrow (n\sigma_n)^{-1}$ as $\|z\| = t/m$ goes to 0, we get

$$\liminf_{t \rightarrow 0, m \rightarrow m_0} \frac{P_R(z, x)}{P(z, x)} \geq 1 - m_0^{n-2} \geq 1 - l^{n-2}.$$

We know that the ratio $P(z, x)/P(z, y)$ converges uniformly in x and y to 1 as $\|z\| \rightarrow 0$, and $P(x, y) \geq P_R(x, y)$. Therefore,

$$\liminf_{t \rightarrow 0, m \rightarrow m_0} \frac{P_R(z, x)}{P_R(z, y)} \geq 1 - l^{n-2}$$

and

$$\liminf_{t \rightarrow 0, m \rightarrow m_0} \phi(t, m) \geq 1 - l^{n-2}.$$

Therefore, there is a constant $c = c(k, l, n) > 0$ such that on the set $K = \{(t, m) : k \leq m \leq l, 2t \leq m\}$ the function $\phi(t, m) \geq c$.

Since

$$\frac{\Gamma(\|z_0\|)}{\Gamma(t)} \geq \omega(z_0, E, R(t, 1)) \geq \frac{b}{\Gamma(t)},$$

there is a constant $l = l(b, n) < 1$ such that $\|z_0\| \leq l$, and according to the assumptions of our lemma $2t \leq k \leq \|z_0\|$. Hence $P_R(z_1, x) \geq cP_R(z_1, y)$. Since

$$\omega(z_1, E^*, R) = \int_{E^*} P_R(z_1, x) s(dx),$$

where $E^* = I(E)$, we see that

$$\frac{\omega(z_0, E, R)}{\omega(z_0, S(0, t), R)} = \frac{\omega(z_1, E^*, R)}{\omega(z_1, S, R)} \leq \frac{c^{-2}s(E^*)}{s(S)} = \frac{t^{1-n}s(E)}{nc^2\sigma_n}.$$

But $\omega(z_0, E, R) \geq b/\Gamma(t)$ and $\omega(z_0, S(0, t), R) = \Gamma(\|z_0\|)/\Gamma(t)$, so

$$s(E) \geq \frac{n\sigma_n c^2 t^{n-1} b}{\Gamma(z_0)} \geq dt^{n-1},$$

where $d = n\sigma_n c^2 b/\Gamma(k)$. □

The following theorem is an analog of Harnack's inequality for arbitrary domains.

Theorem 2.5. *Let $D \subset B$ be an open set containing the origin, and let a, b, k , and m be positive constants with $m \geq a$. Suppose that for some point $z_0 \in D$ we have $\|z_0\| > k$ and $G_D(z_0, 0) \geq b$. Then there is a constant $c = c(a, k, m, n) > 0$ with the following property: for any positive sph function u on D which is harmonic and less than or equal to m on $D \cap B(0, k)$ and whose boundary values on $\partial D \cap B(0, k)$ are equal to 0, we have $u(z_0) \geq cb$ when $u(0) \geq a$.*

Proof. When $n > 2$ we consider the cone $C(\alpha) \subset \mathbb{R}^n$, which in polar coordinates (r, ϕ) is described as the set of points where $\phi \leq \alpha$. Let $C(\alpha, r) = C(\alpha) \cap \overline{B}(r, k/2)$. Using Lemma 2.4 we find $d = d(k/2, b/2, n)$ and take $\alpha > 0$ such that $s(C(\alpha) \cap S) = d$. If $B' = B(0, k/2)$ then the function $\psi(r) = \omega(0, C(\alpha, r), B')$ is a decreasing function for $0 < r < k$, and $\psi(r)$ converges to 1 as $r \rightarrow 0+$. The latter statement follows from the fact that by Theorem 8.27 of [H] the origin is a regular boundary point for the domain $B' \setminus C(\alpha, 0)$.

When $n = 2$ we take $\psi(r) = \omega(0, I_r, B')$, where $I_r = [r, k/2]$ is the interval on the real axis. In this case also $\psi(r) \rightarrow 1$ as $r \rightarrow 0+$.

Let $r = r(a, k, m, n) > 0$ be the maximal number such that $r \leq k/2$ and

$$(1) \quad \frac{a\psi(r)}{4} + m(1 - \psi(r)) \leq \frac{a}{2}.$$

Fix a sph function u on D satisfying the conditions of the theorem. Let $E \subset D$ be the set of points x where $u(x) \leq a/4$, and let $E_t = E \cap S(0, t)$. In the first case we assume that

$$\omega(z_0, E_t, D) < \frac{b}{2\Gamma(t)}$$

for some $r \leq t < k$. Then by Lemma 2.1

$$\omega(z_0, S(0, t) \setminus E_t, D) \geq \omega(z_0, S(0, t), D) - \omega(z_0, E_t, D) \geq \frac{b}{2\Gamma(t)},$$

and by Lemma 2.3

$$u(z_0) \geq \frac{ab}{8\Gamma(t)} \geq \frac{ab}{8\Gamma(r)}.$$

Thus in this case the theorem holds with $c = a/8\Gamma(r)$.

In the second case we assume that our previous assumption is not true, i.e.,

$$\omega(z_0, E_t, D) \geq \frac{b}{2\Gamma(t)}$$

for all $r \leq t < k$. Then the sets E_t are non-empty. By Lemma 2.4, $s(E_t) \geq dt^{n-1}$ when $n > 2$. Let $E' = E \cap \overline{B}(r, k/2)$. By Corollary 2 of [BT], $\omega(0, E', B') \geq \psi(r)$ when $n > 2$, and the same inequality holds for $n = 2$ by the Milloux–Schmidt inequality ([N], Theorem 1, p. 107). Consider the domain $D' = B' \setminus E'$ and the harmonic function v on D' with boundary values equal to $a/4$ on $E' \cap B'$ and to m on $\partial B' \setminus E'$. Clearly $u \leq v$ on D' and

$$v(0) \leq \frac{a}{4}\omega(0, E', B') + m(1 - \omega(0, E', B')) \leq \frac{a}{4}\psi(r) + m(1 - \psi(r)).$$

By inequality (1) $u(0) \leq a/2$ and we have a contradiction, which proves that the second case cannot occur and the theorem is proved. \square

In the next section we will need the following fact.

Corollary 2.6. *Let D be an open set in B containing the origin, and let $a > 0$. Fix $z_1, z_2 \in D$ and let $r = \min\{\|z_1\|, \|z_2\|\} > 0$. There exists a constant $c = c(a, r, n) > 0$ such that $G_D(z_1, z_2) \geq c$ if $\min\{G_D(0, z_1), G_D(0, z_2)\} \geq a$.*

Proof. In Theorem 2.5 let $u(x) = G_D(x, z_2)$ and $k = r/2$. Then u is harmonic on $D \cap B(0, k)$, its boundary values are equal to 0 on $\partial D \cap B(0, k)$, and u does not exceed some constant m depending only on k on $D \cap B(0, k)$. Since $u(0) \geq a$ and $G_D(0, z_1) \geq a$, by the previous theorem $u(z_1) = G_D(z_1, z_2) \geq c = c(a, r, n)$. \square

3. HARMONIC FUNCTIONS AND RESTORING COVERINGS

A positive Borel measure μ on a compact set $X \subset \mathbb{R}^n$ is called a Jensen measure with barycenter $z \in X$ if

$$(2) \quad u(z) \geq \int_X u(x) \mu(dx)$$

for every continuous sph function u defined on a neighborhood of X .

Since every sph function on an open set D is the limit of an increasing sequence of continuous sph functions on a set $G \subset D$, we see that subaveraging property (2) holds for all sph functions. Clearly $\mu(X) = 1$ and

$$u(z) = \int_X u(x) \mu(dx),$$

when u is harmonic on a neighborhood of X .

We denote by $J_z(X)$ the set of all Jensen measures on X with barycenter z . Clearly this set is convex and weak-* compact. Let $J(X)$ be the set of all Jensen measures on X . This set is also weak-* compact.

We denote by $h(X)$ the *real* linear space of continuous functions on X which can be uniformly approximated by functions harmonic in neighborhoods of X . We say that a continuous function h on X is harmonic if

$$h(z) = \int_X h(x) \mu(dx)$$

for every $z \in X$ and $\mu \in J_z(X)$. This definition is completely justified by the following theorem.

Theorem 3.1. *A continuous function f on X belongs to $h(X)$ if and only if it is harmonic on X .*

Proof. Suppose that there are harmonic functions u_j defined on neighborhoods D_j of X and converging to f uniformly on X . Then for every $z \in X$ and $\mu \in J_z(X)$ we have

$$f(z) = \lim u_j(z) = \lim \int_X u_j(x) \mu(dx) = \int_X f(x) \mu(dx).$$

To prove the converse statement we extend f continuously on \mathbb{R}^n as \tilde{f} . Let D_j be the j^{-1} -neighborhood of X and μ_x^j the harmonic measure on D_j relative to x . The function

$$u_j(x) = \int_{\partial D_j} \tilde{f}(y) \mu_x^j(dy)$$

is harmonic on D_j .

Suppose that there are points $z_j \in X$ and a $\delta > 0$ such that $|u_j(z_j) - f(z_j)| > \delta$ for all j . Switching if necessary to a subsequence, we may assume that $\lim z_j = z$ and that the measures $\mu_j = \mu_{z_j}^j$ weak-* converge to a measure μ . If v is a continuous sph function in a neighborhood of X , then v is sph on D_j when j is sufficiently large. Therefore,

$$v(z_j) \geq \int_{\partial D_j} v(x) \mu_j(dx).$$

Thus

$$v(z) \geq \int_X v(x) \mu(dx)$$

and $\mu \in J_z(X)$. Since the measures μ_j weak-* converge to μ , the sequence of

$$u_j(z_j) = \int_{\partial D_j} \tilde{f}(y) \mu_j(dy)$$

converges to

$$\int_X \tilde{f}(y) \mu(dy) = f(z).$$

But $f(z_j)$ also converges to $f(z)$, and this contradiction proves the theorem. \square

Let X be a compact set in \mathbb{R}^n . We define the harmonic measure of a closed set $E \subset \mathbb{R}^n$ relative to $x \in X$ as

$$\omega(x, E, X) = \lim_{\epsilon \rightarrow 0} \omega(x, E, X_\epsilon),$$

where $X_\epsilon = \{x \in \mathbb{R}^n : \text{dist}(x, X) < \epsilon\}$ is the open ϵ -neighborhood of X .

Lemma 3.2. *If E is a closed set in X , then*

$$\omega(z, E, X) = \inf_V \sup_\mu \mu(V),$$

where μ runs through all $J_z(X)$ and V through all open sets containing E .

Proof. Let us fix some $\epsilon > 0$ and take an open set $V \supset E$. Then

$$\mu(V) \leq \int_X \omega(x, \overline{V}, X_\epsilon) \mu(dx) \leq \omega(z, \overline{V}, X_\epsilon)$$

for all $\mu \in J_z(X)$. Hence

$$\omega(z, E, X) \geq \inf_V \sup_\mu \mu(V).$$

On the other hand, for the same V choose an open set $W \subset V$ containing E and a sequence of positive ϵ_j converging to 0 such that the sequence of harmonic measures μ_j relative to $X_{\epsilon_j} \setminus \overline{W}$ and z weak-* converges to some measure μ . Then

$$\mu_j(V) \geq \mu_j(\overline{W}) = \omega(z, \overline{W}, X_{\epsilon_j}) \geq \omega(z, E, X)$$

and $\mu(V) \geq \omega(z, E, X)$. If u is a continuous sph function in a neighborhood Z of X , then

$$u(z) \leq \int_{\mathbb{R}^n} u(x) \mu_j(dx)$$

when $\overline{X}_{\epsilon_j} \subset Z$. Since μ is the weak-* limit of μ_j , the latter inequality holds for μ also, and this means that μ is a Jensen measure on X with barycenter z . Thus

$$\sup_{\mu} \mu(V) \geq \omega(z, E, X)$$

for every open set $V \supset E$ when μ runs through all $J_z(X)$. \square

For a point $z \in X$ we consider the set

$$I_X(z) = I(z) = \{x \in X : \omega(z, \overline{B}(x, r), X) > 0 \text{ for all } r > 0\}.$$

By Lemma 3.2 if $\mu \in J_z(X)$, then $\text{supp } \mu \subset I(z)$. The set $I(z)$ is closed and $z \in I(z)$, so the sets $I(z)$ form a covering of X . The presence of points other than z in $I(z)$ can be deduced from

Theorem 3.3. *Let X be a compact set in \mathbb{R}^n and $z \in X$. The following statements are equivalent:*

- (1) $I(z) = \{z\}$.
- (2) If $E_r = X \setminus B(z, r)$, then $\omega(z, E_r, X) = 0$ for all $r > 0$.
- (3) The complement Y of X in \mathbb{R}^n is nonthin at z .

Proof. (1) \Rightarrow (2). Suppose that $\omega(z, E_r, X) > 0$ for some $r > 0$. The harmonic measure $\omega(z, A, X)$ is subadditive in A and, therefore, for any finite cover of E_r by balls B_j^1 of radius $r_1 = r/2$ with centers in E_r the harmonic measure at z of at least one ball, say B_1^1 , is greater than 0. Covering B_1^1 by balls of radius $r_2 = r_1/2$, choosing a ball with positive harmonic measure at z , and continuing this process at infinity, we get a sequence of points $x_j \in E_r$ such that harmonic measures at z of balls of radii $r_j = 2^{-j}r$ with centers at x_j are positive. Clearly any limit point of this sequence is in $I(z)$, and hence $I(z) \neq \{z\}$. Thus, $\omega(z, E_r, X) = 0$.

(2) \Rightarrow (3). Suppose that Y is thin at z . By [H, Theorem 10.3] we can find a positive sph function u on a ball $B_1 = B(z, r)$ such that $u(x) \geq 1 > u(z) = a$ for $x \in Y \cap B_1$. The function $v = 1 - u$ is subharmonic and less than 1 on B_1 , negative on $Y \cap B_1$, and $v(z) = 1 - a > 0$. Therefore, any positive sph function on X_ϵ , which is greater than 1 on $E_{r/2}$, is greater than v on X_ϵ . Hence, $\omega(z, E_r, X_\epsilon)$ and, consequently, $\omega(z, E_r, X)$ are greater than $1 - a > 0$.

(3) \Rightarrow (2). Suppose that Y is nonthin at z . Let D_j , $j = 1, 2, \dots$, be a sequence of decreasing regular domains such that $\bigcap \overline{D_j} = X$. Functions $u_j(x) = \omega(x, E_r, D_j)$ are harmonic on $D_j \cap B(z, r)$ and equal to 0 on $\partial D_j \cap B(z, r)$. If we let $u_j(x) = 0$ for $x \in B(z, r) \setminus \overline{D_j}$, then u_j will be subharmonic on $B(z, r)$. Since the sequence of u_j decreases, its limit u exists and is a subharmonic function on $B(z, r)$. By [H, Theorem 10.3] $\limsup_{x \rightarrow z, x \in Y} u(x) = u(z)$. But if $x \in Y \cap B(z, r)$, then $x \in B(z, r) \setminus \overline{D_j}$ when j is large and, therefore, $u_j(x) = 0$. So $u(x) = 0$ and $u(z) = \omega(z, E_r, X) = 0$.

(2) \Rightarrow (1). For arbitrary $x \neq z$ in X find positive t and r such that $B(x, t) \subset \mathbb{R}^n \setminus B(z, r)$. Then $\omega(z, \overline{B}(x, t), X) = 0$ and $x \notin I(z)$. Thus $I(z) = \{z\}$. \square

The following theorem shows that $\{I(z)\}$ is a restoring covering for $h(X)$.

Theorem 3.4. *A continuous function f belongs to $h(X)$ if and only if it belongs to $h(I(z))$ for all $z \in X$.*

Proof. The “only if” part of the theorem is trivial. Suppose that a continuous function $f \in h(I(z))$ for all $z \in X$. If $\mu \in J_z(X)$, then by Lemma 3.2 $\text{supp } \mu \subset I(z)$.

If $u(x)$ is a continuous sph function defined on a neighborhood D of $I(z)$, then for every set $G \subset D$ containing $I(z)$ there is a sph function v on \mathbb{R}^n such that $v \equiv u$ on G . To get such an extension one may assume that the open set $\mathbb{R}^n \setminus \overline{G}$ is regular and take a function

$$v(x) = \min(u(x), \alpha(V(x) - \beta)),$$

where $V(x)$ is the capacitary potential of \overline{G} (see [H, p. 138]) and α, β are appropriate constants. Then

$$\int_X u(x) \mu(dx) = \int_X v(x) \mu(dx) \leq u(z).$$

Thus $\mu \in J_z(I(z))$. Therefore,

$$f(z) = \int_X f(x) \mu(dx)$$

and by Theorem 3.1 the function $f \in h(X)$. \square

Green functions $G_{X_\epsilon}(x, y)$ decrease as ϵ decreases to 0 and, hence, converge to the function $G_X(x, y)$, which we will call the Green function on compacta. This function is, generally, not lower but upper semicontinuous and symmetric, i.e., $G_X(x, y) = G_X(y, x)$. The following theorem states that the properties of harmonic measures proved in Section 2 stay true when the open set D is replaced by a compact set X . From now on, we will refer to theorems from Section 2 when we need an analogous result for compacta.

Theorem 3.5. *Lemma 2.1, Theorem 2.2, and Corollary 2.6 stay true when the open set D in their statements is replaced by a compact set $X \subset B$ containing the origin.*

Proof. We may assume that open sets $X_\epsilon \subset B$. For proofs of the new versions of the listed results it is enough to note that the numbers $\omega(x, \overline{B}(0, t), X_\epsilon)$ and $G_{X_\epsilon}(x, 0)$ decrease to $\omega(x, \overline{B}(0, t), X)$ and $G_X(x, 0)$, respectively, as ϵ decreases to 0. \square

We say that $x \sim y$ for points $x, y \in X$ if $G_X(x, y) > 0$. Let us introduce the set $Q(z) = \{x \in X : G_X(x, z) > 0\}$.

The following theorem describes the geometry of the sets $Q(z)$ and $I(z)$, and its first three parts plus Theorem 3.4 provide the proof for Theorem 1.1.

Theorem 3.6. *Let X be a compact set in \mathbb{R}^n and $z \in X$.*

- (1) *The set $I(z)$ is the closure of the set $Q(z)$.*
- (2) *The relation \sim is an equivalence relation.*
- (3) *(The maximum principle) If $f \in h(X)$ and $f(y) = \sup_{x \in I(z)} f(x)$, for some $y \in Q(z)$, then f is constant on $I(z)$.*
- (4) *Either $Q(z) = \{z\}$ or $m(Q(z)) > 0$.*

Proof. 1) We may assume that the diameter of X is less than 1. Hence some translation moves any point of X into the origin leaving X in B . If $G_X(x, z) = a > 0$ then $G_X(z, x) = a$, and by Lemma 2.1 $\omega(z, \overline{B}(x, t), X) \geq a/\Gamma(t)$ for all sufficiently small t . So $Q(z) \subset I(z)$.

If $x \in I(z)$, then by Theorem 2.2 every neighborhood of x contains points y where $G_X(y, z) > 0$. So $I(z) \subset \overline{Q(z)}$.

2) We need to prove the transitivity, which follows immediately from Corollary 2.6.

3) By (2) $Q(z) = Q(y)$, so we may assume that $y = z$. Suppose that $x \in I(z)$ and $f(x) < f(z)$. Take a ball $B' = B(x, r)$ such that $f(y) < f(z)$ when $y \in X \cap B'$. Since $\omega(z, B', X) > 0$, by Lemma 3.2 there is a measure $\mu \in J_z(X)$ such that $\mu(B') > 0$. Then

$$f(z) = \int_X f(y) \mu(dy) < f(z),$$

and this contradiction proves part 3).

4) If $Q(z)$ contains more than one point, then by Lemma 2.1 and Theorem 2.2 every neighborhood of these points has positive measure. \square

Theorem 3.4 and the first and fourth part of the previous theorem yield the Hartogs–Rosenthal theorem.

Theorem 3.7. *If the measure of a compact set $X \subset \mathbb{R}^n$ is zero, then $h(X) = C_{\mathbb{R}}(X)$.*

The theorem above provides only a sufficient criterion for $h(X) = C_{\mathbb{R}}(X)$. The following theorem does it in the “necessary and sufficient” form. Statements (3) and (4) are due, respectively, to M. V. Keldysh [Ga] and D. Khavinson [Kh].

Theorem 3.8. *For a compact set $X \subset \mathbb{R}^n$ the following statements are equivalent:*

- (1) $h(X) = C_{\mathbb{R}}(X)$;
- (2) $I(z) = \{z\}$ for all $z \in X$;
- (3) $\mathbb{R}^n \setminus X$ is nonthin at all points of X ;
- (4) $f(x) = \|x\|^2$ is in $h(X)$.

Proof. (1) and (2) are equivalent by Theorem 3.4. (2) and (3) are equivalent by Theorem 3.3. Clearly (1) implies (4). If (4) is true, then $-||x - z||^2 \in h(X)$ for all $z \in X$. Hence by the third part of Theorem 3.6 $Q(z) = \{z\}$. So $I(z) = \{z\}$ and (1) is also true. \square

Definition 3.9. A compact set $X \subset \mathbb{R}^n$ is called a *Jensen set* if $X = I(z)$ for some point $z \in X$.

The simplest example of a Jensen set is the closed unit ball in \mathbb{R}^n . Also, if Y is a compact set, $z \in Y$, and $X = I(z)$, then X is a Jensen set. Indeed, $J_z(X) = J_z(Y)$ because all measures from $J_z(Y)$ are supported by X . Hence by Lemma 3.2 $\omega(z, E, X) = \omega(z, E, Y)$ for all closed sets $E \subset X$. This means that $I_X(z) = I_Y(z) = X$.

The following example provides a non-trivial Jensen set whose interior is empty.

Example 3.10. Take, in B , a countable dense set $A = \{z_1, z_2, \dots\}$ which does not contain the origin and a point $x \notin A$ with $\|x\| > 1/2$. Let $B_j = B(x, 1/2^j)$, $j = 1, 2, \dots$. We choose $t_1 > 0$ such that $x \notin B(z_1, t_1)$ and $\omega(0, \overline{B_1}, \overline{D_1}) > \omega(0, \overline{B_1}, \overline{B})/2$, where $D_1 = \overline{B} \setminus B(z_1, t_1)$. If t_1, \dots, t_{n-1} have been chosen, choose $t_n > 0$ such that $x \notin B(z_n, t_n)$ and $\omega(0, \overline{B_j}, \overline{D_n}) > \omega(0, \overline{B_j}, \overline{D_j})/2$, where $j = 1, \dots, n$ and $D_j = D_{j-1} \setminus B(z_j, t_j)$. Let $Y = \bigcap D_j$. Then $x \in Y$. For every $\epsilon > 0$ the compact set $D_j \subset Y_\epsilon$ when j is sufficiently large and, therefore, $\omega(0, B_j, Y) > 0$ for all j . Thus $x \in I_Y(0)$. Take $X = I_Y(0)$. The interior of X is empty and X is Jensen.

Now we are able to establish analytic properties of elements of a restoring covering. These elements are Jensen sets. From the maximum principle we get

Corollary 3.11. *If X is a Jensen set in \mathbb{R}^n , then every non-negative function $f \in h(X)$ equal to 0 on an open non-void set is equal to 0.*

As the following example shows, a point w in $I(z) \setminus Q(z)$ can have a non-trivial set $Q(w)$ when $X \subset \mathbb{R}^n$, $n \geq 3$.

Example 3.12. Let $\phi(x)$ be a positive continuous function on $(-\infty, 0)$ decreasing so fast that for the set

$$E = \overline{B}(0, 1) \setminus \{x < 0, y^2 + z^2 < \phi(x)\}$$

in $\mathbb{R}^3 = (x, y, z)$ the origin is an irregular point. Let

$$F = \overline{B}(0, 1) \cap \{x < 0, y^2 + z^2 \leq \frac{\phi(x)}{2}\}$$

and $X = E \cup F$. If $w_1 = (-1/2, 0, 0)$ and $w_2 = (1/2, 0, 0)$, then $0 \in I(w_1) \setminus Q(w_1)$, but $G_X(0, w_2) > 0$, so $w_2 \in I(0)$.

However for $n = 2$ we have

Theorem 3.13. *If X is a compact set in \mathbb{C} , $z \in X$ and $w \in I(z)$, then either $Q(z) = Q(w)$ or $Q(w) = \{w\}$.*

The proof of this theorem follows immediately from

Lemma 3.14. *Suppose that z is a point of a compact set $X \subset \mathbb{C}$, $w \in I(z)$, $y \notin Q(z)$, $y \neq w$. Then $G_X(w, y) = 0$.*

Proof. If $Q(z) \cap S(w, t) = \emptyset$ for some $t < r = \|z - w\|$, then $G_X(x, z) \equiv 0$ on $B(w, t)$ and, hence, $w \notin I(z)$. So $Q(z) \cap S(w, t) \neq \emptyset$ for $0 < t < r$. Let $s = \min\{r, \|w - y\|/2\}$ and $E = Q(z) \cap B(w, s)$.

Let G_ϵ be the Green function on X_ϵ . We fix $\delta > 0$ and for every $\epsilon > 0$ we consider the closure F_ϵ of the set

$$\{x \in B(w, s) \cap X_\epsilon : G_\epsilon(x, y) \leq \delta\}.$$

Since the union of F_ϵ , $\epsilon > 0$, contains E , by the Milloux-Schmidt inequality,

$$\lim_{\epsilon \rightarrow 0} \omega(w, F_\epsilon, B(w, s) \cap X_\epsilon) = 1.$$

So $\omega(w, F_\epsilon, B(w, s) \cap X_\epsilon) > 1 - \delta$ for some $\epsilon > 0$, and since $G_\epsilon(x, y)$ is less than some constant c on $B(w, s)$, we see that $G_\epsilon(w, y) \leq \delta(1 - \delta) + c\delta$. Thus $G_X(w, y) \leq \delta(1 - \delta) + c\delta$ for any $\delta > 0$, and $G_X(w, y) = 0$. \square

Definition 3.15. A Jensen set X is called a *Wermer set* if for every $z \in X$ either $I(z) = X$ or $I(z) = \{z\}$.

The previous theorem implies

Corollary 3.16. *Every Jensen set $X \subset \mathbb{C}$ is a Wermer set.*

Definition 3.17. A uniform algebra A on a compact set X is *analytic* if every function $f \in A$ equal to 0 on an open non-void set in X is equal to 0.

If X is a compact set in \mathbb{C} , we denote by $H(X)$ the *complex* uniform algebra of continuous functions on X which can be uniformly approximated on X by functions holomorphic in neighborhoods of X .

The following theorem is a simple consequence of the Two Constants Theorem.

Theorem 3.18. *If X is a Jensen set in \mathbb{C} , then $H(X)$ is analytic.*

Proof. Let $h \in H(X)$ be equal to 0 on a non-void open set $V \subset X$. We may assume that $X = I(z)$ and $z \in V$. There is a sequence of open sets $U_j \supset X$ and holomorphic functions h_j on U_j such that h_j converge uniformly to h on X . We may assume that $|h_j| \leq 1$ on U_j . If $w \in Q(z)$ then

$$\omega(w, \overline{V}, U_j) \geq \omega(w, \overline{V}, X) = a > 0.$$

Let b_j be the supremum of $|h_j|$ on \overline{V} . By the Two Constants Theorem $|h_j(w)| \leq b_j^a$, and therefore $h(w) = 0$ for $w \in Q(z)$. Since $Q(z)$ is dense in X , $h \equiv 0$. \square

This result above yields the following:

Corollary 3.19. *If a compact set X in \mathbb{C} has a nontrivial Jensen measure (not equal to δ_z), then X contains a nontrivial compact set Y with analytic algebra $H(Y)$.*

Proof. Let z be the barycenter of a nontrivial Jensen measure on X . Then $Y = I(z)$ is a nontrivial Jensen set. \square

It follows from [P] that a compact set $X \subset \mathbb{C}^n$ is Jensen (with respect to z) if and only if X is the cluster of a perfect sequence centered at z . Combining this result with Theorem 3.18, we get the following:

Corollary 3.20. *If a compact set $X \subset \mathbb{C}$ is the cluster of a perfect sequence of mappings of the unit disk into \mathbb{C} , then $H(X)$ is analytic.*

In \mathbb{C}^n the situation is more complicated. The Wermer example X is the cluster of a perfect sequence with analytic algebra $H(X)$. This follows from the proof of analyticity of $H(X)$ in [G]. But the compact set $Y = Y_1 \cup Y_2$, where $Y_1 = \{(z, w) \in \mathbb{C}^2 : |w| \leq 1, |z| = 1\}$ and $Y_2 = \{(z, w) \in \mathbb{C}^2 : w = 0, |z| \leq 1\}$ is also the cluster of a perfect sequence (see [P]). However, the algebra $H(Y)$ is not analytic because $f(z, w) = w$ is equal to 0 on Y_2 .

REFERENCES

- [BT] A. Baernstein II, B. A. Taylor, *Spherical rearrangements, subharmonic functions, and *-functions in n -space*, Duke Math. J., 43, (1976), pp. 245–268. MR **53**:5906
- [Gam] T. Gamelin, *Uniform Algebras*, Prentice-Hall, Inc., 1984. MR **53**:14137 (1st ed.)
- [Ga] P. M. Gauthier, *Uniform approximation*, In: Complex Potential Theory, Kluwer Acad. Publ., 1994, pp. 235–271. MR **96d**:30048
- [G] H. Goldmann, *An analytic algebra without analytic structure in the spectrum*, Ark. Mat., 27 (1989), 89–95. MR **90e**:46042
- [H] L. L. Helms, *Introduction to Potential Theory*, Wiley-Interscience, 1969. MR **41**:5638
- [Kh] D. Khavinson, *On uniform approximation by harmonic functions*, Mich. Math. J., 34 (1987), 465–473. MR **89a**:14026
- [M] R. McKissick, *A nontrivial normal sup algebra*, Bull. Amer. Math. Soc., 69 (1963), 391–395. MR **26**:4166
- [N] R. Nevanlinna, *Analytic functions*, Springer-Verlag, 1970. MR **43**:5003
- [P] E. A. Poletsky, *Analytic geometry on compacta in \mathbb{C}^n* , Math. Z., 222 (1996), 407–424. MR **97e**:32015
- [S] E. L. Stout, *The Theory of Uniform Algebras*, Bogden & Quigley, Inc., Tarrytown-on-Hudson, NY, 1971. MR **54**:11066
- [W] J. Wermer, *Polynomially convex hulls and analyticity*, Ark. Mat., 20 (1982), 129–135. MR **84b**:32021

DEPARTMENT OF MATHEMATICS, 215 CARNEGIE HALL, SYRACUSE UNIVERSITY, SYRACUSE, NEW YORK 13244