

ASYMPTOTIC FORMULAE WITH REMAINDER ESTIMATES FOR EIGENVALUE BRANCHES OF THE SCHRÖDINGER OPERATOR $H - \lambda W$ IN A GAP OF H

S. Z. LEVENDORSKIĬ

ABSTRACT. The Floquet theory provides a decomposition of a periodic Schrödinger operator into a direct integral, over a torus, of operators on a basic period cell. In this paper, it is proved that the same transform establishes a unitary equivalence between a multiplier by a decaying potential and a pseudo-differential operator on the torus, with an operator-valued symbol. A formula for the symbol is given.

As applications, precise remainder estimates and two-term asymptotic formulas for spectral problems for a perturbed periodic Schrödinger operator are obtained.

0.1. There are a number of articles on the discrete spectrum in a gap of the essential spectrum of the Schrödinger operator $H = -\Delta + V$ (see [1, 3, 4, 7, 8, 9, 12, 13, 17, 20] and the bibliography in [1, 3, 4]). The majority of them [1, 3, 4, 7, 8, 9, 21] deal with the existence of eigenvalue branches and with estimates or asymptotics for the number $N_{\pm}(t; H - E; W)$ of branches of $H \mp \mu W$ with $0 < \mu < t$ which cross the energy level $E \in \mathbb{R} - \sigma(H)$ (here W is a potential decaying at infinity). In [12, 13, 17], the principal term of the asymptotics of a series of eigenvalues accumulating to a boundary point \mathcal{E} of the essential spectrum was computed, without a remainder estimate.

For motivation from solid state physics (impurities in crystals), see [1] and the bibliography there.

In this paper, we obtain the principal term of the asymptotics with remainder estimate for $N_{\pm}(t; H - E; W)$; W is smooth and non-negative, and V is periodic.

The main idea of the paper is a reduction to a pseudodifferential operator (pdo) on a torus, with operator-valued symbol. For $N_{-}(t; H - E; W)$, further reductions allow one to obtain bounds in terms of the counting function of the spectrum of a pdo acting in sections of a finite-dimensional fibering over a torus. These bounds give the principal terms of the asymptotics as $t \rightarrow \infty$, with remainder estimates which are as sharp as those for the corresponding pdo; if for the latter an asymptotic formula with two terms of the asymptotics is known, then the same formula is valid for $N_{-}(t; H - E; W)$.

In [1], only the principal term of the asymptotics of $N_{-}(t; H - E; W)$ was found, without a remainder estimate, and W was assumed to stabilize to a positive spherically homogeneous potential; on the other hand, V was not necessarily periodic.

Received by the editors May 15, 1995 and, in revised form, December 9, 1995.

1991 *Mathematics Subject Classification.* Primary 35P20.

The author was supported in part by ISF grant RNH 000.

Note that the reduction to an operator with operator-valued symbol explains why the classical Weyl formula with the ordinary symbol predicts false asymptotics for $N_-(t; H; W)$ (as was observed in [1]): for the ordinary symbol under consideration, the uncertainty principle fails since derivatives of this symbol do not admit good estimates (so that the function h , which plays a crucial role in calculi of pseudodifferential operators [10, Chapter 18], is not small outside some compact), whereas the operator-valued symbol enjoys more favourable properties. Similar effects are well-known in the theory of degenerate elliptic operators—see, for example, the surveys [5, 15, 19].

The same reduction and the approximate spectral projection method for pseudodifferential operators with operator-valued symbols [16, Section 22] allows one to obtain the principal term of the asymptotics of $N_+(t; H - E; W)$ when W is not a short-range potential. The principal term of the asymptotics for a short-range potential W , was computed in [9], [3], where it was shown that it agreed with the classical Weyl formula; if W is not short-range, then the classical Weyl formula fails and analogues of strong and intermediate degeneration take place (for a discussion of the types of degenerate elliptic operators, see [5, 15, 19]).

The reduction to an operator on a torus (with a subsequent reduction to different operators) also gives asymptotic formulae with remainder estimates or even two terms of the asymptotics for a series of eigenvalues of $H + W$ accumulating to a boundary point \mathcal{E} of the essential spectrum of $H + W$; the corresponding results will be published elsewhere.

In [4], the asymptotics of $N_\pm(t; H - \mathcal{E}; W)$ was computed; this case has special subtle features.

Variational arguments allow one to deduce from the results of this paper the principal term of the asymptotics for non-smooth potentials—see e.g. [19, Section 11], where the general scheme (based on results by M.S.Birman and M.Z.Solomyak) is described.

0.2. Now we formulate conditions on V and W .

Let $\{\alpha^j\}_{j=1}^n$ be a basis in \mathbb{R}^n , and

$$\Gamma = \{x \in \mathbb{R}^n \mid x = \sum_{j=1}^n p_j \alpha^j, p_j \in \mathbb{Z}, j = 1, \dots, n\}$$

the lattice associated with $\{\alpha^j\}$.

Let V be a real-valued Γ -periodic function of the class $L_\infty(\mathbb{R}^n)$; the corresponding multiplier will also be denoted by V .

$H = -\Delta + V$ is an operator in $L_2(\mathbb{R}^n)$.

Let W satisfy the following conditions:

(a) There is $\rho \in (0, 1]$ such that

$$(0.1) \quad |(D^\alpha W)(x)| \leq C_\alpha W(x) \langle x \rangle^{-\rho|\alpha|}, \quad \forall \alpha,$$

where $\langle x \rangle = (1 + |x|^2)^{1/2}$, C_α are independent of x

(b) There are positive constants m_-, m, d, d_1 such that

$$(0.2) \quad d_1 \langle x \rangle^{-m_-} \leq W(x) \leq d \langle x \rangle^{-m}.$$

In the last subsections of this Introduction, we consider non-negative W and W having singularities.

0.3. We need the following construction which is usually used to describe $\sigma_{\text{ess}}(H) = \sigma(H)$ (see e.g. [18, Sect.XIII,16], and [19]).

Let $\{\beta^j\}_{j=1}^n$ be a basis dual to $\{\alpha^j\}_{j=1}^n$:

$$(\alpha^j, \beta^k) = 2\pi\delta_{jk},$$

and set

$$\Gamma^* = \{x \in \mathbb{R}^n \mid x = \sum_{j=1}^n p_j \beta^j, \ p_j \in \mathbb{Z}, \ j = 1, \dots, n\}.$$

Set

$$\Omega = \{x = \sum_{j=1}^n t_j \alpha^j \mid t_j \in [0, 1)\}, \quad \Omega^* = \{\theta = \sum_{j=1}^n \theta_j \beta^j \mid \theta_j \in [0, 1)\}.$$

We use the usual flat metrics on \mathbb{R}^n/Γ and $\mathbb{R}^{n^*}/\Gamma^*$; when we integrate or do local considerations we identify \mathbb{R}^n/Γ (resp. $\mathbb{R}^{n^*}/\Gamma^*$) with Ω (resp. Ω^*).

For fixed $\theta \in \Omega^*$, denote by $\mathcal{H}(\theta)$ an operator $-\Delta + V$ on Ω with the boundary conditions

$$u(x + \alpha^j) = \exp\{i(\alpha^j, \theta)\}u(x), \quad \partial_{t_j} u(x + \alpha^j) = \exp\{i(\alpha^j, \theta)\}\partial_{t_j} u(x)$$

($x, x + \alpha^j \in \partial\Omega, j = 1, \dots, n$). This is a self-adjoint semibounded (from below) operator with discrete spectrum. Let $E_1(\theta) < E_2(\theta) \leq \dots$ be its eigenvalues counted with multiplicity.

Introduce

$$\mathfrak{L} := \int_{\mathbb{R}^{n^*}/\Gamma^*} \oplus L_2(\Omega) \frac{d\theta}{\text{vol } \Omega^*}, \quad \mathcal{H} := \int_{\mathbb{R}^{n^*}/\Gamma^*} \oplus \mathcal{H}(\theta) \frac{d\theta}{\text{vol } \Omega^*}.$$

One can identify \mathfrak{L} with $L_2(\mathbb{R}^{n^*}/\Gamma^*; L_2(\Omega))$ ($\cong L_2(\Omega^*, d\theta/\text{vol } \Omega^*; L_2(\Omega, dx))$), where $d\theta$ and dx are the Lebesgue measures over Ω^* and Ω respectively, and \mathcal{H} with multiplication by the operator-valued function $\mathcal{H}(\cdot)$.

Define an operator $\mathcal{U} : L_2(\mathbb{R}^n) \rightarrow \mathfrak{L}$ by

$$(0.3) \quad (\mathcal{U}f)(x, \theta) = \sum_{\gamma \in \Gamma} \exp\{-i(\theta, \gamma)\}f(x + \gamma).$$

This is an isometry, and $\mathcal{U}H\mathcal{U}^* = \mathcal{H}$. Hence

$$\sigma(H) = \bigcup_{j \geq 1} \bigcup_{\theta \in \mathbb{R}^{n^*}/\Gamma^*} E_j(\theta).$$

0.4. Reduction to a pdo on a torus. In the paper, the computation of the spectral asymptotics is based on the following theorem, which will be proved in Section 1.

Theorem 0.1. *Let W satisfy (0.1). Then*

$$\mathcal{W} := \mathcal{U}W\mathcal{U}^* \in L_{\rho,0}^{-m}(\mathbb{R}^{n^*}/\Gamma^*; \text{End } L_2(\Omega))$$

(i.e. it is a pdo on $\mathbb{R}^{n^*}/\Gamma^*$, of Hörmander's class $L_{\rho,0}^{-m}$, with the symbol, call it w , taking values in $\text{End } L_2(\Omega)$), and w admits an asymptotic expansion

$$w(\xi) = W(-\xi) + \sum_{j=1}^n (\cdot)_j (\partial_j W)(-\xi) + \dots,$$

where $(\cdot)_j$ is a “multiplication by x_j ” operator in $L_2(\Omega)$.

If in addition W satisfies (0.2), then

$$\mathcal{W}^{-1} := \mathcal{U}W^{-1}\mathcal{U}^* \in L_{\rho,0}^{m-}(\mathbb{R}^{n^*}/\Gamma^*; \text{End } L_2(\Omega)),$$

and its symbol, call it w^{-1} , admits an asymptotic expansion

$$(0.4) \quad w^{-1}(\xi) = W(-\xi)^{-1} + \sum_{j=1}^n (\cdot)_j (\partial_j W^{-1})(-\xi) + \dots$$

For a detailed formulation and a formula for the full asymptotic expansion of $w^{\pm 1}$, see Section 1.

0.5. Asymptotics of $N_-(t; H - E; W)$: a reduction to a spectral problem for a pdo acting in sections of a finite-dimensional fibering over a torus.

Let $(\mathcal{E}^-, \mathcal{E}^+)$ be a gap in $\sigma(H)$, and fix $E \in (\mathcal{E}^-, \mathcal{E}^+)$. Let $\mathcal{E}^- > -\infty$; then there is $p \geq 1$ such that

$$(0.5) \quad E_1(\theta) < E_2(\theta) \leq \dots \leq E_p(\theta) < E < E_{p+1}(\theta) \leq \dots, \quad \forall \theta \in \mathbb{R}^{n^*}/\Gamma^*.$$

Let $\mathcal{P}_1(\theta)$ stand for a spectral projection $P_{(-\infty, E)}(\mathcal{H}(\theta))$, and set $\mathcal{L}(\theta) = \text{Ran } \mathcal{P}_1(\theta)$ and $\mathcal{L} = \bigcup_{\theta \in \mathbb{R}^{n^*}/\Gamma^*} \mathcal{L}(\theta)$. Due to (0.5), $\mathcal{P}_1 \in C^\infty(\mathbb{R}^{n^*}/\Gamma^*; \text{End } L_2(\Omega))$ and $\dim \text{Ran } \mathcal{P}_1(\theta) = p$ for all $\theta \in \mathbb{R}^{n^*}/\Gamma^*$. Hence, \mathcal{L} is a p -dimensional fibering over $\mathbb{R}^{n^*}/\Gamma^*$, which inherits a structure of an Hermitian fibering from $L_2(\mathbb{R}^{n^*}/\Gamma^*; L_2(\Omega))$. Clearly, the operators

$$u(\cdot, \theta) \mapsto \mathcal{P}_1(\theta)(E - \mathcal{H}(\theta))^s u(\cdot, \theta), \quad s \in \mathbb{R},$$

and

$$u(\cdot, \theta) \mapsto \mathcal{P}_1(\theta)(\mathcal{W}^{-1}u)(\cdot, \theta)$$

act in $C^\infty(\mathbb{R}^{n^*}/\Gamma^*; \mathcal{L})$; we denote them by $(E - \mathcal{H})_1^s$ and \mathcal{W}_1^{-1} , respectively.

\mathcal{W}_1^{-1} is a pdo due to Theorem 0.1, and $(E - \mathcal{H})_1^s$ is locally a multiplier by a matrix-valued function which is positive-definite due to (0.5). Hence, $\mathcal{W}_1^{-1} := (E - \mathcal{H})_1^{1/2} \mathcal{W}_1^{-1} (E - \mathcal{H})_1^{1/2}$ is a pdo acting in $C^\infty(\mathbb{R}^{n^*}/\Gamma^*; \mathcal{L})$. To calculate its local symbols, take an open set $U \subset \mathbb{R}^{n^*}/\Gamma^*$. If $\text{diam } U$ is small then there exists an orthonormal basis $\{u_j(\cdot, \theta)\}_{j=1}^p$ in $\text{Ran } \mathcal{P}_1(\theta)$, $\theta \in U$, with $u_j \in C^\infty(\bar{U}; L_2(\Omega))$. Thus, we can define a trivialization $\chi : \mathcal{L}|_U \rightarrow U \times \mathbb{C}^p$ by

$$\chi^{-1}((f_1(\theta), \dots, f_p(\theta))) = \sum_{j=1}^p u_j(\cdot, \theta) f_j(\theta).$$

By using (0.4) and the composition theorem for pdo (see Theorem 5.8), one obtains an asymptotic expansion for a local symbol of \mathcal{W}_1^{-1} , call it $w_{11,U,\chi}^{-1}$:

$$(0.6) \quad w_{11,U,\chi}^{-1} = w_{11,U,\chi}^{-1,0} + w_{11,U,\chi}^{-1,-1} + \dots,$$

with each term decaying faster than the previous one as $\xi \rightarrow \infty$. For detailed version of (0.6) see Section 2; here we need formulae for the first two terms only:

$$w_{11,U,\chi}^{-1,0}(\theta, \xi) = (E - \mathcal{H}(\theta))_{11}^1 W(-\xi)^{-1},$$

where

$$(E - \mathcal{H}(\theta))_{11}^r = [\langle u_j(\cdot, \theta), (E - \mathcal{H}(\theta))^r u_l(\cdot, \theta) \rangle_{L_2(\Omega)}]_{j,l=1}^p,$$

and

$$w_{11,U,\chi}^{-1,-1}(\theta, \xi) = [w_{11,U,\chi;l;j}^{-1,-1}(\theta, \xi)]_{l,j=1}^p,$$

$$w_{11,U,\chi;l;j}^{-1,-1}(\theta, \xi) = \sum_{r=1}^n (iD_r W^{-1})(-\xi) \langle (E - \mathcal{H}(\theta))_1^{1/2} u_l(\cdot, \theta), \\ (-D_{\theta_r} + (\cdot)_r)((E - \mathcal{H}(\theta))_1^{1/2} u_j(\cdot, \theta)) \rangle_{L_2(\Omega)},$$

where $D_j = -i\partial_j$.

It follows from (0.6) and (0.2) that \mathcal{W}_{11}^{-1} is a symmetric hypoelliptic pdo of the Hörmander class $H_{\rho,0}^{m-,m}(\mathbb{R}^{n^*}/\Gamma^*; \mathcal{L})$, with symbol positive definite for large $|\xi|$. Hence, the closure of \mathcal{W}_{11}^{-1} , call it $\overline{\mathcal{W}_{11}^{-1}}$, is a semibounded (from below) self-adjoint operator with discrete spectrum (see Theorem 5.12).

For $C \in \mathbb{R}$, set $W_C^{-1}(x) = W(x)^{-1} - CW(x)^{-1}\langle x \rangle^{-2\rho}$, and construct $\mathcal{W}_{11,C}^{-1}$ in the way \mathcal{W}_{11}^{-1} was constructed (but starting from W_C^{-1} instead of W^{-1}). Theorem 0.1 gives for the symbol of $\mathcal{W}_{11,C}^{-1}$ an asymptotic expansion (0.6) with the same first terms (only the tail differs from the one for \mathcal{W}_{11}^{-1}); hence the above consideration shows that $\overline{\mathcal{W}_{11,C}^{-1}}$, the closure of $\mathcal{W}_{11,C}^{-1}$, is semibounded from below with discrete spectrum.

Denote by $N(t; A)$ the counting function of the spectrum of a semibounded (from below) operator A .

Theorem 0.2. *Let (0.1), (0.2) and (0.5) hold.*

Then there exists $C > 0$ such that

$$(0.7) \quad N(t; \overline{\mathcal{W}_{11}^{-1}}) \leq N_-(t; H - E; W) \leq N(t; \overline{\mathcal{W}_{11,C}^{-1}}).$$

0.6. Asymptotics of $N_-(t; H - E; W)$: “hypoelliptic case”. The first two terms of the asymptotic expansion (0.6) for the symbol of $\mathcal{W}_{11,C}^{-1}$ are independent of C ; hence results for hypoelliptic pdo (see Theorem 6.4) give the same asymptotic formulae for the left- and right-most parts of (0.7) (“the same” means that only a constant in the O -term (or o -term) depends on C). Therefore, the same formulae hold for $N_-(t; H - E; W)$.

In a general “hypoelliptic case” (0.1), (0.2), we have the following theorem.

Theorem 0.3. *Let (0.1), (0.2) and (0.5) hold, and let*

$$c_-(t, H - E, W) \\ = (2\pi)^{-n} \sum_{j=1}^p \text{meas}\{(\theta, \xi) \in \Omega^* \times \mathbb{R}^n \mid 1 + tW(-\xi)(E_j(\theta) - E)^{-1} < 0\}$$

satisfy a Tauberian condition: for any $\epsilon > 0$, there is $\omega > 0$ such that

$$(0.8) \quad c_-(t \pm t^{1-\epsilon}, H - E, W) - c_-(t, H - E, W) \\ = O(c_-(t, H - E, W)t^{-\omega}), \quad \text{as } t \rightarrow +\infty.$$

Then there exists $\omega > 0$ such that as $t \rightarrow +\infty$,

$$(0.9) \quad N_-(t; H - E; W) = c_-(t, H - E, W)(1 + O(t^{-\omega})).$$

For the detailed proof, see Section 2.

0.7. Asymptotics of $N_-(t; H - E; W)$: “elliptic case”. Let W be “elliptic”, i.e. let it admit a decomposition

$$(0.10) \quad W(x) = W_{-m}(x) + O(|x|^{-m-1}), \quad \text{as } |x| \rightarrow +\infty,$$

where $W_{-m} \in C^\infty(\mathbb{R}^n \setminus 0; \mathbb{R}_+)$ is positively homogeneous of degree $-m < 0$. Then (0.6) gives

$$w_{11,U,\chi}^{-1}(\theta, \xi) = (E - \mathcal{H}(\theta))_{11}^1 W_{-m}(-\xi)^{-1} + O(|\xi|^{m-1}), \quad \text{as } |\xi| \rightarrow +\infty,$$

and the well-known results for elliptic pdo [11, 19] give the following theorem.

Theorem 0.4. *Let (0.5) hold, let $(0 <) W \in L_\infty$ satisfy (0.10) with $m > 0$, and let $W^{-1} \in L_{\infty,loc}(\mathbb{R}^n)$.*

Then

$$(0.11) \quad N_-(t; H - E; W) = c_-(H - E; W_{-m}) t^{n/m} + O(t^{(n-1)/m}),$$

where

$$\begin{aligned} c_-(H - E; W_{-m}) \\ = (2\pi)^{-n} n^{-1} \sum_{j=1}^p \int_{S_{n-1}} (W(x)_{-m})^{n/m} dS(x) \int_{\mathbb{R}^{n^*}/\Gamma^*} (E - E_j(\theta))^{-n/m} d\theta. \end{aligned}$$

For details, see Section 2.

Now, let

$$(0.12) \quad W(x) = W_{-m}(x) + W_{-m-1}(x) + O(|x|^{-m-1-\epsilon}), \quad \text{as } |x| \rightarrow +\infty,$$

where $W_j \in C^\infty(\mathbb{R}^n \setminus 0)$ are positively homogeneous of degree j ($j = -m, -m-1$), $\epsilon > 0$, and W_{-m} is positive. Then (0.6) implies that $\mathcal{W}_{11,C}^{-1}$ is (modulo terms of order less than $m-1$) a positive definite classical elliptic pdo of order m , with the principal symbol

$$(0.13) \quad w_{11,U,\chi;m}^{-1}(\theta, \xi) = (E - \mathcal{H}(\theta))_{11}^1 W_{-m}(-\xi)^{-1},$$

and the next term of the asymptotic expansion of the symbol defined locally by

$$(0.14) \quad w_{11,U,\chi;m-1}^{-1}(\theta, \xi) = -(E - \mathcal{H}(\theta))_{11}^1 (W_{-m}^{-2} W_{-m-1})(-\xi) + w'_{11,U,\chi}(\theta, \xi),$$

where

$$\begin{aligned} w'_{11,U,\chi}(\theta, \xi) &= [w'_{11,U,\chi;l_j}(\theta, \xi)]_{l,j=1}^p, \\ w'_{11,U,\chi;l_j}(\theta, \xi) &= \sum_{r=1}^n (iD_r W_{-m}^{-1})(-\xi) \\ &\quad \times \langle (E - \mathcal{H}(\theta))_1^{1/2} u_l(\cdot, \theta), (-D_{\theta_r} + (\cdot)_r) ((E - \mathcal{H}(\theta))_1^{1/2} u_j(\cdot, \theta)) \rangle_{L_2(\Omega)}. \end{aligned}$$

For details, see Section 2.

If one knows the first two terms of the asymptotic expansion of the symbol of a positive definite pdo $\mathcal{W}_{11,C}^{-1}$ acting in sections of a finite-dimensional fibering over a compact manifold without boundary, one can compute both the principal and subprincipal symbols of a power $\mathcal{W}_1 := (\mathcal{W}_{11}^{-1})^{1/m}$:

$$(0.15) \quad w_{1,U,\chi;p} = (w_{11,U,\chi;m}^{-1})^{1/m},$$

$$(0.16) \quad w_{1,U,\chi,s} = w_{1,U,\chi;0} + \frac{i}{2} \sum_{|\alpha|=1} \partial_\theta^\alpha \partial_\xi^\alpha w_{1,U,\chi;p},$$

where $w_{1,U,\chi;0}$ is the term of order zero in the asymptotic expansion of the symbol of \mathcal{W}_1 ; for the formula for $w_{1,U,\chi;0}$, see [20]. Formulae (0.13)-(0.16) taken together provide complete information about the principal and subprincipal symbols of \mathcal{W}_1 . In particular, the eigenvalues of the principal symbol of \mathcal{W}_1 are given by

$$\mu_j(\theta, \xi) = (E - E_j(\theta))^{1/m} (W_{-m}^{-1}(-\xi))^{1/m}, \quad j = 1, \dots, p.$$

Under certain non-periodicity condition on bicharacteristics of μ_j , there exist general two-term asymptotic formulas for the counting function of the spectrum of elliptic operators [11] (see also [19]), but in the case of non-constant multiplicities of μ_j , the result is only announced in [11].

By applying results of [11] to (0.7), we obtain the following theorem (modulo reservations in the case of non-constant multiplicities of μ_j).

Theorem 0.5. *Let (0.5) hold, let $(0 <) W \in L_\infty(\mathbb{R}^n)$ satisfy (0.12) with $m > 0$, let $W^{-1} \in L_{\infty,loc}(\mathbb{R}^n)$, and let the bicharacteristics of μ_j , $j = 1, \dots, p$, satisfy a non-periodicity condition of [11, 19].*

Then

$$(0.17) \quad \begin{aligned} N_-(t; H - E; W) &= c_-(H - E; W_{-m}) t^{n/m} \\ &+ c_1(H - E; W_{-m}; W_{-m-1}) t^{(n-1)/m} + o(t^{(n-1)/m}). \end{aligned}$$

For a (rather complicated) formula for $c_1(H - E; W_{-m}; W_{-m-1})$ in terms of $w_{1,p}$ and $w_{1,s}$, see [11].

0.8. Asymptotics of $N_+(t; H - E; W)$. For the case W a short-range potential, the principal term of asymptotics was calculated in [9, 3]; V was not necessarily periodic. In [9], $W(x) \leq C\langle x \rangle^{-m}$, $m > 2$, and in [3], $W \in L_{n/2}(\mathbb{R}^n)$, $n \geq 3$. The result of [9, 3] reads

$$(0.18) \quad N_+(t; H - E; W) = t^{n/2} (2\pi)^{-n} |v_n| \int_{\mathbb{R}^n} W^{n/2}(x) dx + o(t^{n/2}), \text{ as } t \rightarrow +\infty,$$

where $|v_n|$ is the volume of the unit ball in \mathbb{R}^n .

If W satisfies (0.1), (0.2) and there exist $C, c > 0$ and a cone K such that

$$(0.19) \quad W(x) \geq c\langle x \rangle^{-m}, \quad \forall x \in K, |x| \geq C,$$

this integral converges if and only if $m > 2$. We calculate the principal term of the asymptotics, with a remainder estimate, in cases when $m \leq 2$.

Theorem 0.6. *Let W obey (0.1) and (0.2) with $m = 2$, and let W admit a representation*

$$(0.20) \quad W(x) = W_{-2}(x) + W'(x),$$

where W_{-2} is positively homogeneous of degree -2 , positive on some set, and

$$(0.21) \quad |W'(x)| \leq C|x|^{-2-\epsilon},$$

for some $\epsilon > 0$.

Then

$$(0.22) \quad N_+(t; H - E; W) = c(W_{-2}) t^{n/2} (\ln t + O(1)), \quad \text{as } t \rightarrow +\infty,$$

where

$$(0.23) \quad c(W_{-2}) = (2\pi)^{-n} |v_n| 2^{-1} \int_{S_{n-1}} W_{-2}(x)^{n/2} dx.$$

Theorem 0.7. *Let W obey (0.1) and (0.2) with $m \in (0, 2)$ and admit a representation*

$$(0.24) \quad W(x) = W_{-m}(x) + W'(x),$$

where W_{-m} is positively homogeneous of degree $-m$, positive on some set, and

$$(0.25) \quad |W'(x)| \leq C|x|^{-m-\epsilon},$$

for some $\epsilon > 0$.

Then there exists $\omega > 0$ such that

$$(0.26) \quad N_+(t; H - E; W) = c_+(H - E, W_{-m}) t^{n/m} (1 + O(t^{-\omega})), \text{ as } t \rightarrow +\infty,$$

where

$$(0.27) \quad \begin{aligned} c_+(H - E, W_{-m}) &= (2\pi)^{-n} n^{-1} \int_{S_{n-1}} W_{-m}(x)^{n/m} dx \int_{\mathbb{R}^{n*}/\Gamma^*} \sum_{j=p+1}^{+\infty} (E_j(\theta) - E)^{-n/m} d\theta, \end{aligned}$$

and p is the same as in (0.5).

Remark 0.1. The series in (0.27) converges if and only if $m < 2$, and the integral in (0.18) converges if and only if $m > 2$, so $m = 2$ is an “intermediate case” between these two essentially different cases. Similar three cases arise in the theory of degenerate elliptic operators (see [5, 15, 19]).

Remark 0.2. Theorem 0.6 is valid for any $V \in L_\infty(\mathbb{R}^n)$.

Theorem 0.6 will be proved in Subsection 3.1, using a rather elementary method. The “strong degeneration case” $m < 2$, when the principal term of asymptotics is expressed via an operator-valued symbol, requires a heavier machinery. In Subsection 3.2, we shall use a modification of the proof of the general theorem of the approximate spectral projection method [16, Section 22], for pdo with operator-valued symbols. In fact, we shall prove the following result which is a bit more general than Theorem 0.7, and then derive Theorem 0.7 from it.

Theorem 0.8. *Let W obey (0.1), (0.2) and (0.19) with $m \in (0, 2)$, and let*

$$(0.28) \quad \begin{aligned} c_+(t; H - E; W) &= (2\pi)^{-n} \sum_{j=p+1}^{+\infty} \text{meas}\{(x, \theta) \in \mathbb{R}^n \times \mathbb{R}^{n*}/\Gamma^* \mid 1 - tW(x)(E_j(\theta) - E)^{-1} < 0\} \end{aligned}$$

where p is the same as in (0.5), satisfy a Tauberian condition (0.8).

Then there exists $\omega > 0$ such that

$$(0.29) \quad N_+(t; H - E; W) = c_+(t; H - E; W)(1 + O(t^{-\omega})), \text{ as } t \rightarrow +\infty.$$

0.9. The case of non-negative W .

Theorem 0.9. *Let $W \geq 0$ satisfy (0.19) and the condition*

$$(0.30) \quad |D^\alpha W(x)| \leq C_\alpha \langle x \rangle^{-m-|\alpha|\rho}, \quad \forall \alpha.$$

Then:

- a) if $m > 0$ and (0.8) holds, then (0.9) holds;*
- b) if $m = 2$, and (0.20) and (0.21) hold, then (0.22) holds but with $o(\ln t)$ instead of $O(1)$;*
- c) if $m \in (0, 2)$ and $c_+(t; H - E; W)$ satisfies (0.8), then (0.29) holds; in particular, if (0.24) and (0.25) hold then (0.26) holds.*

0.10. Singular perturbations. For $\gamma \in \Gamma$ and $W_1 : \mathbb{R}^n \rightarrow \mathbb{R}$, define a multiplier in $L_2(\Omega)$:

$$W_1(\cdot + \gamma) : u(x) \mapsto W_1(x + \gamma)u(x),$$

and set

$$(\delta_j W_1)(\cdot + \gamma) = W_1(\cdot + \gamma + \alpha^j) - W_1(\cdot + \gamma).$$

Theorem 0.10. *Let W satisfy (0.1), (0.2), and let $W_1 \geq 0$ satisfy*

$$(0.31) \quad \|\delta_{i_1} \cdots \delta_{i_l} W_1(\cdot + \gamma)\|_{\text{End } L_2(\Omega)} \leq C_p W(\gamma) \langle \gamma \rangle^{-\epsilon - \rho l},$$

where $\epsilon > 0$ is independent of l, i_1, \dots, i_l .

Then all the statements of Theorems 0.3 and 0.6–0.9 remain valid for $N_\pm(t; H - E; W + W_1)$, the right-hand sides of all the asymptotic formulae remaining expressed via W .

The sharp remainder estimate (0.11) holds if $\epsilon = 1$, and a two-termed asymptotic formula (0.17) holds if $\epsilon > 1$.

0.11. In Section 1, we study a pdo on a torus. We prove Theorems 0.2–0.5 in Section 2, Theorems 0.6–0.8 in Section 3, and Theorems 0.9 and 0.10 in Section 4.

In Section 5 we list necessary theorems of a calculus of pdo with operator-valued symbols, and in Section 6 we state two general theorems of the approximate spectral projection method for pdo in \mathbb{R}^n [16] and prove two analogues of one of them for pdo on a torus.

ACKNOWLEDGEMENTS

The author thanks Tony Sudbery of the University of York, and Jochen Brüning of Universität Augsburg, for their kind hospitality during the first and last stages of the work, respectively.

1. A STUDY OF A PDO ON A TORUS

1.1. Let $\rho \in (0, 1]$, and let $W : \mathbb{R}^n \rightarrow \mathbb{C}$ and $f \in C^1(\mathbb{R}^n)$ satisfy the following conditions:

- (a) For any $l \in \mathbb{Z}_+$,

(1.1)

$$c_l(W; f; \rho) := \max_{\gamma \in \Gamma} \|\delta_{j_1} \cdots \delta_{j_l} W(\cdot + \gamma)\|_{\text{End } L_2(\Omega)} f(\gamma)^{-1} \langle \gamma \rangle^{\rho l} < \infty,$$

with max taken over all l -tuples of integers belonging to $\{1, \dots, n\}$.

- (b) There is C such that

$$(1.2) \quad |(\nabla f)(x)| \leq C f(x) \langle x \rangle^{-\rho}.$$

(c) There are $d_1 > 0, d, m_1, m$ such that

$$(1.3) \quad d_1 \langle x \rangle^{-m_1} \leq f(x) \leq d \langle x \rangle^{-m}.$$

We write $W \in \mathcal{F}_\rho(f)$. $\mathcal{F}_\rho(f)$ is a Frechet space with $c_j(\cdot; f; \rho)$ as seminorms.

Let \mathcal{U} be defined by (0.3). We set $\mathcal{W} := \mathcal{U}W\mathcal{U}^*$. It acts as follows:

$$(\mathcal{W}u)(x, \theta) = \frac{1}{\text{vol } \Omega^*} \sum_{\gamma \in \Gamma} W(x + \gamma) \int_{\Omega^*} \exp\{i(\theta' - \theta, \gamma)\} u(x, \theta') d\theta'.$$

We shall regard \mathcal{W} as an operator in $L_2(\mathbb{R}^{n^*}/\Gamma^*; L_2(\Omega))$. The aim of this section is to show that \mathcal{W} is a pdo on $\mathbb{R}^{n^*}/\Gamma^*$, a compact manifold without boundary, with the symbol taking values in $\text{End } L_2(\Omega)$, the space of bounded linear operators in $L_2(\Omega)$. We shall use local charts

$$\kappa_j : (\mathbb{R}^n \supset) U_j \rightarrow U_j^1 (\subset \mathbb{R}^{n^*}/\Gamma^*)$$

such that $\kappa_s^{-1} \circ \kappa_r$ is the restriction of a linear operator in \mathbb{R}^n ; for κ a local chart,

$$\kappa^* : C_0^\infty(U^1) \rightarrow C_0^\infty(U), \quad (\kappa^{-1})^* : C_0^\infty(U) \rightarrow C_0^\infty(U^1)$$

stand for the corresponding maps.

We shall also need the following simple corollary of (1.2)-(1.3).

Lemma 1.1. *There are $C = C(f), N = N(f)$ such that for all x, y ,*

$$(1.4) \quad f(y) \leq C f(x) (1 + |x - y|)^N.$$

Proof. Clearly, there is $C_1 > 0$ such that

$$\langle x \rangle^\rho \langle y \rangle^{-\rho} \leq C_1, \quad \text{provided } |x - y| < \langle x \rangle^\rho / 2.$$

Fix $c < \min\{1/2, (CC_1)^{-1}\}$. Taking the supremum over a set of all (x, y) with $|x - y| < c \langle x \rangle^\rho$ and applying the Lagrange formula and (1.2), we obtain

$$\begin{aligned} \sup f(x)^{-1} f(y) &\leq 1 + \sup f(x)^{-1} |f(y) - f(x)| \\ &\leq 1 + c \langle x \rangle^\rho f(x)^{-1} \sup |\nabla f(y)| \\ &\leq 1 + cC \langle x \rangle^\rho f(x)^{-1} \sup f(y) \langle y \rangle^{-\rho} \\ &\leq 1 + cCC_1 \sup f(x)^{-1} f(y). \end{aligned}$$

Hence,

$$f(y) \leq (1 - cCC_1)^{-1} f(x), \quad \text{if } |x - y| \leq c \langle x \rangle^\rho.$$

Clearly, f^{-1} also satisfies (1.2). Therefore, applying the previous argument to f^{-1} , we find $c > 0$ and C_2 such that

$$f(y) \leq C_2 f(x), \quad \text{if } |x - y| \leq c \langle y \rangle^\rho.$$

Thus, (1.4) holds if $|x - y| \leq c \max\{\langle x \rangle^\rho, \langle y \rangle^\rho\}$, with some $c > 0$, but if $|x - y| \geq c \max\{\langle x \rangle^\rho, \langle y \rangle^\rho\}$, then we deduce (1.4) from (1.3):

$$f(y) f(x)^{-1} \leq C \langle x \rangle^{m-} \langle y \rangle^{-m} \leq CC_2 (1 + |x - y|)^{N_2},$$

where $N_2 = 2 \max\{0, -m, m_-\}/\rho$, $C_2 = \max\{1, 2c^{-2/\rho}\}$. \square

The next lemma describes an especially nice subclass of $\mathcal{F}_\rho(f)$.

Lemma 1.2. *For all j , let*

$$(1.5) \quad c_j^0(W; f; \rho) := \sup_{|\alpha| \leq j} \sup_{\mathbb{R}^n} |D^\alpha W(x)| f(x)^{-1} \langle x \rangle^{\rho|\alpha|} < \infty.$$

Then (1.1) holds.

Proof. Without loss of generality we may assume that $\{\alpha^j\}$ is the canonical basis of \mathbb{R}^n . Then

$$(\delta_j W)(x + \gamma) = \int_0^1 (\partial_j W)(x + \gamma + t\alpha^j) dt,$$

$$(\delta_{j_1} \cdots \delta_{j_p} W)(x + \gamma) = \int_0^1 \cdots \int_0^1 (\partial_{j_1} \cdots \partial_{j_p} W)(x + \gamma + t_1 \alpha^{j_1} + \cdots + t_p \alpha^{j_p}) dt_1 \cdots dt_p,$$

and due to (1.5)

$$\begin{aligned} & \|(\delta_{j_1} \cdots \delta_{j_p} W)(\cdot + \gamma)\|_{\text{End } L_2} \\ & \leq C_p \sup_{|x - \gamma| \leq p+1} f(x) \langle x \rangle^{-p\rho} \leq C_{1,p} f(\gamma) \langle \gamma \rangle^{-p\rho} \end{aligned}$$

(the last inequality follows from the similar inequality obtained in the proof of Lemma 1.1). \square

If W satisfies (1.5) we write $W \in \mathcal{F}_\rho^0(f)$; this is a Fréchet space with $c_j^0(\cdot; f; \rho)$ as seminorms.

1.2. Some definitions of the theory of pdo on a torus. Let B be a Banach space. $L^{-\infty}(\mathbb{R}^{n^*}/\Gamma^*; B)$ stands for a class of integral operators whose kernels are of the class $C^\infty(\mathbb{R}^{n^*}/\Gamma^* \times \mathbb{R}^{n^*}/\Gamma^*; B)$.

Let f satisfy (1.2)-(1.3), and set $\Lambda(x) = \langle x \rangle$. We say that $a \in S_{\rho,0}(f; \mathbb{R}^n; B)$ if and only if

$$n_m(a; f; \rho) := \sup_{|\alpha|+|\beta| \leq m} \sup_{\mathbb{R}^{2n}} \|D_\xi^\alpha D_x^\beta a(x, \xi)\|_B f(-\xi)^{-1} \langle \xi \rangle^{\rho|\alpha|} < \infty, \quad m = 0, 1, \dots$$

It is a Fréchet space with $n_m(\cdot; f; \rho)$ as seminorms. We set

$$S^{-\infty}(\mathbb{R}^n; B) := \bigcap_m S_{\rho,0}(\Lambda^m; \mathbb{R}^n; B).$$

If A is a pdo with symbol of the class $S_{\rho,0}(f; \mathbb{R}^n; B)$, we write $A \in L_{\rho,0}(f; \mathbb{R}^n; B)$.

We say that $A \in L_{\rho,0}(f; \mathbb{R}^{n^*}/\Gamma^*; B)$ if and only if the following two conditions hold

a) for any $\phi, \psi \in C^\infty(\mathbb{R}^{n^*}/\Gamma^*)$ with $\text{supp } \phi \cap \text{supp } \psi = \emptyset$,

$$\phi A \psi \in L^{-\infty}(\mathbb{R}^{n^*}/\Gamma^*; B);$$

b) for a local chart $\kappa: U \rightarrow U^1$ and any $\phi, \psi \in C_0^\infty(U^1)$,

$$\phi A \psi = (\kappa^{-1})^*(\phi \circ \kappa) A_\kappa (\psi \circ \kappa) \kappa^*,$$

where $A_\kappa \in L_{\rho,0}(f; \mathbb{R}^n; B)$.

The symbol a_κ of A_κ is called a local symbol of A . Under the conditions imposed on the local charts, local symbols give rise to the global symbol a , which is a well-defined function on $T^*(\mathbb{R}^{n^*}/\Gamma^*)$ modulo symbols of the class $S^{-\infty}$. We write $a \in S_{\rho,0}(f; \mathbb{R}^{n^*}/\Gamma^*; B)$.

If $B = \mathbb{C}$, one omits B in all the notation.

For further information about pdo of these classes, see Section 5.

1.3. Calculation of the symbol of \mathcal{W} . We want to show that \mathcal{W} belongs to $L_{\rho,0}(f; \mathbb{R}^{n^*}/\Gamma^*; \text{End } L_2(\Omega))$ and calculate its symbol. In doing so, we may use $\phi, \psi \in C^\infty(\mathbb{R}^{n^*}/\Gamma^*)$ with sufficiently small $\text{diam supp } \phi, \text{diam supp } \psi$.

Lemma 1.3. *Let $\phi, \psi \in C^\infty(\mathbb{R}^{n^*}/\Gamma^*)$ satisfy the following condition: there are $j \in \{1, \dots, n\}$ and $c > 0$ such that*

$$(1.6) \quad |\exp\{-i(\theta' - \theta, \alpha^j)\} - 1| > c, \quad \forall \theta \in \text{supp } \phi, \theta' \in \text{supp } \psi.$$

Then $\phi \mathcal{W} \psi \in L^{-\infty}(\mathbb{R}^{n^}/\Gamma^*; \text{End } L_2(\Omega))$.*

Proof. Take $\theta \in \text{supp } \phi$ and j from (1.6). We have

$$\begin{aligned} & \sum_{\gamma \in \Gamma} \int_{\mathbb{R}^{n^*}/\Gamma^*} W(\cdot + \gamma) \exp\{i(\theta' - \theta, \gamma)\} \psi(\theta') u(\cdot, \theta') d\theta' \\ &= \sum_{\gamma \in \Gamma} \int_{\mathbb{R}^{n^*}/\Gamma^*} W(\cdot + \gamma + \alpha^j) \exp\{i(\theta' - \theta, \gamma + \alpha^j)\} \psi(\theta') u(\cdot, \theta') d\theta' \\ &= \sum_{\gamma \in \Gamma} \int_{\mathbb{R}^{n^*}/\Gamma^*} W(\cdot + \gamma) \exp\{i(\theta' - \theta, \gamma)\} \exp\{i(\theta' - \theta, \alpha^j)\} \psi(\theta') u(\cdot, \theta') d\theta' \\ &+ \sum_{\gamma \in \Gamma} \int_{\mathbb{R}^{n^*}/\Gamma^*} (\delta_j W)(\cdot + \gamma) \exp\{i(\theta' - \theta, \gamma)\} \exp\{i(\theta' - \theta, \alpha^j)\} \psi(\theta') u(\cdot, \theta') d\theta'. \end{aligned}$$

By substituting $W(\cdot + \gamma)(1 - \exp\{i(\theta' - \theta, \alpha^j)\})^{-1}$ for $W(\cdot + \gamma)$, we obtain

$$\begin{aligned} & \sum_{\gamma \in \Gamma} \int_{\mathbb{R}^{n^*}/\Gamma^*} W(\cdot + \gamma) \exp\{i(\theta' - \theta, \gamma)\} \psi(\theta') u(\cdot, \theta') d\theta' \\ &= \sum_{\gamma \in \Gamma} \int_{\mathbb{R}^{n^*}/\Gamma^*} (\exp\{-i(\theta' - \theta, \alpha^j)\} - 1)^{-1} (\delta_j W)(\cdot + \gamma) \\ &\quad \times \exp\{i(\theta' - \theta, \gamma)\} \psi(\theta') u(\cdot, \theta') d\theta'. \end{aligned}$$

Iterating this procedure of “summation by parts” we get, for any N ,

$$\begin{aligned} & \sum_{\gamma \in \Gamma} \int_{\mathbb{R}^{n^*}/\Gamma^*} W(\cdot + \gamma) \exp\{i(\theta' - \theta, \gamma)\} \psi(\theta') u(\cdot, \theta') d\theta' \\ &= \sum_{\gamma \in \Gamma} \int_{\mathbb{R}^{n^*}/\Gamma^*} (\exp\{-i(\theta' - \theta, \alpha^j)\} - 1)^{-N} (\delta_j^N W)(\cdot + \gamma) \\ &\quad \times \exp\{i(\theta' - \theta, \gamma)\} \psi(\theta') u(\cdot, \theta') d\theta'. \end{aligned}$$

On the support of the integrand, the first factor is bounded due to (1.6), and the second one admits an estimate

$$\|(\delta_j^N W)(\cdot + \gamma)\| \leq C_N \langle \gamma \rangle^{-m-\rho N}, \quad \forall \gamma \in \Gamma,$$

due to (1.1). Take $N > (n-m)/\rho$ and $u \in C^\infty(\mathbb{R}^{n^*}/\Gamma^*; L_2(\Omega))$; then we can do the summation first to see that $\phi \mathcal{W} \psi$ is an integral operator with the kernel

$$\begin{aligned} (1.7) \quad & K(W, \phi, \psi; \theta, \theta') \\ &= \frac{1}{\text{vol } \Omega^*} \sum_{\gamma \in \Gamma} (\exp\{-i(\theta' - \theta, \alpha^j)\} - 1)^{-N} \phi(\theta) \psi(\theta') (\delta_j^N W)(\cdot + \gamma) \exp\{i(\theta' - \theta, \gamma)\}. \end{aligned}$$

Hence

$$\|K(W, \phi, \psi; \theta, \theta')\|_{\text{End } L_2(\Omega)} \leq C(W, \phi, \psi) \sum_{\gamma \in \Gamma} \langle \gamma \rangle^{-\rho N - m} = C_1(W, \phi, \psi)$$

uniformly in θ, θ' .

To obtain the similar estimate for a derivative $D_\theta^\alpha D_{\theta'}^\beta K(W, \phi, \psi; \theta, \theta')$, we take $N > (n - m + |\alpha| + |\beta|)/\rho$ and differentiate (1.7). \square

Lemma 1.4. *Let $\kappa : U \rightarrow U^1$ be a local chart with small $\text{diam } U$, and let $\phi, \psi \in C_0^\infty(U^1)$.*

Then

$$(1.8) \quad \phi \mathcal{W} \psi = (\kappa^{-1})^*(\phi \circ \kappa)(B_{\kappa,0} + B_{\kappa,\infty})(\psi \circ \kappa) \kappa^*,$$

where $B_{\kappa,0} \in L_{\rho,0}(f; \mathbb{R}^n; \text{End } L_2(\Omega))$ has the symbol $b_{\kappa,0} = b_{\kappa,0}(\xi)$, which is independent of θ , and $B_{\kappa,\infty} \in L^{-\infty}(\mathbb{R}^n; \text{End } L_2(\Omega))$.

Proof. Fix $\mu \in C_0^\infty(\mathbb{R})$, $0 \leq \mu \leq 1$, $\mu(t) = 1$ ($|t| < 1$), $\mu(t) = 0$ ($|t| > 2$), and for $M > 1$, set

$$W_M(x) = \mu(|x|/M)W(x), \quad W_{-M} = W - W_M, \quad \mathcal{W}_{\pm M} = \mathcal{U}W_{\pm M}\mathcal{U}^*.$$

Then $W_{\pm M}$ satisfies (1.1) uniformly in $M > 1$. Take $u \in C^\infty(\mathbb{R}^{n^*}/\Gamma^*; L_2(\Omega))$. Integration by parts gives

$$\begin{aligned} & (\mathcal{W}_{-M}\psi u)(\cdot, \theta) \\ &= \frac{1}{\text{vol } \Omega^*} \sum_{\gamma \in \Gamma} W_{-M}(\cdot + \gamma) \int_{\mathbb{R}^{n^*}/\Gamma^*} \exp\{i(\theta' - \theta, \gamma)\} (1 + |\gamma|^2)^{-N} \\ & \quad \times (1 - \Delta_{\theta'})^N \psi(\theta') u(x, \theta') d\theta'. \end{aligned}$$

Take $N > n - m$; since $|x| \geq M$ for $x \in \text{supp } W_{-M}$, we have

$$\sup_{\theta \in \mathbb{R}^{n^*}/\Gamma^*} \|(\mathcal{W}_{-M}\psi u)(\cdot, \theta)\|_{L_2(\Omega)} \leq C(W, N) \sum_{|\gamma| \geq M} \langle \gamma \rangle^{-m-N} \rightarrow 0,$$

as $M \rightarrow \infty$. To obtain similar estimates for $D_\theta^\alpha (\mathcal{W}_{-M}\psi u)$, we take $N > |\alpha| + n - m$. Thus, $\mathcal{W}_{-M}u \rightarrow 0$ in $C^\infty(\mathbb{R}^{n^*}/\Gamma^*; L_2(\Omega))$ as $M \rightarrow \infty$.

It follows that (1.8) will be proved together with the formula

$$(1.9) \quad \phi \mathcal{W}_M \psi = (\kappa^{-1})^*(\phi \circ \kappa)(B_{\kappa,M,0} + B_{\kappa,M,\infty})(\psi \circ \kappa) \kappa^*,$$

where

(a) $B_{\kappa,M,0} \in L_{\rho,0}(f; \mathbb{R}^n; \text{End } L_2(\Omega))$ and $B_{\kappa,M,\infty} \in L^{-\infty}(\mathbb{R}^n; \text{End } L_2(\Omega))$ uniformly in M ,

(b) $B_{\kappa,M,0}$ has the symbol $b_{\kappa,M,0} = b_{\kappa,M,0}(\xi)$ which is independent of θ , and

(c) The limits

$$(1.10) \quad b_{\kappa,0}(\xi) = \lim_{M \rightarrow \infty} b_{\kappa,M,0}(\xi),$$

$$(1.11) \quad b_{\kappa,\infty}(\theta, \xi) = \lim_{M \rightarrow \infty} b_{\kappa,\infty,0}(\theta, \xi).$$

exists uniformly on any compact set in \mathbb{R}^n and $U \times \mathbb{R}^n$ respectively.

Take $\chi_1, \chi_2 \in C_0^\infty(U)$ such that $(\phi \circ \kappa)\chi_1 = \phi \circ \kappa$, $(\psi \circ \kappa)\chi_2 = \psi \circ \kappa$, $\chi_2(\theta) = 1$ for $\theta \in \text{supp } \chi_1$, and put

$$\begin{aligned} & (\tilde{W}_M u)(\cdot, \theta') \\ &= \frac{1}{\text{vol } \Omega^*} \sum_{\gamma \in \Gamma} W_M(\cdot + \gamma) \int_{\mathbb{R}^n} \chi_1(\theta) \chi_2(\theta') \exp\{i(\theta' - \theta, \gamma)\} u(x, \theta') d\theta'. \end{aligned}$$

This is an integral operator with the kernel

$$K(W_M, \chi_1, \chi_2; \theta, \theta') = \frac{1}{\text{vol } \Omega^*} \sum_{\gamma \in \Gamma} W_M(\cdot + \gamma) \chi_1(\theta) \chi_2(\theta') \exp\{i(\theta' - \theta, \gamma)\}.$$

Therefore, \tilde{W}_M is a pdo with the symbol

$$\begin{aligned} (1.12) \quad \tilde{b}_M(\theta, \xi) &= \int_{\mathbb{R}^n} \chi_1(\theta) \chi_2(\theta + \theta') K(W_M, \chi_1, \chi_2; \theta, \theta + \theta') \exp\{i(\theta', \xi)\} d\theta' \\ &= \frac{1}{\text{vol } \Omega^*} \int_{\mathbb{R}^n} \chi_1(\theta) \chi_2(\theta + \theta') \sum_{\gamma \in \Gamma} W_M(\cdot + \gamma) \exp\{i(\theta', \gamma + \xi)\} d\theta'. \end{aligned}$$

Take $\chi \in C_0^\infty(\mathbb{R}^n)$ such that $\chi(\theta') = 1$ on the support of the integrand in (1.12), notice that

$$\chi_1(\theta) \chi_2(\theta + \theta') = \chi_1(\theta) + r(\theta, \theta'),$$

where

$$(1.13) \quad |D_\theta^\nu D_{\theta'}^\omega r(\theta, \theta')| \leq C(\nu, \omega, N) |\theta'|^N,$$

for all N and all multi-indices ν, ω , and set

$$b_{M,0}(\xi) = \frac{1}{\text{vol } \Omega^*} \int_{\mathbb{R}^n} \chi(\theta') \sum_{\gamma \in \Gamma} W_M(\cdot + \gamma) \exp\{i(\theta', \gamma + \xi)\} d\theta',$$

$$\tilde{b}_{M,0}(\theta, \xi) = \chi_1(\theta) b_{M,0}(\xi),$$

$$\tilde{b}_{M,\infty}(\theta, \xi) = \frac{1}{\text{vol } \Omega^*} \int_{\mathbb{R}^n} r(\theta, \theta') \chi(\theta') \sum_{\gamma \in \Gamma} W_M(\cdot + \gamma) \exp\{i(\theta', \gamma + \xi)\} d\theta'.$$

Clearly,

$$\tilde{b}_M(\theta, \xi) = \tilde{b}_{M,0}(\theta, \xi) + \tilde{b}_{M,\infty}(\theta, \xi).$$

Integration by parts gives

$$\begin{aligned} & (\xi^\sigma D_\theta^\alpha D_\xi^\beta \tilde{b}_{M,\infty})(\theta, \xi) \\ &= \frac{1}{\text{vol } \Omega^*} \int_{\mathbb{R}^n} D_\theta^\alpha r(\theta, \theta') \chi(\theta') \sum_{\gamma \in \Gamma} W_M(\cdot + \gamma) \theta'^\beta \exp\{i(\theta', \gamma)\} D_{\theta'}^\sigma \exp\{i(\theta', \xi)\} d\theta' \\ &= \frac{1}{\text{vol } \Omega^*} \int_{\mathbb{R}^n} (-1)^{|\sigma|} D_{\theta'}^\sigma (D_\theta^\alpha r(\theta, \theta') \chi(\theta') \theta'^\beta) \\ & \quad \times \exp\{i(\theta', \gamma)\} \sum_{\gamma \in \Gamma} W_M(\cdot + \gamma) \exp\{i(\theta', \xi)\} d\theta'. \end{aligned}$$

The estimate (1.13) allows us to sum by parts as in Lemma 1.3 and obtain

$$\| \xi^\sigma D_\theta^\alpha D_\xi^\beta \tilde{b}_{M,\infty}(\theta, \xi) \|_{\text{End } L_2(\Omega)} \leq C(\sigma, \alpha, \beta) \sum_{\gamma \in \Gamma} \langle \gamma \rangle^{-n-1} = C_1(\sigma, \alpha, \beta),$$

uniformly in θ, ξ ; similarly, for $M_1 > M > 1$,

$$\| \xi^\sigma D_\theta^\alpha D_\xi^\beta (\tilde{b}_{M,\infty}(\theta, \xi) - \tilde{b}_{M_1,\infty}(\theta, \xi)) \|_{\text{End } L_2(\Omega)} \leq C(\sigma, \alpha, \beta) \sum_{|\gamma| \geq M} \langle \gamma \rangle^{-n-1} \rightarrow 0$$

as $M \rightarrow \infty$, uniformly in θ, ξ from a given compact.

Hence, $\tilde{b}_{M,\infty} \in S^{-\infty}(\mathbb{R}^n; \text{End } L_2(\Omega))$ uniformly in $M > 1$, and the limit (1.11) exists with $b_\infty \in S^{-\infty}(\mathbb{R}^n; \text{End } L_2(\Omega))$.

Now we consider

$$D_\xi^\beta b_{M,0}(\xi) = \frac{1}{\text{vol } \Omega^*} \int_{\mathbb{R}^n} \chi(\theta') \sum_{\gamma \in \Gamma} W_M(\cdot + \gamma) \theta'^\beta \exp\{i(\theta', \gamma + \xi)\} d\theta'.$$

Set

$$\delta^\beta = \delta_1^{\beta_1} \cdots \delta_n^{\beta_n}, \quad f_\beta(\theta') = \prod_{1 \leq j \leq n} ((\exp\{-i(\theta'_j, a^j)\} - 1)^{-1} \theta'_j)^{\beta_j} \chi(\theta'),$$

sum by parts and integrate by parts:

$$\begin{aligned} D_\xi^\beta b_{M,0}(\xi) &= \frac{1}{\text{vol } \Omega^*} \int_{\mathbb{R}^n} ((1 - \Delta_{\theta'})^{N_1} f_\beta(\theta')) \\ &\quad \times \sum_{\gamma \in \Gamma} (\delta^\beta W_M)(\cdot + \gamma) \langle \gamma + \xi \rangle^{-2N_1} \exp\{i(\theta', \gamma + \xi)\} d\theta'. \end{aligned}$$

Let N be as in (1.4). Then due to (1.1) and (1.4),

$$\begin{aligned} \| (\delta^\beta W_M)(\cdot + \gamma) \langle \gamma + \xi \rangle^{-2N_1} \| &\leq C_{\beta, N_1} f(\gamma) \langle \gamma \rangle^{-\rho|\beta|} \langle \gamma + \xi \rangle^{-2N_1} \\ &\leq C_{1, \beta, N_1} f(-\xi) \langle \gamma + \xi \rangle^{N-2N_1} \langle \gamma \rangle^{-\rho|\beta|} \leq C_{2, \beta, N_1} f(-\xi) \langle \xi \rangle^{-\rho|\beta|} \langle \gamma + \xi \rangle^{N+\rho|\beta|-2N_1}. \end{aligned}$$

Taking $N_1 > (n + \rho|\beta| + N)/2$, we obtain

$$\| D_\xi^\beta b_{M,0}(\xi) \|_{\text{End } L_2(\Omega)} \leq C_1(\beta, N) f(-\xi) \langle \xi \rangle^{-\rho|\beta|}.$$

Similarly, for $M_1 > M > 1$,

$$\| D_\xi^\beta (b_{M,0}(\xi) - b_{M_1,0}(\xi)) \|_{\text{End } L_2(\Omega)} \leq C(\beta, N) \sum_{|\gamma| > M} f(\gamma) \langle \gamma \rangle^{-\rho|\beta|} \langle \gamma + \xi \rangle^{-2N_1} \rightarrow 0$$

as $M \rightarrow \infty$, uniformly in ξ from a given compact.

Hence, $b_{M,0} \in S_{\rho,0}(f; \mathbb{R}^n; \text{End } L_2(\Omega))$ uniformly in $M > 1$, and the limit (1.10) exists with $b_0 \in S_{\rho,0}(f; \mathbb{R}^n; \text{End } L_2(\Omega))$.

This completes the proof of Lemma 1.3. \square

Lemma 1.5. *Put*

(1.14)

$$w(\xi) = \lim_{M \rightarrow \infty} \lim_{t \rightarrow \infty} \frac{1}{\text{vol } \Omega^*} \int_{\mathbb{R}^n} \sum_{\gamma \in \Gamma} \mu(|\theta'|/t) W_M(\cdot + \gamma) \exp\{i(\theta', \gamma + \xi)\} d\theta'.$$

Then $b_0 - w \in S^{-\infty}(\mathbb{R}^n; \text{End } L_2(\Omega))$.

Proof. The arguments from Lemma 1.4 show that

(1.15)

$$b_0(\xi) = \lim_{M \rightarrow \infty} \frac{1}{\text{vol } \Omega^*} \int_{\mathbb{R}^n} \chi(\theta') \sum_{\gamma \in \Gamma} W_M(\cdot + \gamma) \exp\{i(\theta', \gamma + \xi)\} d\theta'.$$

Hence

$$w(\xi) - b_0(\xi) = \lim_{M \rightarrow \infty} \lim_{t \rightarrow \infty} \frac{1}{\text{vol } \Omega^*} \int_{\mathbb{R}^n} (\mu(|\theta'|/t) - \chi(\theta')) \sum_{\gamma \in \Gamma} W_M(\cdot + \gamma) \exp\{i(\theta', \gamma + \xi)\} d\theta'.$$

By integrating by parts, we obtain

$$\begin{aligned} w(\xi) - b_0(\xi) &= \lim_{M \rightarrow \infty} \lim_{t \rightarrow \infty} \frac{1}{\text{vol } \Omega^*} \int_{\mathbb{R}^n} \sum_{\gamma \in \Gamma} (1 + |\gamma + \xi|^2)^{-N} W_M(\cdot + \gamma) \exp\{i(\theta', \gamma + \xi)\} \\ &\quad \times (1 - \Delta_{\theta'})^N (\mu(|\theta'|/t) - \chi(\theta')) d\theta' \\ &= \lim_{M \rightarrow \infty} \lim_{t \rightarrow \infty} \frac{1}{\text{vol } \Omega^*} \int_{\mathbb{R}^n} \sum_{\gamma \in \Gamma} (1 + |\gamma + \xi|^2)^{-N} W_M(\cdot + \gamma) \exp\{i(\theta', \gamma + \xi)\} \\ &\quad \times (\mu(|\theta'|/t) - \chi(\theta')) d\theta' \\ &\quad + \lim_{M \rightarrow \infty} \lim_{t \rightarrow \infty} \frac{1}{\text{vol } \Omega^*} \int_{\mathbb{R}^n} \sum_{\gamma \in \Gamma} (1 + |\gamma + \xi|^2)^{-N} W_M(\cdot + \gamma) \exp\{i(\theta', \gamma + \xi)\} \\ &\quad \times \{(1 - \Delta_{\theta'})^N - 1\} (-\chi(\theta')) d\theta' \\ &\quad + \lim_{M \rightarrow \infty} \lim_{t \rightarrow \infty} \frac{1}{\text{vol } \Omega^*} \int_{\mathbb{R}^n} \sum_{\gamma \in \Gamma} (1 + |\gamma + \xi|^2)^{-N} W_M(\cdot + \gamma) \exp\{i(\theta', \gamma + \xi)\} \\ &\quad \times \{(1 - \Delta_{\theta'})^N - 1\} (\mu(|\theta'|/t) - \chi(\theta')) d\theta'. \end{aligned}$$

Fix ξ and $N > (n - m)/(2\rho)$; then all the integrals converge absolutely and are uniformly bounded. Passing to the limit under the integral sign, we obtain

$$\begin{aligned} w(\xi) - b_0(\xi) &= \lim_{M \rightarrow \infty} \frac{1}{\text{vol } \Omega^*} \int_{\mathbb{R}^n} \sum_{\gamma \in \Gamma} (1 + |\gamma + \xi|^2)^{-N} W_M(\cdot + \gamma) \exp\{i(\theta', \gamma + \xi)\} \times \\ &\quad \times (1 - \Delta_{\theta'})^N (1 - \chi(\theta')) d\theta'. \end{aligned}$$

Since $1 - \chi(\theta') = 0$ in a neighbourhood of 0, we can sum by parts and prove that $w - b_0 \in S^{-\infty}(\mathbb{R}^n; \text{End } L_2(\Omega))$. \square

By combining Lemmas 1.3-1.5, we obtain the following theorem.

Theorem 1.6. $\mathcal{W} \in L_{\rho,0}(f; \mathbb{R}^{n^*}/\Gamma^*; \text{End } L_2(\Omega))$, and its symbol is given by (1.14).

1.4. **A formula for w in the case $W \in \mathcal{F}_\rho^0(f)$.** Define the formal power series $L_j \in \mathbb{C}[[s, t]] = \mathbb{C}[[s_1, \dots, s_n, t_1, \dots, t_n]]$ by

$$L_j(s, t) = t_j(\exp\{is_j\} - 1)/(\exp\{it_j\} - 1),$$

and set

$$\begin{aligned} L &= (L_1, \dots, L_n), \quad L^1(s, t) = s - L(s, t), \\ L^k(s, t) &= L^{k-1}(s, L(s, t)), \quad k = 2, 3, \dots \end{aligned}$$

Since $L_j(s, t) \in \sum_{l=1}^n s_l \mathbb{C}[[s, t]]$, there exists the limit

$$L^\infty(s) = \lim_{k \rightarrow \infty} L^k(s, t).$$

Set

$$\hat{L}(D) = \frac{1}{\text{vol } \Omega} \int_{\Omega} (1 - \exp\{i(x, L^\infty(-D))\}) dx \in \mathbb{C}[[D_1, \dots, D_n]].$$

Since $L_j(s, t) = s_j +$ terms of order greater than 1, we have $L^1(s, t) = 0 \bmod s$, and $\hat{L}(D) = I +$ terms of order greater than 1.

Theorem 1.7. *Let $W \in \mathcal{F}_\rho^0(f)$. Then*

$$(1.16) \quad w(\xi) \sim \sum_{|\alpha|+s \geq 0} \frac{(\cdot)^\alpha}{\alpha!} (\hat{L}(D)^s (iD)^\alpha W)(-\xi),$$

where

$$(\cdot)^\alpha : L_2(\Omega) \ni u(x) \mapsto x^\alpha u(x) \in L_2(\Omega).$$

The asymptotic sum in (1.16) is understood in the usual sense: let

$$\sum_{|\alpha|+s \geq 0} \frac{(\cdot)^\alpha}{\alpha!} \hat{L}(D)^s (iD)^\alpha = \sum_{|\beta| \geq 0} c_\beta(\cdot) D^\beta,$$

then (1.16) means that, for any N ,

$$(1.17) \quad w(\xi) = \sum_{|\beta| < N} c_\beta(\cdot) (D^\beta W)(-\xi) + r_N(\xi),$$

where $r_N \in S_{\rho,0}(f\Lambda^{-N\rho}; \mathbb{R}^{n*}/\Gamma^*; \text{End } L_2(\Omega))$. In particular,

$$(1.18) \quad w(\xi) = W(-\xi) + \sum_{j=1}^n (\cdot)_j (iD_j W)(-\xi) + r_2(\xi).$$

Proof. The argument used in the proof of Lemma 1.4 shows that it suffices to prove (1.17) for $W \in \mathbb{C}_0^\infty(\mathbb{R}^n)$, provided for any j

$$(1.19) \quad n_j(r_N; f\Lambda^{-\rho N}; \rho) \leq C c_l^0(W; f; \rho),$$

where C, l depend on j, ρ, f, N , and n , but not on W .

Due to Lemma 1.5 (see (1.15)), we have modulo $S^{-\infty}(\mathbb{R}^n; \text{End } L_2(\Omega))$:

$$\begin{aligned} w(\xi) &= \frac{1}{\text{vol } \Omega^*} \int_{\mathbb{R}^n} d\theta' \chi(\theta') \sum_{\gamma \in \Gamma} W(\cdot + \gamma) \exp\{i(\theta', \gamma + \xi)\} \\ (1.20) \quad &= \sum_{|\alpha| < N} \frac{(\cdot)^\alpha}{\alpha!} w_\alpha(\xi) + \sum_{|\alpha| = N} \frac{(\cdot)^\alpha}{\alpha!} r_\alpha(\xi), \end{aligned}$$

where

$$w_\alpha(\xi) = \frac{1}{\text{vol } \Omega^*} \int_{\mathbb{R}^n} d\theta' \chi(\theta') \sum_{\gamma \in \Gamma} ((iD)^\alpha W)(\gamma) \exp\{i(\theta', \gamma + \xi)\},$$

$$r_\alpha(\xi) = \frac{1}{\text{vol } \Omega^*} \int_{\mathbb{R}^n} d\theta' \chi(\theta') \int_0^1 dt \sum_{\gamma \in \Gamma} ((iD)^\alpha W)(t(\cdot) + \gamma) \exp\{i(\theta', \gamma + \xi)\}.$$

In (1.20) and below, “modulo $S^{-\infty}$ ” has a meaning similar to (1.19), namely, that $S^{-\infty}$ -seminorms of an “error term” admit estimates via seminorms of $W \in \mathcal{F}_\rho^0(f)$.

Due to (1.5), $((iD)^\alpha W)(t(\cdot) + \gamma) \in \mathcal{F}_\rho(f\Lambda^{-|\alpha|\rho})$ uniformly in $t \in [0, 1]$. Therefore Theorem 1.6 gives

$$(1.21) \quad r_\alpha \in S_{\rho,0}(f\Lambda^{-|\alpha|\rho}; \mathbb{R}^{n^*}/\Gamma^*; \text{End } L_2(\Omega))$$

with r_α satisfying (1.19).

For $V \in C_0^\infty(\mathbb{R}^n)$, set $V_1(x) = V(-x)$. We have

$$\begin{aligned} & \frac{1}{\text{vol } \Omega^*} \int_{\mathbb{R}^n} d\theta' \chi(\theta') \sum_{\gamma \in \Gamma} V(\gamma) \exp\{i(\theta', \xi + \gamma)\} \\ &= \frac{1}{\text{vol } \Omega^*} \int_{\mathbb{R}^n} d\theta' \chi(\theta') \sum_{\gamma \in \Gamma} V_1(\gamma) \exp\{i(\theta', \xi - \gamma)\} \\ (1.22) \quad &= v_1(\xi) + v_2(\xi), \end{aligned}$$

where

$$\begin{aligned} v_1(\xi) &= \frac{1}{\text{vol } \Omega^*} \int_{\mathbb{R}^n} d\theta' \chi(\theta') \sum_{\gamma \in \Gamma} \frac{1}{\text{vol } \Omega} \int_{\Omega} dx V_1(\gamma + x) \exp\{i(\theta', \xi - \gamma - x)\}, \\ v_2(\xi) &= \frac{1}{\text{vol } \Omega^*} \int_{\mathbb{R}^n} d\theta' \chi(\theta') \sum_{\gamma \in \Gamma} \frac{1}{\text{vol } \Omega} \int_{\Omega} dx \times \\ & \quad \times (V_1(\gamma) - V_1(\gamma + x) \exp\{-i(\theta', x)\}) \exp\{i(\theta', \xi - \gamma)\}. \end{aligned}$$

Clearly, modulo $S^{-\infty}(\mathbb{R}^n)$,

$$\begin{aligned} v_1(\xi) &= (2\pi)^{-n} \int_{\mathbb{R}^n} d\theta' \chi(\theta') \int_{\mathbb{R}^n} dx V_1(x) \exp\{i(\theta', \xi - x)\} \\ (1.23) \quad &= V_1(\xi) = V(-\xi), \end{aligned}$$

and

$$\begin{aligned} v_2(\xi) &\sim \frac{1}{\text{vol } \Omega^*} \int_{\mathbb{R}^n} d\theta' \chi(\theta') \sum_{\gamma \in \Gamma} \frac{1}{\text{vol } \Omega} \int_{\Omega} dx \\ (1.24) \quad &\times ((I - \exp\{i(x, D_x - \theta')\})V_1)(\gamma) \exp\{i(\theta', \xi - \gamma)\}, \end{aligned}$$

the RHS in (1.24) being understood in the sense explained in Section 5.

Write the summation by parts formula in the form

$$(1.25) \quad \sum_{\gamma \in \Gamma} V_1(\gamma) \theta^\omega \exp\{-i(\theta, \gamma)\} = \sum_{\gamma \in \Gamma} (L(D, \theta)^\omega V_1)(\gamma) \exp\{-i(\theta, \gamma)\},$$

where

$$L(D, \theta)^\omega = L_1(D, \theta)^{\omega_1} \cdots L_n(D, \theta)^{\omega_n},$$

$$L_j(D, \theta) = \theta_j(\exp\{iD_j\} - 1)/(\exp\{i\theta_j\} - 1).$$

Set

$$L^1(s, t) = s - L(s, t), \quad L^k(s, t) = L^{k-1}(s, L(s, t)), \quad k = 2, 3, \dots$$

By substituting (1.25) into (1.24), we obtain

$$(1.26) \quad v_2(\xi) \sim \frac{1}{\text{vol } \Omega^*} \int_{\mathbb{R}^n} d\theta' \chi(\xi) \sum_{\gamma \in \Gamma} \frac{1}{\text{vol } \Omega} \int_{\Omega} dx \\ \times ((I - \exp\{i(x, L^1(D, \theta))\})V_1)(\gamma) \exp\{i(\theta', \xi - \gamma)\}.$$

By iterating this procedure and using Theorem 5.6, we obtain (1.26) with L^k instead of L^1 , $k = 2, 3, \dots$. Since $L_l(s, t) \in \sum_{j=1}^n s_j \mathbb{C}[[s, t]]$, we have

$$(1.27) \quad v_2(\xi) \sim \frac{1}{\text{vol } \Omega^*} \int_{\mathbb{R}^n} d\theta' \chi(\xi) \sum_{\gamma \in \Gamma} (\tilde{L}(D)V_1)(\gamma) \exp\{i(\theta', \xi - \gamma)\},$$

where

$$\tilde{L}(D) = \frac{1}{\text{vol } \Omega} \int_{\Omega} (I - \exp\{i(x, L^\infty(D))\}) dx.$$

It follows from (1.22), (1.27) and (1.23), that

$$(1.28) \quad \frac{1}{\text{vol } \Omega^*} \int_{\mathbb{R}^n} d\theta' \chi(\theta') \sum_{\gamma \in \Gamma} ((I - \tilde{L}(D))V_1)(\gamma) \exp\{i(\theta', \xi - \gamma)\} \sim V_1(\xi).$$

Replacing V_1 in (1.28) by $\sum_{j \geq 0} \tilde{L}(D)^j V_1$ and using Theorem 5.5, we obtain

$$(1.29) \quad \frac{1}{\text{vol } \Omega^*} \int_{\mathbb{R}^n} d\theta' \chi(\theta') \sum_{\gamma \in \Gamma} V(\gamma) \exp\{i(\theta', \xi + \gamma)\} \\ \sim \sum_{j \geq 0} (\tilde{L}(D)^j V_1)(\xi) = \sum_{j \geq 0} (\hat{L}(D)^j V)(-\xi).$$

We apply (1.29) with $V = (iD)^\alpha W$ to (1.20); then (1.20) and (1.21) give (1.16). \square

2. ASYMPTOTICS OF $N_-(t; H - E; W)$

2.1. Auxiliary variational lemmas. We shall need the following definition and lemmas.

Let \mathfrak{A} be a quadratic form in a Hilbert space \mathfrak{H} , with a domain $D(\mathfrak{A})$, and let $V \subset D(\mathfrak{A})$ be a subspace. We assume that \mathfrak{A} is closable, and we set

$$\mathcal{N}(\lambda, \mathfrak{A}, V) = \sup\{\dim L \mid L \subset V, \mathfrak{A}[u] < \lambda \|u\|_{\mathfrak{H}}^2 \quad \forall (0 \neq) u \in L\}.$$

For A an operator in \mathfrak{H} and $\mathfrak{A}[u] = \langle Au, u \rangle_{\mathfrak{H}}$, $D(\mathfrak{A}) = D(A)$, write $\mathcal{N}(\lambda, A, V)$ instead of $\mathcal{N}(\lambda, \mathfrak{A}, V)$.

For A a self-adjoint operator and $(a, b) \subset \mathbb{R}$, set $N((a, b); A) = \dim \text{Ran } P_{(a, b)}(A)$, where $P_{(a, b)}(A)$ is a spectral projection of A . Set also $N(\lambda; A) = N((-\infty, \lambda); A)$.

The following lemmas are well-known and widely used; for proofs, see e.g. [6, 16, 18].

Lemma 2.1. *Let A be a semibounded from below self-adjoint operator with the domain $D(A)$.*

Then $N(\lambda; A) = \mathcal{N}(\lambda; A; D(A))$.

Lemma 2.2. $\mathcal{N}(\lambda; \mathfrak{A}; V)$ is independent of V provided V is a core of the form \mathfrak{A} .

Lemma 2.3. If $\mathfrak{A}[u] \leq \mathfrak{A}_1[u] \quad \forall u \in V$, then

$$\mathcal{N}(\lambda; \mathfrak{A}_1; V) \leq \mathcal{N}(\lambda; \mathfrak{A}; V).$$

Lemma 2.4. If $V \subset V_1$, then

$$\mathcal{N}(\lambda; \mathfrak{A}; V) \leq \mathcal{N}(\lambda; \mathfrak{A}; V_1).$$

Lemma 2.5. Let H_s be a Hilbert space, \mathfrak{A}_s a quadratic form in H_s , $V_s \subset D(\mathfrak{A}_s)$ a subspace ($s = 0, 1, \dots, r$), let

$$l: \bigoplus_{1 \leq s \leq r} V_s \rightarrow V_0$$

be an isomorphism, and let

$$\mathfrak{A}_0[l(u_1, \dots, u_r)] = \sum_{s=1}^r \mathfrak{A}_s[u_s].$$

Then

$$\mathcal{N}(\lambda; \mathfrak{A}_0; V_0) = \sum_{s=1}^r \mathcal{N}(\lambda; \mathfrak{A}_s; V_s).$$

2.2. Proof of Theorem 0.2. The Birman-Schwinger principle implies that $N_{\pm}(t; H - E; W)$ is equal to the number of positive eigenvalues counted, with multiplicity, of a problem

$$(2.1) \quad u = \pm \mu W^{1/2} (H - E)^{-1} W^{1/2} u, \quad u \in L_2(\mathbb{R}^n),$$

belonging to $(0, t)$ (see [1, 9]). Let \mathcal{U} be an isometry defined by (0.3), and set $\mathcal{H} = \mathcal{U}H\mathcal{U}^*$, $\mathcal{W} = \mathcal{U}W\mathcal{U}^*$. Then we can deal with the problem

$$(2.2) \quad u = \pm \mu \mathcal{W}^{1/2} (\mathcal{H} - E)^{-1} \mathcal{W}^{1/2} u$$

instead, and Lemmas 2.1, 2.2 and 2.5 give

$$(2.3) \quad N_{\pm}(t; H - E; W) = \mathcal{N}(0; \mathcal{W}^{-1} \mp t(\mathcal{H} - E)^{-1}; C^{\infty}(\mathbb{R}^{n*}/\Gamma^*; L_2(\Omega))).$$

Denote by $\mathcal{P}_1(\theta)$ a spectral projection $\mathcal{P}_{(-\infty, E)}(\mathcal{H}(\theta))$, and set $\mathcal{P}_2(\theta) = I - \mathcal{P}_1(\theta)$. We have $\mathcal{P}_j \in C^{\infty}(\mathbb{R}^{n*}/\Gamma^*; \text{End } L_2(\Omega))$. For $j = 1, 2$, define an orthoprojection \mathcal{P}_j in $L_2(\mathbb{R}^{n*}/\Gamma^*; L_2(\Omega))$ by

$$(\mathcal{P}_j u)(x, \theta) = \mathcal{P}_j(\theta) u(x, \theta),$$

and set $\mathcal{L}_j = \mathcal{P}_j(C^{\infty}(\mathbb{R}^{n*}/\Gamma^*; L_2(\Omega)))$.

It follows from (2.3) and Lemma 2.4 that

$$(2.4) \quad N_{-}(t; H - E; W) \geq \mathcal{N}(0; \mathcal{W}^{-1} + t(\mathcal{H} - E)^{-1}; \mathcal{L}_1),$$

which is the estimate from below in Theorem 0.2.

To obtain the estimate from above, we note that it follows from (1.18) and Theorem 5.8 that

$$[\mathcal{W}^{-1}, \mathcal{P}_j] \in L_{\rho, 0}(\mathcal{W}^{-1} \Lambda^{-\rho}; \mathbb{R}^{n*}/\Gamma^*; \text{End } L_2(\Omega));$$

therefore, for $u \in C^{\infty}(\mathbb{R}^{n*}/\Gamma^*; L_2(\Omega))$,

$$\langle \mathcal{W}^{-1} u, u \rangle = \sum_{j=1,2} \langle \mathcal{W}^{-1} u_j, u_j \rangle + \langle K u_1, u_2 \rangle,$$

where $u_j = \mathcal{P}_j u$ and $K \in L_{\rho,0}(W^{-1}\Lambda^{-\rho}; \mathbb{R}^{n^*}/\Gamma^*; \text{End } L_2(\Omega))$.

Set

$$W_{r,s}(x) = W(x)^r \langle x \rangle^s, \quad \mathcal{W}_{r,s} = \mathcal{U} W_{r,s} \mathcal{U}^*.$$

By Theorem 1.6, $\mathcal{W}_{r,s} \in L_{\rho,0}(W_{r,s}; \mathbb{R}^{n^*}/\Gamma^*; \text{End } L_2(\Omega))$; therefore Theorem 5.8 gives

$$K_1 := \mathcal{W}_{1/2,0} K \in L_{\rho,0}(W_{-1/2,-\rho}; \mathbb{R}^{n^*}/\Gamma^*; \text{End } L_2(\Omega)).$$

For any $\epsilon > 0$, there exists $C_\epsilon > 0$ such that

$$|\langle K u_1, u_2 \rangle| \leq \|K_1 u_1\| \|\mathcal{W}_{-1/2,0} u_2\| \leq \epsilon \langle \mathcal{W}_{-1/2,0}^* \mathcal{W}_{-1/2,0} u_2, u_2 \rangle + C_\epsilon \langle K_1^* K_1 u_1, u_1 \rangle.$$

By construction, $\mathcal{W}_{-1/2,0}^* \mathcal{W}_{-1/2,0} = \mathcal{W}^{-1}$; therefore

$$\langle \mathcal{W}^{-1} u, u \rangle \geq \langle (\mathcal{W}^{-1} (1 - \epsilon) u_2, u_2 \rangle + \langle (\mathcal{W}^{-1} - C_\epsilon K_1^* K_1) u_1, u_1 \rangle.$$

Clearly,

$$\langle (\mathcal{H} - E)^{-1} u, u \rangle = \sum_{j=1,2} \langle (\mathcal{H} - E)^{-1} u_j, u_j \rangle,$$

whence

$$\begin{aligned} & \langle (\mathcal{W}^{-1} + t(\mathcal{H} - E)^{-1}) u, u \rangle \\ & \geq \langle (\mathcal{W}^{-1} (1 - \epsilon) + t\mathcal{P}_2 (\mathcal{H} - E)^{-1}) u_2, u_2 \rangle + \langle (\mathcal{W}^{-1} - C_\epsilon K_1^* K_1 + t\mathcal{P}_1 (\mathcal{H} - E)^{-1}) u_1, u_1 \rangle, \end{aligned}$$

and we deduce from (2.3) and Lemmas 2.5, 2.3 that

$$\begin{aligned} N_-(t; H - E; W) & \leq \mathcal{N}(0; \mathcal{W}^{-1} (1 - \epsilon) + t\mathcal{P}_2 (\mathcal{H} - E)^{-1}; \mathcal{L}_2) \\ (2.5) \quad & + \mathcal{N}(0; \mathcal{W}^{-1} - C_\epsilon K_1^* K_1 + t\mathcal{P}_1 (\mathcal{H} - E)^{-1}; \mathcal{L}_1). \end{aligned}$$

Fix $\epsilon \in (0, 1)$. Then $\mathcal{W}^{-1} (1 - \epsilon)$ is positive definite, and since $\mathcal{P}_2 (\mathcal{H} - E)^{-1} \geq 0$, the first term in the right-hand side of (2.5) is zero. Now Theorems 5.8 and 5.9 give

$$K_2 := \mathcal{W}_{1/2,\rho} K_1^* K_1 \mathcal{W}_{1/2,\rho} \in L_{\rho,0}(1; \mathbb{R}^{n^*}/\Gamma^*; \text{End } L_2(\Omega)),$$

and by Theorem 5.10, K_2 is bounded in $L_2(\mathbb{R}^{n^*}/\Gamma^*; L_2(\Omega))$. Hence, there exists $C > 0$ such that

$$-C_\epsilon K_1^* K_1 \geq -C \mathcal{W}_{-1,-2\rho},$$

and due to Lemma 2.3, we can replace $-C_\epsilon K_1^* K_1$ in (2.5) by $-C \mathcal{W}_{-1,-2\rho}$. Thus,

$$N_-(t; H - E; W) \leq \mathcal{N}(0; \mathcal{W}^{-1} - C \mathcal{W}_{-1,-2\rho} + t(\mathcal{H} - E)^{-1}; \mathcal{L}_1),$$

which is the estimate from above in Theorem 0.2.

Theorem 0.2 is proved.

2.3. Proof of Theorem 0.3. By applying Theorems 1.7 and 5.8, we obtain for local symbols of $\mathcal{W}_{11,C}^{-1}$ a representation (0.6) with

$$w_{11,U,\chi}^{-1,-1} \in S_{\rho,0}(W_{-1,-\rho}; \mathbb{R}^n; \text{End } \mathbb{C}^p)$$

and the tail belonging to $S_{\rho,0}(W_{-1,-2\rho}; \mathbb{R}^n; \text{End } \mathbb{C}^p)$. This means that conditions (6.4)–(6.8) of Theorem 6.4 hold with

$$M = \mathbb{R}^{n^*}/\Gamma^*, \quad f(\xi) = W(-\xi)^{-1}, \quad \mu_j(\theta, \xi) = (E - E_j(\theta))W(-\xi)^{-1}, \quad j = 1, \dots, p,$$

and (0.8) is imposed in both Theorems 6.4 and 0.3. Hence, all the conditions of Theorem 6.4 hold, and Theorem 0.2 together with (6.10) gives (0.9).

Theorem 0.3 is proved.

2.4. Proof of Theorem 0.4. If W admits a full asymptotic expansion

$$W(x) = W_{-m}(x) + W_{-m-1}(x) + \cdots, \quad \text{as } |x| \rightarrow +\infty,$$

then Theorem 0.2 is applicable with $\rho = 1$. By applying Theorems 1.7 and 5.8, we find that $\mathcal{W}_{11,C}^{-1}$ is an elliptic classical pdo with the principal symbol having eigenvalues

$$(2.6) \quad \mu_j(\theta, \xi) = (E_j(\theta) - E)W_{-m}(-\xi)^{-1}.$$

Well-known results for elliptic pdo (see e.g. [11, 19]) give

$$(2.7) \quad N(t; \overline{\mathcal{W}_{11,C}^{-1}}) = c_-(H - E; W_{-m})t^{n/m} + O(t^{(n-1)/m}),$$

with the constant in the O -term depending on C .

Thus, under an additional assumption Theorem 0.4 is proved, and the proof in the general case is reduced to this special one as follows. Let $\tilde{W} \in C^\infty(\mathbb{R}^n; \mathbb{R}_+)$ be a function which coincides with W_{-m} outside the unit ball. Then due to (0.10), there exists $C_1 > 0$ such that

$$(2.8) \quad \tilde{W}_{C_1}^{-1} := \tilde{W}^{-1} - C_1 \langle \cdot \rangle^{m-1} \leq W^{-1} \leq W_{-C_1}^{-1} := \tilde{W}^{-1} + C_1 \langle \cdot \rangle^{m-1}.$$

Set $\tilde{\mathcal{W}}_{\pm C_1}^{-1} = \mathcal{U} \tilde{W}_{\pm C_1}^{-1} \mathcal{U}^*$. Using Lemmas 2.1–2.3 and (2.8), and arguing as at the beginning of the proof of Theorem 0.2, we obtain

$$(2.9) \quad \begin{aligned} & \mathcal{N}(0; \tilde{\mathcal{W}}_{-C_1}^{-1} + t(\mathcal{H} - E)^{-1}; C^\infty(\mathbb{R}^{n^*}/\Gamma^*; L_2(\Omega))) \\ & \leq N_-(t; H - E; W) \\ & \leq \mathcal{N}(0; \tilde{\mathcal{W}}_{C_1}^{-1} + t(\mathcal{H} - E)^{-1}; C^\infty(\mathbb{R}^{n^*}/\Gamma^*; L_2(\Omega))). \end{aligned}$$

The functions $\tilde{W}_{\pm C_1}^{-1}$ satisfy the conditions

$$\begin{aligned} |\partial^\alpha \tilde{W}_{\pm C_1}^{-1}(x)| & \leq C_\alpha \langle x \rangle^{m-|\alpha|}, \quad \forall \alpha, \\ \tilde{W}_{\pm C_1}^{-1} & \geq c_2 \langle x \rangle^m \quad \text{for } |x| \geq C_2, \end{aligned}$$

where $c_2 > 0$. Therefore we can argue as in the proof of Theorem 0.2 and obtain

$$(2.10) \quad N(t; \overline{\tilde{\mathcal{W}}_{11,-C_1}^{-1}}) \leq \mathcal{N}(0; \tilde{\mathcal{W}}_{-C_1}^{-1} + t(\mathcal{H} - E)^{-1}; C^\infty(\mathbb{R}^{n^*}/\Gamma^*; L_2(\Omega))),$$

$$(2.11) \quad N(t; \overline{\tilde{\mathcal{W}}_{11,C_2}^{-1}}) \geq \mathcal{N}(0; \tilde{\mathcal{W}}_{C_1}^{-1} + t(\mathcal{H} - E)^{-1}; C^\infty(\mathbb{R}^{n^*}/\Gamma^*; L_2(\Omega))),$$

with some $C_2 > 0$. Here $\overline{\tilde{\mathcal{W}}_{11,\pm C_1}^{-1}}$ are constructed in the way $\overline{\mathcal{W}_{11,C}^{-1}}$ were constructed.

By applying Theorems 1.7 and 5.6 we find that $\tilde{\mathcal{W}}_{11,C}^{-1}$ is an elliptic classical pdo with principal symbol having eigenvalues (2.6). Hence, for $N(t; \overline{\tilde{\mathcal{W}}_{11,C}^{-1}})$ the asymptotic formula (2.7) holds.

Now, (2.7) and (2.9)–(2.11) give (0.11), and Theorem 0.4 is proved.

2.5. Proof of Theorem 0.5. If W admits a full asymptotic expansion then W satisfies conditions (0.1) and (0.2) with $m = m_-, \rho = 1$, and W^{-1} admits the asymptotic expansion

$$W^{-1}(x) = W_{-m}^{-1}(x) - W_{-m}^{-2}(x)W_{-m-1}(x) + \cdots.$$

Hence, Theorems 1.7 and 5.6 say that $\mathcal{W}_{11,C}^{-1}$ is a classical pdo with the first two terms of the asymptotic expansion of the symbol given by (0.13) and (0.14). The principal symbol being positive, we may add a smoothing operator (an operator of order $-\infty$) and obtain a positive definite operator (this leaves both the principal and the next term of asymptotics unchanged); after that, we can consider the counting function of $1/m$ -th power of the (closure of the) operator under consideration. The principal and subprincipal symbols of this power are given by (0.15) and (0.16), and the results of [11] give (0.17) (modulo the reservations mentioned before Theorem 0.5).

Thus, Theorem 0.5 is proved provided W admits a full asymptotic expansion. In the general case we set

$$\tilde{W}_C^{-1} = W_{-m}^{-1} - W_{-m}^{-2}W_{-m-1} - C\langle \cdot \rangle^{m-1-\epsilon},$$

notice that due to (0.12) there exists $C > 0$ such that

$$\tilde{W}_C^{-1} \leq W^{-1} \leq \tilde{W}_{-C}^{-1},$$

and reduce the proof to the one for the special case considered above as in the proof of Theorem 0.4.

3. THE ASYMPTOTICS OF $N_+(t; H - E; W)$

3.1. Proof of Theorem 0.6. Define an isometry $\mathcal{U}_0 : L_2(\mathbb{R}^{n^*}/\Gamma^*; L_2(\Omega)) \rightarrow L_2(\mathbb{R}^{n^*}/\Gamma^*; L_2(\mathbb{R}^n/\Gamma))$ by $u(x, \theta) \mapsto \exp\{-i(\theta, x)\}u(x, \theta)$, and set

$$\mathcal{H}_1 = \mathcal{U}_0 \mathcal{H} \mathcal{U}_0^*.$$

It is multiplication by the operator-valued function

$$\mathcal{H}_1(\theta) = (i\nabla - \theta)^*(i\nabla - \theta) + V : H^2(\mathbb{R}^n/\Gamma) \rightarrow L_2(\mathbb{R}^n/\Gamma).$$

Set $\mathcal{W}_1 = \mathcal{U}_0 \mathcal{W} \mathcal{U}_0^*$. It acts as follows:

$$(\mathcal{W}_1 u)(x, \theta) = \frac{1}{\text{vol } \Omega^*} \sum_{\gamma \in \Gamma} W_1(x + \gamma) \int_{\Omega^*} \exp\{i(\theta' - \theta, x + \gamma)\} u(x, \theta') d\theta',$$

and trivial modifications of the proofs of Theorems 1.6 and 1.7 give

Theorem 3.1. *Let W satisfy (1.1). Then $\mathcal{W}_1 \in L_{\rho,0}(f; \mathbb{R}^{n^*}/\Gamma^*; \text{End } L_2(\mathbb{R}^n/\Gamma))$, and if W satisfies (1.5) then its symbol admits a decomposition*

$$w_1(\xi) = W(-\xi) + r(\xi),$$

where $r \in S_{\rho,0}(f\Lambda^{-\rho}; \mathbb{R}^{n^*}/\Gamma^*; \text{End } L_2(\mathbb{R}^n/\Gamma))$.

Thus, we can use $\mathcal{W}_1, \mathcal{H}_1$ almost as we used \mathcal{W}, \mathcal{H} before, the difference being that \mathcal{H}_1 is not a smooth function on $\mathbb{R}^{n^*}/\Gamma^*$ but all the $\mathcal{H}_1(\theta)$ have the same domain; note that \mathcal{H}_1 is a smooth function on $\overline{\Omega^*}$.

To simplify the notation, we use labels \mathcal{W}, \mathcal{H} for $\mathcal{W}_1, \mathcal{H}_1$.

Lemma 3.2. *Let $V \in L_\infty(\mathbb{R}^n)$ and let E be in a gap of $H = -\Delta + V$.*

Then there exists $C_1 > 0$ such that

$$(3.1) \quad (-\Delta + 1)^{-1} - C_1(-\Delta + 1)^{-3/2} \leq (H - E)^{-1} \leq (-\Delta + 1)^{-1} + C_1(-\Delta + 1)^{-3/2}.$$

Proof. Set $B = -\Delta + 1$, $W_1 = 1 - E + V$, and consider

$$T := (-\Delta + V - E)^{-1} - B^{-1} = B^{-1}W_1(-\Delta + V - E)^{-1} = B^{-3/4}T_1B^{-3/4},$$

where

$$\begin{aligned} T_1 &:= B^{-1/4}W_1(-\Delta + V - E)^{-1}B^{3/4} \\ &= B^{-1/4}W_1B^{-1/4} + B^{-1/4}W_1TB^{3/4} \\ &= B^{-1/4}W_1B^{-1/4} + B^{-1/4}W_1(\Delta + V - E)^{-1}W_1B^{-1/4}. \end{aligned}$$

We see that T_1 is bounded in $L_2(\mathbb{R}^n)$. Hence there exists $C_1 > 0$ such that

$$|\langle Tu, u \rangle_{L_2}| \leq C_1 \|B^{-3/4}u\|_{L_2}^2,$$

and (3.1) follows. \square

Define an operator \mathcal{A} in $L_2(\mathbb{R}^{n^*}/\Gamma^*; L_2(\mathbb{R}^n/\Gamma))$ by

$$(\mathcal{A}u)(x, \theta) = (-\Delta_x + 1)u(x, \theta),$$

and set $\mathcal{B} = \mathcal{U}_0\mathcal{U}B\mathcal{U}^*\mathcal{U}_0^* = (i\nabla_x - \theta)^2 + 1$. For any $s, m \in \mathbb{R}$,

$$(-\Delta_x + 1)^s, \quad ((i\nabla_x - \theta)^2 + 1)^s : H^m(\mathbb{R}^n/\Gamma) \rightarrow H^{m-2s}(\mathbb{R}^n/\Gamma)$$

are bounded (the latter uniformly in $\theta \in \Omega^*$). Hence there exists $C_2 > 0$ such that $\mathcal{B}^{-3/2} \leq C_2\mathcal{A}^{-3/2}$. Further, similarly to the proof of Lemma 3.2, there exists $C_3 > 0$ such that

$$|\langle (\mathcal{B}^{-1} - \mathcal{A}^{-1})u, u \rangle_{L_2}| \leq C_3 \langle \mathcal{A}^{-3/2}u, u \rangle_{L_2},$$

and using (3.1), we find $C > 0$ such that

$$(3.2) \quad -C\mathcal{A}^{-3/2} + \mathcal{A}^{-1} \leq (\mathcal{H} - E)^{-1} \leq \mathcal{A}^{-1} + C\mathcal{A}^{-3/2}.$$

The Birman-Schwinger principle, Lemma 2.3 and (3.2) show that

$$\begin{aligned} N(0; I - t\mathcal{W}^{1/2}(\mathcal{A}^{-1} - C\mathcal{A}^{-3/2})\mathcal{W}^{1/2}) &\leq N_+(t; H - E; W) \leq \\ (3.3) \quad &\leq N(0; I - t\mathcal{W}^{1/2}(\mathcal{A}^{-1} + C\mathcal{A}^{-3/2})\mathcal{W}^{1/2}). \end{aligned}$$

Using Lemma 2.2, we obtain

$$\begin{aligned} &N(0; I - t\mathcal{W}^{1/2}(\mathcal{A}^{-1} \pm C\mathcal{A}^{-3/2})\mathcal{W}^{1/2}) \\ &= \mathcal{N}(0; \mathcal{W}^{-1} - t(\mathcal{A}^{-1} \pm C\mathcal{A}^{-3/2}); C^\infty(\mathbb{R}^{n^*}/\Gamma^*; L_2(\mathbb{R}^n/\Gamma))). \end{aligned}$$

Let $\mathcal{W}^{-1,0}$ be a symmetric positive invertible scalar pdo with the symbol $W(-\xi)^{-1}$, and $\overline{\mathcal{W}^{-1,0}}$ its closure in $L_2(\mathbb{R}^{n^*}/\Gamma^*)$. Due to (0.2) there exist $C, \rho_1 > 0$ such that

$$W(x)^{-1}\langle x \rangle^{-\rho} \leq CW(x)^{-1-\rho_1}.$$

Theorem 5.13 shows that $(\overline{\mathcal{W}^{-1,0}})^{\pm(1-\rho_1)/2} \in L_{\rho,0}(W^{\mp(1-\rho_1)/2}; \mathbb{R}^{n^*}/\Gamma^*)$, and Theorems 3.1 and 5.6 give

$$(\overline{\mathcal{W}^{-1,0}})^{-(1-\rho_1)/2}(\mathcal{W}^{-1} - \mathcal{W}_{-1,0})(\overline{\mathcal{W}^{-1,0}})^{-(1-\rho_1)/2} \in L_{\rho,0}(1; \mathbb{R}^{n^*}/\Gamma^*).$$

By Theorem 5.10, this operator is bounded in $L_2(\mathbb{R}^{n^*}/\Gamma^*; L_2(\mathbb{R}^n/\Gamma))$. Hence, there exists $C > 0$ such that

$$\overline{\mathcal{W}^{-1,0}} - C(\overline{\mathcal{W}^{-1,0}})^{1-\rho_1} \leq \overline{\mathcal{W}^{-1}} \leq \overline{\mathcal{W}^{-1,0}} + C(\overline{\mathcal{W}^{-1,0}})^{1-\rho_1}.$$

Now Lemma 2.3 and the Birman-Schwinger principle allow us to deduce from (3.3) that

$$\begin{aligned} & N(0; \overline{\mathcal{W}^{-1,0}} + C_2(\overline{\mathcal{W}^{-1,0}})^{1-\rho_1} - t(\mathcal{A}^{-1} - C\mathcal{A}^{-3/2})) \\ & \leq N_+(t; H - E; W) \\ (3.4) \quad & \leq N(0; \overline{\mathcal{W}^{-1,0}} - C_2(\overline{\mathcal{W}^{-1,0}})^{1-\rho_1} - t(\mathcal{A}^{-1} + C\mathcal{A}^{-3/2})), \end{aligned}$$

where $\overline{\mathcal{W}^{-1,0}}$ is regarded as an operator in $L_2(\mathbb{R}^{n^*}/\Gamma^*; L_2(\mathbb{R}^n/\Gamma))$, with scalar symbol.

It is well-known that the counting function of the spectrum of the Laplacian on \mathbb{R}^n/Γ has the asymptotics

$$N(t; -\Delta) = c_0 t^{n/2} + O(t^{(n-1)/2}), \quad \text{as } t \rightarrow +\infty$$

(see, e.g., [5, 19]), where

$$c_0 = (2\pi)^{-n} |v_n| \text{vol } \Omega.$$

Hence, $\lambda_j, j = 1, 2, \dots$, the eigenvalues of $(-\Delta_x + 1)^{-1}$, obey

$$(3.5) \quad \lambda_j = c_0^{2/n} j^{-2/n} + O(j^{-3/n}), \quad \text{as } j \rightarrow +\infty.$$

W admits a representation (0.20), (0.21); therefore Theorem 6.4 gives for $\overline{\mathcal{W}^{-1,0}}$, an operator in $L_2(\mathbb{R}^{n^*}/\Gamma^*)$,

$$(3.6) \quad N(t; \overline{\mathcal{W}^{-1,0}}) = c_1(W_{-2})t^{n/2} + O(t^{n/2-\epsilon_1}), \quad \text{as } t \rightarrow +\infty,$$

for some $\epsilon_1 > 0$, where

$$c_1(W_{-2}) = (2\pi)^{-n} \text{vol } \Omega^* n^{-1} \int_{S_{n-1}} W_{-2}(\xi)^{n/2} dS(\xi).$$

Let $\mu_1 \leq \mu_2 \leq \dots$ be the eigenvalues of an operator $\mathcal{W}^{-1,0}$ in $L_2(\mathbb{R}^{n^*}/\Gamma^*)$, counted with multiplicity. Then (3.6) implies

$$\mu_j^{-1} = c_1(W_{-2})^{2/n} j^{-2/n} + O(j^{-2/n-\epsilon_2}), \quad \text{as } j \rightarrow +\infty,$$

for some $\epsilon_2 > 0$.

Now (3.4) and (3.5) show that there are C_2, C_3 and $C_4 > 0$ such that

$$\begin{aligned} & \text{card}\{(i, j) \mid i \geq 1, j \geq 1, \\ & 1 < t((c_0 j^{-1})^{2/n} - C_2 j^{-3/n})((c_1(W_{-2})i^{-1})^{2/n} - C_3 i^{-2/n-\epsilon_2})\} - C_4 t^{n/2} \\ & \leq N_+(t; H - E; W) \\ & \leq C_4 t^{n/2} + \text{card}\{(i, j) \mid i \geq 1, j \geq 1, \\ (3.7) \quad & 1 < t((c_0 j^{-1})^{2/n} + C_2 j^{-3/n})((c_1(W_{-2})i^{-1})^{2/n} + C_3 i^{-2/n-\epsilon_2})\}. \end{aligned}$$

It is easy to show that

$$\begin{aligned} & \text{card}\{(i, j) \mid i \geq 1, j \geq 1, \\ & 1 - t((c_0 j^{-1})^{2/n} \pm C_2 j^{-3/n})((c_1(W_{-2})i^{-1})^{2/n} \pm C_3 i^{-2/n-\epsilon_2}) < 0\} \\ & = \text{meas}\{(x, y) \mid x > 1, y > 1, 1 - t^{n/2} c_0 c_1(W_{-2}) x^{-1} y^{-1} < 0\} + O(t^{n/2}) \\ & = \frac{n}{2} c_0 c_1(W_{-2}) t^{n/2} \ln t + O(t^{n/2}); \end{aligned}$$

since $\text{vol } \Omega^* \text{vol } \Omega = (2\pi)^n$, we have

$$\frac{n}{2} c_0 c_1(W_{-2}) = c(W_{-2}),$$

where $c(W_{-2})$ is given by (0.23), and (3.7) gives (0.22).

This completes the proof of Theorem 0.6.

Note that as far as V was concerned we used only the condition $V \in L_\infty(\mathbb{R}^n)$.

3.2. Proof of Theorem 0.8. Due to (2.1) and Lemmas 2.2 and 2.5 we have

$$(3.8) \quad N_+(t; H - E; W) = \mathcal{N}(0; W^{-1} - t(H - E)^{-1}; C_0^\infty(\mathbb{R}^n)),$$

and the right-hand side in (3.8) can be estimated as follows.

Fix $\epsilon > 0$ and $\mu \in C_0^\infty(\mathbb{R})$ such that $0 \leq \mu \leq 1$, $\mu(t) = 1$ for $|t| < 1$, $\mu(t) = 0$ for $|t| > 2$. For $c \in [1/4, 2]$, set

$$\tilde{\mu}_{t,\epsilon,c}(x) = \mu(W(x)^{-1}c^{-1}t^{-\epsilon}), \quad \tilde{\mu}_{t,\epsilon,-c} = 1 - \tilde{\mu}_{t,\epsilon,c},$$

$$\mu_{t,\epsilon,\pm c} = \tilde{\mu}_{t,\epsilon,\pm c} \left(\sum_{j=\pm c} \tilde{\mu}_{t,\epsilon,j}^2 \right)^{-1/2},$$

$$W_{t,\epsilon}^{-1}(x) = W(x)^{-1} \tilde{\mu}_{t,\epsilon,-1/4}(x) + t^\epsilon \tilde{\mu}_{t,\epsilon,1/4}.$$

Due to (0.1) and (0.2),

$$|D^\alpha \tilde{\mu}_{t,\epsilon,c}(x)| \leq C_\alpha (|x| + t^{\epsilon/m-})^{-\rho|\alpha|}, \quad \forall \alpha,$$

$$(3.9) \quad |D^\alpha \mu_{t,\epsilon,c}(x)| \leq C_\alpha (|x| + t^{\epsilon/m-})^{-\rho|\alpha|}, \quad \forall \alpha,$$

$$(3.10) \quad |D^\alpha W_{t,\epsilon}^{-1}(x)| \leq C_\alpha W_{t,\epsilon}^{-1}(x) (|x| + t^{\epsilon/m-})^{-\rho|\alpha|}, \quad \forall \alpha.$$

Using (3.9), one obtains

$$[H, \mu_{t,\epsilon,\pm c}] = \sum_{1 \leq j \leq n} a_{t,\epsilon,c,j}(x) \partial_j + a'_{t,\epsilon,c},$$

where

$$|a_{t,\epsilon,c,j}(x)| \leq Ct^{-\epsilon/m-}, \quad |a'_{t,\epsilon,c}(x)| \leq Ct^{-2\epsilon/m-};$$

hence for any $E_0 < \inf \sigma(H)$ there exists $C > 0$ such that

$$\begin{aligned} & | \langle [(H - E)^{-1}, \mu_{t,\epsilon,\pm c}] u, u \rangle_{L_2} | \\ &= | \langle (H - E_0)^{1/2} (H - E)^{-1} [\mu_{t,\epsilon,\pm c}, H] (H - E)^{-1} u, (H - E_0)^{-1/2} u \rangle_{L_2} | \\ (3.11) \quad & \leq Ct^{-\epsilon/m-} \langle (H - E_0)^{-1} u, u \rangle_{L_2}. \end{aligned}$$

Set $u_{t,\epsilon,\pm 1} = \mu_{t,\epsilon,\pm 1} u$. (3.11) gives

$$\begin{aligned} - \langle (H - E)^{-1} u, u \rangle_{L_2} &= - \langle (H - E)^{-1} \sum_{j=\pm 1} \mu_{t,\epsilon,j}^2 u, u \rangle_{L_2} \\ &\geq - \sum_{j=\pm 1} \langle (H - E)^{-1} u_{t,\epsilon,j}, u_{t,\epsilon,j} \rangle_{L_2} - Ct^{-\epsilon/m-} \langle (H - E_0)^{-1} u, u \rangle_{L_2}. \end{aligned}$$

Take $C_0 > C$ and apply the same procedure to $(H - E)^{-1} - C_0 t^{-\epsilon/m-} (H - E_0)^{-1}$. The result is

$$(3.12) \quad - \langle (H - E)^{-1} u, u \rangle_{L_2} \geq - \sum_{j=\pm 1} \langle ((H - E)^{-1} + C_0 t^{-\epsilon/m-} (H - E_0)^{-1}) u_{t,\epsilon,j}, u_{t,\epsilon,j} \rangle_{L_2},$$

for $t > t_0$, if t_0 is sufficiently large.

Using (3.12) and Lemmas 2.5 and 2.3, we obtain

$$\begin{aligned}
 (3.13) \quad & \mathcal{N}(0, W^{-1} - t(H - E)^{-1}; C_0^\infty(\mathbb{R}^n)) \\
 & \leq \mathcal{N}(0, W^{-1} - t((H - E)^{-1} + C_0 t^{-\epsilon/m-}(H - E_0)^{-1}); C_0^\infty(\Omega_{t,\epsilon,5})) \\
 & \quad + \mathcal{N}(0, W^{-1} - t((H - E)^{-1} + C_0 t^{-\epsilon/m-}(H - E_0)^{-1}); C_0^\infty(\Omega_{t,\epsilon,-1})),
 \end{aligned}$$

where

$$\Omega_{t,\epsilon,c} = \{x \mid W(x)^{-1} < ct^\epsilon\}, \quad \Omega_{t,\epsilon,-c} = \{x \mid W(x)^{-1} > ct^\epsilon\}.$$

On $\Omega_{t,\epsilon,-1}$, we have $W^{-1} = W_{t,\epsilon}^{-1}$; hence we can rewrite (3.13) as

$$\begin{aligned}
 (3.14) \quad & \mathcal{N}(0; W^{-1} - t(H - E)^{-1}; C_0^\infty(\mathbb{R}^n)) \\
 & \leq \mathcal{N}(0, W^{-1} - t((H - E)^{-1} + C_0 t^{-\epsilon/m-}(H - E_0)^{-1}); C_0^\infty(\Omega_{t,\epsilon,5})) \\
 & \quad + \mathcal{N}(0, W_{t,\epsilon}^{-1} - t((H - E)^{-1} + C_0 t^{-\epsilon/m-}(H - E_0)^{-1}); C_0^\infty(\mathbb{R}^n)).
 \end{aligned}$$

Similarly, there exists $C_0 > 0$ such that

$$\begin{aligned}
 (3.15) \quad & \mathcal{N}(0; W_{t,\epsilon}^{-1} - t((H - E)^{-1} - C_0 t^{-\epsilon/m-}(H - E_0)^{-1}); C_0^\infty(\mathbb{R}^n)) \\
 & \leq \mathcal{N}(0; W_{t,\epsilon}^{-1} - t(H - E)^{-1}; C_0^\infty(\Omega_{t,\epsilon,5})) + \mathcal{N}(0; W_{t,\epsilon}^{-1} - t(H - E)^{-1}; C_0^\infty(\Omega_{t,\epsilon,-1})) \\
 & \leq \mathcal{N}(0; W_{t,\epsilon}^{-1} - t(H - E)^{-1}; C_0^\infty(\Omega_{t,\epsilon,5})) + \mathcal{N}(0; W^{-1} - t(H - E)^{-1}; C_0^\infty(\mathbb{R}^n)).
 \end{aligned}$$

There exist $C_1, c_1 > 0$ such that $W^{-1}(x) \geq c_1 > 0$, and

$$\begin{aligned}
 W^{-1} - t((H - E)^{-1} + C_0 t^{-\epsilon/m-}(H - E_0)^{-1}) & \geq c_1 - C_1 t(-\Delta + 1)^{-1}, \\
 W_{t,\epsilon}^{-1} - t(H - E)^{-1} & \geq c_1 - C_1 t(-\Delta + 1)^{-1},
 \end{aligned}$$

and Lemma 2.3 gives

$$\begin{aligned}
 (3.16) \quad & \mathcal{N}(0; W^{-1} - t((H - E)^{-1} + C_0 t^{-\epsilon/m-}(H - E_0)^{-1}); C_0^\infty(\Omega_{t,\epsilon,5})) \\
 & \leq \mathcal{N}(0; c_1 - C_1 t(-\Delta + 1)^{-1}; C_0^\infty(\Omega_{t,\epsilon,5})),
 \end{aligned}$$

$$\begin{aligned}
 (3.17) \quad & \mathcal{N}(0; W_{t,\epsilon}^{-1} - t(H - E)^{-1}; C_0^\infty(\Omega_{t,\epsilon,5})) \leq \mathcal{N}(0; c_1 - C_1 t(-\Delta + 1)^{-1}; C_0^\infty(\Omega_{t,\epsilon,5})).
 \end{aligned}$$

But $(-\Delta + 1)^{-1}$ is a classical elliptic pdo of order -2 ; hence Theorem 6.2 gives

$$\begin{aligned}
 (3.18) \quad & \mathcal{N}(0; c_1 - C_1 t(-\Delta + 1)^{-1}; C_0^\infty(\Omega_{t,\epsilon,5})) \\
 & \leq C_2 \text{meas } \Omega_{t,\epsilon,6} t^{n/2} \leq C_3 t^{n/2+n\epsilon/m}.
 \end{aligned}$$

(See also [14], where asymptotic formulae for pdo of negative order were derived from Theorem 6.2.) By gathering (3.14)–(3.18) we obtain

$$\begin{aligned}
 (3.19) \quad & -C_3 t^{n/2+n\epsilon/m} + \mathcal{N}(0; W_{t,\epsilon}^{-1} - t((H - E)^{-1} - C_0 t^{-\epsilon/m-}(H - E_0)^{-1}); C_0^\infty(\mathbb{R}^n)) \\
 & \leq \mathcal{N}(0; W^{-1} - t(H - E)^{-1}; C_0^\infty(\mathbb{R}^n)) \\
 & \leq C_3 t^{n/2+n\epsilon/m} + \mathcal{N}(0; W_{t,\epsilon}^{-1} - t((H - E)^{-1} + C_0 t^{-\epsilon/m-}(H - E_0)^{-1}); C_0^\infty(\mathbb{R}^n)).
 \end{aligned}$$

We choose $\epsilon > 0$ so that $n/2 + n\epsilon/m < n/m$; this is possible due to the condition $m \in (0, 2)$.

3.3. Using Lemmas 2.5 and 2.2, we obtain

$$\begin{aligned}
 (3.20) \quad & \mathcal{N}(0; W_{t,\epsilon}^{-1} - t((H - E)^{-1} \pm C_0 t^{-\epsilon/m-} (H - E_0)^{-1}); C_0^\infty(\mathbb{R}^n)) \\
 &= \mathcal{N}(0; I - t(W_{t,\epsilon}^{-1})^{-1/2}((H - E)^{-1} \pm C_0 t^{-\epsilon/m-} (H - E_0)^{-1})(W_{t,\epsilon}^{-1})^{-1/2}; C_0^\infty(\mathbb{R}^n)) \\
 &= \mathcal{N}(0; I - t(W_{t,\epsilon}^{-1})^{-1/2}((H - E)^{-1} \pm C_0 t^{-\epsilon/m-} (H - E_0)^{-1})(W_{t,\epsilon}^{-1})^{-1/2}; L_2(\mathbb{R}^n)) \\
 &= \mathcal{N}(0; I - t(\mathcal{W}_{t,\epsilon}^{-1})^{-1/2}((\mathcal{H} - E)^{-1} \pm C_0 t^{-\epsilon/m-} (\mathcal{H} - E_0)^{-1})(\mathcal{W}_{t,\epsilon}^{-1})^{-1/2}; \\
 &\quad L_2(\mathbb{R}^{n^*}/\Gamma^*; L_2(\mathbb{R}^n/\Gamma))) \\
 &= \mathcal{N}(0; I - t(\mathcal{W}_{t,\epsilon}^{-1})^{-1/2}((\mathcal{H} - E)^{-1} \pm C_0 t^{-\epsilon/m-} (\mathcal{H} - E_0)^{-1})(\mathcal{W}_{t,\epsilon}^{-1})^{-1/2}; \\
 &\quad C^\infty(\mathbb{R}^{n^*}/\Gamma^*; L_2(\mathbb{R}^n/\Gamma))) \\
 &= \mathcal{N}(0; \mathcal{W}_{t,\epsilon}^{-1} - t((\mathcal{H} - E)^{-1} \pm C_0 t^{-\epsilon/m-} (\mathcal{H} - E_0)^{-1}); C^\infty(\mathbb{R}^{n^*}/\Gamma^*; L_2(\mathbb{R}^n/\Gamma))),
 \end{aligned}$$

where $\mathcal{W}_{t,\epsilon}^{-1} = \mathcal{U}W_{t,\epsilon}^{-1}\mathcal{U}^*$.

Due to (3.10), the derivatives of $W_{t,\epsilon}^{-1}$ decay as $t \rightarrow +\infty$. Therefore the same estimates as in Section 2 now give

1) for $\phi, \psi \in C^\infty(\mathbb{R}^{n^*}/\Gamma^*)$ with $\text{supp } \phi \cap \text{supp } \psi = \emptyset$, $\psi \mathcal{W}_{t,\epsilon}^{-1} \psi$ is an integral operator with kernel $K_{t,\epsilon}(\theta, \theta')$ for all N, α, β ,

$$(3.21) \quad \|D_\theta^\alpha D_{\theta'}^\beta K_{t,\epsilon}(\theta, \theta')\|_{\text{End } L_2(\mathbb{R}^n/\Gamma)} \leq C_{\alpha,\beta,N} t^{-N};$$

2) for a local chart $\kappa : U \rightarrow U^1$ with small $\text{diam } U$ and $\phi, \psi \in C_0^\infty(U^1)$,

$$(3.22) \quad \phi \mathcal{W}_{t,\epsilon}^{-1} \psi = (\kappa^{-1})^*(\phi \circ \kappa)(W_{t,\epsilon}^{-1}(-D) + W_{t,\epsilon}^{-1,-1})(\psi \circ \kappa) \kappa^*,$$

where $W_{t,\epsilon}^{-1,-1}$ is a pdo with symbol satisfying the estimates

$$(3.23) \quad \|D_\xi^\alpha D_\theta^\beta w_{t,\epsilon}^{-1,-1}(\theta, \xi)\|_{\text{End } L_2(\mathbb{R}^n/\Gamma)} \leq C_{\alpha,\beta} t^{-\epsilon/m-} W_{t,\epsilon}^{-1}(-\xi)(|\xi| + t^{\epsilon/m-})^{-\rho|\alpha|},$$

for all α, β .

Identify Ω^* with $\mathbb{R}^{n^*}/\Gamma^*$, fix $\epsilon_1 \in (0, \rho\epsilon/2m_-)$, and cut Ω^* into boxes of size $c_0 t^{-\epsilon_1}$ with $1 < c_0 < 2$. We set

$$U_{t,\epsilon,j}^+ = \{x \mid \text{dist}(x, U_{t,\epsilon,j}) < t^{-2\epsilon_1}\},$$

$$U_{t,\epsilon,j}^- = \{x \in U_{t,\epsilon,j} \mid \text{dist}(x, \partial U_{t,\epsilon,j}) > t^{-2\epsilon_1}\},$$

and construct a partition of unity

$$(3.24) \quad \sum_j \phi_{t,\epsilon,j}^2(\theta) = 1, \quad \forall \theta \in \mathbb{R}^{n^*}/\Gamma^*,$$

with the following properties:

$$(3.25) \quad |D^\alpha \phi_{t,\epsilon,j}(\theta)| \leq C_{\alpha,\epsilon} t^{2\epsilon_1|\alpha|}, \quad \forall \alpha,$$

$$(3.26) \quad 0 \leq \phi_{t,\epsilon,j} \leq 1, \quad \phi_{t,\epsilon,j} = 1 \text{ on } U_{t,\epsilon,j}^-, \quad \text{supp } \phi_{t,\epsilon,j} \subset U_{t,\epsilon,j}^+.$$

Let $\kappa_j : U_j \rightarrow U_j^1$ be local charts such that $\kappa_j^{-1} \circ \kappa_l$ are restrictions of orthogonal operators and $U_{t,\epsilon,j}^+ \subset U_j$. Consider

$$K_{t,\epsilon,j} = [\mathcal{W}_{t,\epsilon}^{-1}, \phi_{t,\epsilon,j}] \phi_{t,\epsilon,j}.$$

Clearly, (3.21) for $\mathcal{W}_{t,\epsilon}^{-1}$ yields the same estimate for $K_{t,\epsilon,j}$; therefore we may consider $K_{t,\epsilon,j}$ in appropriate local charts. Now, (3.10), (3.23) and (3.25) mean that (in the notation of Section 6)

$$W_{t,\epsilon}^{-1}(-D) \in \Psi(W_{t,\epsilon}^{-1}(\cdot, -); g_{t,\epsilon}), \quad W_{t,\epsilon}^{-1,-1} \in \Psi(t^{-\epsilon/m-} W_{t,\epsilon}^{-1}(\cdot, -); g_{t,\epsilon}),$$

$$\phi_{t,\epsilon,j} \in \Psi(1, g_{t,\epsilon}),$$

uniformly in $t \geq 1$, where

$$g_{t,\epsilon;\theta,\xi}(\theta', \xi') = (|\xi| + t^{\epsilon/m-})^{-2\rho} |\xi'|^2 + t^{4\epsilon_1} |\theta'|^2$$

is a metric on \mathbb{R}^{2n} . One easily verifies that when $\epsilon_1 \in (0, \epsilon\rho/2m_-)$ the metric $g_{t,\epsilon}$ is σ -temperate, $W_{t,\epsilon}^{-1}(\cdot, -)$ is a σ, g -temperate function, and

$$h_{t,\epsilon}(\theta, \xi) := (|\xi| + t^{\epsilon/m})^{-\rho} t^{2\epsilon_1} \leq t^{-\epsilon_2},$$

where $\epsilon_2 = \epsilon\rho/m_- - 2\epsilon_1 > 0$. Now Theorem 5.2 gives

$$K_{t,\epsilon,j} \in \Psi(t^{-\epsilon_2} W_{t,\epsilon}^{-1}(\cdot, -); g_{t,\epsilon}).$$

Using once again Theorem 5.2, Theorem 5.13 and (3.21), we obtain for $\psi_j = 1$ on $\text{supp } \phi_{t,\epsilon,j}$

$$\begin{aligned} & (\overline{\mathcal{W}_{t,\epsilon}^{-1}})^{-1/2} K_{t,\epsilon,j} (\overline{\mathcal{W}_{t,\epsilon}^{-1}})^{-1/2} \\ &= t^{-\epsilon_2} (\kappa_j^{-1})^* (\psi_j \circ \kappa_j) K_{t,\epsilon,j}^0 (\phi_j \circ \kappa_j) \kappa_j^* + T_{t,\epsilon,j}^{-\infty}, \end{aligned}$$

where $T_{t,\epsilon,j}^{-\infty}$ satisfies (3.21) and $K_{t,\epsilon,j}^0 \in \Psi(1; g_{t,\epsilon})$ uniformly in $t \geq 1$. Theorem 5.4 (the L_2 -boundedness theorem) provides $C > 0$ such that

(3.27)

$$\| (\overline{\mathcal{W}_{t,\epsilon}^{-1}})^{-1/2} K_{t,\epsilon,j} (\overline{\mathcal{W}_{t,\epsilon}^{-1}})^{-1/2} \|_{\text{End } L_2(\mathbb{R}^{n*}/\Gamma^*; L_2(\mathbb{R}^n/\Gamma))} \leq C t^{-\epsilon_2},$$

and since the number of $\phi_{t,\epsilon,j}$ is $O(t^{n\epsilon_1})$, we obtain

(3.28)

$$\| (\overline{\mathcal{W}_{t,\epsilon}^{-1}})^{-1/2} \sum_j K_{t,\epsilon,j} (\overline{\mathcal{W}_{t,\epsilon}^{-1}})^{-1/2} \|_{\text{End } L_2(\mathbb{R}^{n*}/\Gamma^*; L_2(\mathbb{R}^n/\Gamma))} \leq C_1 t^{n\epsilon_1 - \epsilon_2}.$$

Choose ϵ_1 so that $n\epsilon_1 - (\epsilon\rho/m - 2\epsilon_1) \leq -3\epsilon_1$. Then (3.28) gives

$$|\langle \sum_j K_{t,\epsilon,j} u, u \rangle_{L_2}| \leq C_1 t^{-3\epsilon_1} \langle \mathcal{W}_{t,\epsilon}^{-1} u, u \rangle_{L_2},$$

and we have

$$\begin{aligned} \langle \mathcal{W}_{t,\epsilon}^{-1} u, u \rangle_{L_2} &= \langle \mathcal{W}_{t,\epsilon}^{-1} (1 - t^{-2\epsilon_1}) \sum_j \phi_{t,\epsilon,j}^2 u, u \rangle_{L_2} + t^{-2\epsilon_1} \langle \mathcal{W}_{t,\epsilon}^{-1} u, u \rangle_{L_2} \\ (3.29) \quad &\geq (1 - t^{-2\epsilon_1}) \sum_j \langle \mathcal{W}_{t,\epsilon}^{-1} \phi_{t,\epsilon,j} u, \phi_{t,\epsilon,j} u \rangle_{L_2}. \end{aligned}$$

Similarly to (3.27), we deduce from (3.22) and (3.23) that

$$\begin{aligned} & \langle \mathcal{W}_{t,\epsilon}^{-1} \phi_{t,\epsilon,j} u, \phi_{t,\epsilon,j} u \rangle_{L_2(\mathbb{R}^{n*}/\Gamma^*; L_2(\mathbb{R}^n/\Gamma))} \\ & \geq (1 - C t^{-\epsilon/m-}) \langle W_{t,\epsilon}^{-1}(-D) u_j, u_j \rangle_{L_2(U_{t,\epsilon,j}^+; L_2(\mathbb{R}^n/\Gamma))}, \end{aligned}$$

where $u_j = \kappa_j^* \phi_{t,\epsilon,j} u$; therefore (3.29) yields

$$(3.30) \quad \langle \mathcal{W}_{t,\epsilon}^{-1} u, u \rangle_{L_2} \geq (1 - t^{-\epsilon_1}) \sum_j \langle W_{t,\epsilon}^{-1}(-D) u_j, u_j \rangle_{L_2}$$

provided t is large enough.

Fix $\theta_j = \theta_{t,\epsilon,j} \in U_{t,\epsilon,j}^-$. Due to (3.26) there exists $t_0 > 0$ such that for $t \geq t_0$,

$$(3.31) \quad \begin{aligned} & -\langle ((\mathcal{H} - E)^{-1} + C_0 t^{-\epsilon/m-} (\mathcal{H} - E_0)^{-1}) \phi_{t,\epsilon,j} u, \phi_{t,\epsilon,j} u \rangle_{L_2} \\ & \geq -\langle ((\mathcal{H}(\theta_j) - E)^{-1} + t^{-\epsilon_1} (\mathcal{H}(\theta_j) - E_0)^{-1}) \phi_{t,\epsilon,j} u, \phi_{t,\epsilon,j} u \rangle_{L_2}. \end{aligned}$$

Set

$$\begin{aligned} \mathcal{B}_{t,\epsilon,j}^{\pm} &= (\mathcal{H}(\theta_j) - E)^{-1} \pm t^{-\epsilon_1} (\mathcal{H}(\theta_j) - E_0)^{-1}, \\ \mathcal{A}_{t,\epsilon,j}^{\pm} &= (1 \mp t^{-\epsilon_1}) W_{t,\epsilon}^{-1}(-D) - t \mathcal{B}_{t,\epsilon,j}^{\pm}. \end{aligned}$$

It follows from (3.30), (3.31) and Lemmas 2.5 and 2.3 that

$$(3.32) \quad \begin{aligned} & \mathcal{N}(0; \mathcal{W}_{t,\epsilon}^{-1} - t((\mathcal{H} - E)^{-1} + C_0 t^{-\epsilon/m-} (\mathcal{H} - E_0)^{-1}); C^\infty(\mathbb{R}^{n^*}/\Gamma^*; L_2(\mathbb{R}^n/\Gamma))) \\ & \leq \sum_j \mathcal{N}(0; \mathcal{A}_{t,\epsilon,j}^+; C_0^\infty(U_{t,\epsilon,j}^+; L_2(\mathbb{R}^n/\Gamma))). \end{aligned}$$

Since $\phi_{t,\epsilon,j} = 1$ on $U_{t,\epsilon,j}^-$, for

$$u = (u_j) \in \bigoplus_j C_0^\infty(U_{t,\epsilon,j}^-; L_2(\Omega)) \subset C^\infty(\mathbb{R}^{n^*}/\Gamma^*; L_2(\Omega))$$

we have

$$u = \sum_j \phi_{t,\epsilon,j} u = \sum_j \phi_{t,\epsilon,j} u_j$$

Therefore we can obtain, similarly to (3.32),

$$(3.33) \quad \begin{aligned} & \mathcal{N}(0; \mathcal{W}_{t,\epsilon}^{-1} - t((\mathcal{H} - E)^{-1} - C_0 t^{-\epsilon/m-} (\mathcal{H} - E_0)^{-1}); C^\infty(\mathbb{R}^{n^*}/\Gamma^*; L_2(\mathbb{R}^n/\Gamma))) \\ & \geq \sum_j \mathcal{N}(0; \mathcal{A}_{t,\epsilon,j}^-; C_0^\infty(U_{t,\epsilon,j}^-; L_2(\mathbb{R}^n/\Gamma))). \end{aligned}$$

3.4. Set

$$\begin{aligned} B_{t,\epsilon,j,k}^{\pm} &= (E_k(\theta_j) - E)^{-1} \pm t^{-\epsilon_1} (E_k(\theta_j) - E_0)^{-1}, \\ A_{t,\epsilon,j,k}^{\pm} &= (1 \mp t^{-\epsilon_1}) W_{t,\epsilon}^{-1}(-D) - t B_{t,\epsilon,j,k}^{\pm}. \end{aligned}$$

Since $L_2(\Omega)$ can be decomposed into an orthogonal sum of eigenspaces of $\mathcal{H}(\theta_j)$, we deduce from Lemma 2.5 that

$$(3.34) \quad \mathcal{N}(0; \mathcal{A}_{t,\epsilon,j}^{\pm}; C_0^\infty(U_{t,\epsilon,j}^{\pm}; L_2(\Omega))) = \sum_k \mathcal{N}(0; \mathcal{A}_{t,\epsilon,j,k}^{\pm}; C_0^\infty(U_{t,\epsilon,j}^{\pm})).$$

Due to (3.10) the symbol of $A_{t,\epsilon,j,k}^{\pm}$ satisfies

$$|D_\xi^\alpha a_{t,\epsilon,j,k}^{\pm}(\xi)| \leq C_\alpha (|\xi| + t^{\epsilon/m-})^{-\rho|\alpha|} (W_{t,\epsilon}^{-1}(-\xi) + t(E_k(\theta_j) - E_0)^{-1}), \quad \forall \alpha,$$

uniformly in $t \geq 1, k, j$; therefore Theorem 6.1 is applicable. It tells us that for any $\delta \in (0, 2/3)$ there exist $t_0, c, C, C_1 > 0$ such that for all $t \geq t_0, k, j$,

$$(3.35) \quad \begin{aligned} & |\mathcal{N}(0; \mathcal{A}_{t,\epsilon,j,k}^\pm; C_0^\infty(U_{t,\epsilon,j}^\pm)) - F_{t,\epsilon,k}(U_{t,\epsilon,j}^\pm)| \\ & \leq C(F_{t,\epsilon,k,c,\delta}(U_{t,\epsilon,j}^\pm) - F_{t,\epsilon,k,-c,\delta}(U_{t,\epsilon,j}^\pm)) + C_1 F_{t,\epsilon,k,c,\delta}(\partial U_{t,\epsilon,j,c,\delta}^\pm), \end{aligned}$$

where

$$\begin{aligned} F_{t,\epsilon,k,c,\delta}(U) &= (2\pi)^{-n} \text{meas}\{(\theta, \xi) \in U \times \mathbb{R}^n \mid \\ (1 - t^{-\epsilon_1})W_{t,\epsilon}^{-1}(-\xi) - tB_{t,\epsilon,j,k}^\pm &< ct^{-\rho\delta\epsilon/m-}(W_{t,\epsilon}^{-1}(-\xi) + t(E_k(\theta_j) - E_0)^{-1})\}, \\ F_{t,\epsilon,k}(U) &= F_{t,\epsilon,k,0,0}, \\ \partial U_{t,\epsilon,j,c,\delta}^\pm &= \{x \mid \text{dist}(x, \partial U_{t,\epsilon,j}^\pm) < ct^{-\rho\delta\epsilon/m-}\}. \end{aligned}$$

We may choose ϵ_1 so that $2\rho\epsilon/3m_- > \epsilon_1$; then (3.35) yields

$$(3.36) \quad \begin{aligned} & |\mathcal{N}(0; \mathcal{A}_{t,\epsilon,j,k}^\pm; C_0^\infty(U_{t,\epsilon,j}^\pm)) - F_{t,\epsilon,k,0}^1(U_{t,\epsilon,j})| \\ & \leq C(F_{t,\epsilon,k,c}^1(U_{t,\epsilon,j}) - F_{t,\epsilon,k,-c}^1(U_{t,\epsilon,j})) + C_1 F_{t,\epsilon,k,c}^1(\partial U_{t,\epsilon,j,c}) \end{aligned}$$

(with new $c, C, C_1 > 0$), where

$$\begin{aligned} F_{t,\epsilon,k,c}^1(U) &= (2\pi)^{-n} \text{meas}\{(\theta, \xi) \in U \times \mathbb{R}^n \mid \\ (1 - t^{-\epsilon_1})W_{t,\epsilon}^{-1}(-\xi) - t(E(\theta_j) - E)^{-1} &< ct^{-\epsilon_1}(W_{t,\epsilon}^{-1}(-\xi) + t(E_k(\theta_j) - E_0)^{-1})\}, \\ \partial U_{t,\epsilon,j,c} &= \{x \mid \text{dist}(x, \partial U_{t,\epsilon,j}) < ct^{-\epsilon_1}\}. \end{aligned}$$

Note that $F_{t,\epsilon,k,c}^1(U) = 0$ if $k \leq p$ (i.e. if $E_k(\theta) - E < 0 \quad \forall \theta$), and that if $k \geq p+1$ (and hence $E_k(\theta) - E > 0 \quad \forall \theta$), then there exists $c_1 = c_1(c) > 0$ such that

$$F_k^2(t - c_1 t^{1-\epsilon_1}; U) \leq F_{t,\epsilon,k,-c}^1(U), \quad F_k^2(t + c_1 t^{1-\epsilon_1}; U) \geq F_{t,\epsilon,k,c}^1(U),$$

where $F_k^2(t; U) = (2\pi)^{-n} \text{meas}\{(\theta, \xi) \in U \times \mathbb{R}^n \mid W_{t,\epsilon}^{-1}(-\xi) < t(E_k(\theta) - E)^{-1}\}$. Further, $\text{meas} \partial U_{t,\epsilon,j,c} \leq C_1 t^{-\epsilon_1} \text{meas} U_{t,\epsilon,j}$, and therefore

$$F_{t,\epsilon,k,c}^1(\partial U_{t,\epsilon,j,c}) \leq C_2 t^{-\epsilon_1} F_k^2(t + c_1 t^{1-\epsilon_1}; U_{t,\epsilon,j}).$$

Now we can rewrite (3.36) as

$$(3.37) \quad \begin{aligned} & \mathcal{N}(0; \mathcal{A}_{t,\epsilon,j,k}^\pm; C_0^\infty(U_{t,\epsilon,j}^\pm)) \\ & = F_k^2(t; U_{t,\epsilon,j})(1 + O(t^{-\epsilon_1})) + O(F_k^2(t + c_1 t^{1-\epsilon_1}; U_{t,\epsilon,j}) - F_k^2(t - c_1 t^{1-\epsilon_1}; U_{t,\epsilon,j})), \end{aligned}$$

with the c_1 and the constants in O -terms independent of k, j and $t \geq t_0$. Note that the RHS in (3.37) is zero if $k \leq p$ and t is large.

3.5. Set

$$F_\epsilon(t) = \sum_{j,k} F_k^2(t; U_{t,\epsilon,j}).$$

It follows from (3.8), (3.19), (3.20), (3.32), (3.33) and (3.37) that

$$(3.38) \quad N_+(t; H - E; W) = F_\epsilon(t)(1 + O(t^{-\epsilon_1})) \\ + O(F_\epsilon(t + ct^{1-\epsilon_1}) - F_\epsilon(t - ct^{1-\epsilon_1})) + O(t^{n/2+n\epsilon/m-}).$$

Now, with some $C > 0$,

$$(3.39) \quad F_\epsilon(t) = (2\pi)^{-n} \sum_{k \geq p+1} \text{meas}\{(\theta, \xi) \in \mathbb{R}^{n^*}/\Gamma^* \times \mathbb{R}^n \mid W_{t,\epsilon}^{-1}(-\xi) < t(E_k(\theta) - E)^{-1}\} \\ = (2\pi)^{-n} \sum_{k \geq p+1} \text{meas}\{(\theta, \xi) \in \mathbb{R}^{n^*}/\Gamma^* \times \mathbb{R}^n \mid W^{-1}(-\xi) < t(E_k(\theta) - E)^{-1}\} \\ + O\left(\sum_{k \geq p+1} \text{meas}\{(\theta, \xi) \in \mathbb{R}^{n^*}/\Gamma^* \times \mathbb{R}^n \mid |\xi| \leq Ct^{\epsilon/m}, \quad 1 < t(E_k(\theta) - E)^{-1}\}\right),$$

because if C is sufficiently large then $W^{-1}(-\xi) = W_{t,\epsilon}^{-1}(-\xi)$ for $|\xi| \geq Ct^{\epsilon/m}$.

The first term in the RHS of (3.39) is equal to $c_+(t; H - E; W)$ in (0.28), and the O -term admits an upper bound via

$$C_1 t^{n\epsilon/m} \text{card}\{k \mid k^{2/n} < C_2 t\} \leq C_3 t^{n\epsilon/m+n/2},$$

since $E_k(\theta) \sim \text{const} k^{2/n}$ as $k \rightarrow +\infty$.

Thus, (3.38) and (3.39) give

$$(3.40) \quad N_+(t; H - E; W) = c_+(t; H - E; W)(1 + O(t^{-\epsilon_1})) \\ + O(c_+(t + ct^{1-\epsilon_1}; H - E; W) - c_+(t - ct^{1-\epsilon_1}; H - E; W)) + O(t^{n/2+n\epsilon/m}).$$

Under condition (0.19), there exists $c_0 > 0$ such that

$$c_+(t; H - E; W) \geq c_0 t^{n/m},$$

and since we have chosen $\epsilon > 0$ such that $n/2 + n\epsilon/m < n/m$, the last term in (3.40) is included in the O -term of (0.29). The second term in the RHS of (3.40) is included in the O -term of (0.29) due to (0.8).

Theorem 0.8 is proved.

3.6. Proof of Theorem 0.7. It follows from (0.24) and (0.25) that

$$(3.41) \quad c_+(H - E; W_{-m})t^{n/m} - c_+(t; H - E; W) = O(t^{-\omega+n/m}),$$

with some $\omega > 0$. It follows that $c_+(t; H - E; W)$ satisfies (0.8); (0.19) also follows from (0.24) and (0.25). We see that all the conditions of Theorem 0.8 are satisfied and hence (0.29) holds. Clearly, (0.29) and (3.41) give (0.26).

Theorem 0.7 is proved.

4. THE CASE OF NON-NEGATIVE W

4.1. Here we prove Theorem 0.9.

This time we have to work with (2.1) because a reduction to both (2.3) and (3.8) is, in general, impossible.

Fix $\epsilon > 0$ and $\nu \in (0, \rho)$, take $\mu \in C_0^\infty(\mathbb{R})$ as in the previous section, and for $c \in [1/4, 4]$, set

$$\begin{aligned}\tilde{\mu}_{t,\epsilon,0,c}(x) &= \mu(|x|c^{-1}t^{-\epsilon}), \\ \tilde{\mu}_{t,\epsilon,-1,c}(x) &= (1 - \mu(|x|c^{-1}t^{-\epsilon}))\mu(W(x)\langle x \rangle^{m+\nu}c^{-1}), \\ \tilde{\mu}_{t,\epsilon,1,c}(x) &= (1 - \mu(|x|c^{-1}t^{-\epsilon}))(1 - \mu(W(x)\langle x \rangle^{m+\nu}c^{-1})), \\ \mu_{t,\epsilon,j,c} &= \tilde{\mu}_{t,\epsilon,j,c} \left(\sum_{j=0,\pm 1} \tilde{\mu}_{t,\epsilon,j,c}^2 \right)^{-1/2}, \\ W_{t,\epsilon,0}(x) &= \tilde{\mu}_{t,\epsilon,0,2}(x)W(x), \\ W_{t,\epsilon,-1}(x) &= (1 - \mu(|x|4t^{-\epsilon}))\mu(W(x)\langle x \rangle^{m+\nu}/4)W(x), \\ W_{t,\epsilon,1}(x) &= \langle x \rangle^{-m-\nu}\mu(W(x)\langle x \rangle^{m+\nu}/4) + (1 - \mu(W(x)\langle x \rangle^{m+\nu}/4))W(x), \\ U_{t,\epsilon,0} &= \{x \mid |x| < 2t^\epsilon\}, \quad U_{t,\epsilon,-1} = \{x \mid |x| > t^\epsilon/2, W(x) < 2\langle x \rangle^{-m-\nu}\}, \\ U_{t,\epsilon,1} &= \{x \mid |x| > t^\epsilon/2, W(x) > \langle x \rangle^{-m-\nu}/2\}.\end{aligned}$$

We have

$$(4.1) \quad W_{t,\epsilon,j}(x) = W(x) \quad \forall x \in U_{t,\epsilon,j},$$

and since the derivatives of $\tilde{\mu}_{t,\epsilon,j,c}, \mu_{t,\epsilon,j,c}$ are supported on a set where $|x| > t^\epsilon/2$, we deduce from (0.30) that

$$(4.2) \quad \begin{aligned}|D^\alpha \tilde{\mu}_{t,\epsilon,j,c}(x)| &\leq C_\alpha(|x| + t^\epsilon)^{(\nu-\rho)|\alpha|}, \quad \forall \alpha, \\ |D^\alpha \mu_{t,\epsilon,j,c}(x)| &\leq C_\alpha(|x| + t^\epsilon)^{(\nu-\rho)|\alpha|}, \quad \forall \alpha,\end{aligned}$$

$$(4.3) \quad |D^\alpha W_{t,\epsilon,1}(x)| \leq C_\alpha W_{t,\epsilon,1}(x)(|x| + t^\epsilon)^{(\nu-\rho)|\alpha|}, \quad \forall \alpha.$$

4.2. Take $\epsilon_1 \in (0, \epsilon(\rho - \nu))$, and set

$$u_{t,\epsilon,j}(x) = \mu_{t,\epsilon,j,1}(x)u(x), \quad B_{t,\epsilon,\pm C_0} = (H - E)^{-1} \pm C_0 t^{-\epsilon_1} (H - E_0)^{-1}.$$

Using (4.2) and the equality

$$\sum_{j=\pm 1,0} \mu_{t,\epsilon,j,1}^2 = 1,$$

we obtain, similarly to (3.12),

$$(4.4) \quad \mp \langle (H - E)^{-1}u, u \rangle_{L_2} \geq \mp \sum_{j=0,-1,1} \langle B_{t,\epsilon,\pm C_0} u_{t,\epsilon,j}, u_{t,\epsilon,j} \rangle_{L_2},$$

provided $C_0 > 0$ is large enough. Using (4.4), (4.1) and Lemmas 2.5, 2.3 and 2.4, we deduce from (2.1) the upper bound

$$\begin{aligned}& N_\pm(t; H - E; W) \\ & \leq \sum_{j=-1,0,1} \mathcal{N}(0; I \mp W_{t,\epsilon,j}^{1/2} B_{t,\epsilon,\pm C_0} W_{t,\epsilon,j}^{1/2}; C_0^\infty(U_{t,\epsilon,j}^+)) \\ & \leq \sum_{j=-1,0} \mathcal{N}(0; I \mp W_{t,\epsilon,j}^{1/2} B_{t,\epsilon,\pm C_0} W_{t,\epsilon,j}^{1/2}; C_0^\infty(U_{t,\epsilon,j}^+)) \\ & \quad + \mathcal{N}(0; I \mp W_{t,\epsilon,1}^{1/2} B_{t,\epsilon,\pm C_0} W_{t,\epsilon,1}^{1/2}; C_0^\infty(\mathbb{R}^n)).\end{aligned}\tag{4.5}$$

Similarly, there exists $C_0 > 0$ such that

$$\begin{aligned}
 (4.6) \quad & \mathcal{N}(0; I \mp W_{t,\epsilon,1}^{1/2} B_{t,\epsilon,\mp C_0} W_{t,\epsilon,1}^{1/2}; C_0^\infty(\mathbb{R}^n)) \\
 & \leq \sum_{j=-1,0} \mathcal{N}(0; I \mp W_{t,\epsilon,j}^{1/2} (H - E)^{-1} W_{t,\epsilon,j}^{1/2}; C_0^\infty(U_{t,\epsilon,j}^+)) \\
 & \quad + \mathcal{N}(0; I \mp t W_{t,\epsilon,1}^{1/2} (H - E)^{-1} W_{t,\epsilon,1}^{1/2}; L_2(\mathbb{R}^n)) \\
 & = \sum_{j=-1,0} \mathcal{N}(0; I \mp W_{t,\epsilon,j}^{1/2} (H - E)^{-1} W_{t,\epsilon,j}^{1/2}; C_0^\infty(U_{t,\epsilon,j}^+)) + N_\pm(t; H - E; W).
 \end{aligned}$$

4.3. (4.3) and the estimate

$$\langle x \rangle^{-m-\nu}/2 \leq W_{t,\epsilon,1}(x) \leq C \langle x \rangle^{-m}$$

mean that $W_{t,\epsilon,1}$ satisfies conditions (0.1) and (0.2) (with $\rho - \nu > 0$ instead of ρ) uniformly in $t \geq 1$. Since $W_{t,\epsilon,1}$ differs from W only where $W(x) \leq C \langle x \rangle^{-m-\nu}$, we have

$$(4.7) \quad c_-(t; H - E; W) - c_-(t; H - E; W_{t,\epsilon,1}) = O(t^{n/m-\omega})$$

with some $\omega > 0$. Further, if W satisfies (0.19), there exists $c_0 > 0$ such that

$$(4.8) \quad c_-(t; H - E; W) \geq c_0 t^{n/m};$$

therefore if $c_-(t; H - E; W)$ satisfies (0.8) then $c_-(t; H - E; W_{t,\epsilon,1})$ satisfies it as well. Thus, $W_{t,\epsilon,1}$ satisfies all the conditions of Theorem 0.3, and the proof of this theorem gives

$$\mathcal{N}(0; I + t W_{t,\epsilon,1}^{1/2} (H - E)^{-1} W_{t,\epsilon,1}^{1/2}; L_2(\mathbb{R}^n)) = c_-(t; H - E; W_{t,\epsilon,1})(1 + O(t^{-\omega_1})),$$

where $\omega_1 > 0$. Using (4.7), we obtain, with some $\omega_2 > 0$

$$\mathcal{N}(0; I + t W_{t,\epsilon,1}^{1/2} (H - E)^{-1} W_{t,\epsilon,1}^{1/2}; L_2(\mathbb{R}^n)) = c_-(t; H - E; W)(1 + O(t^{-\omega_1})).$$

Since

$$(4.9) \quad (E_j(\theta) - E)^{-1} \pm C_0 t^{-\epsilon_1} (E_j(\theta) - E_0)^{-1} = (E_j(\theta) - E)^{-1} (1 + O(t^{-\epsilon_1})),$$

we can similarly obtain, for any $C_0 > 0$,

$$(4.10) \quad \mathcal{N}(0; I + t W_{t,\epsilon,1}^{1/2} B_{t,\epsilon,\pm C_0} W_{t,\epsilon,1}^{1/2}; L_2(\mathbb{R}^n)) = c_-(t; H - E; W)(1 + O(t^{-\omega_1})).$$

4.4. By inserting (4.10) into (4.5) and (4.6) we see that (0.9) for W satisfying (0.8), (0.19) and (0.30) will be proved when we show that there exist $C, \omega > 0$ such that for $j = 0, -1$

$$(4.11) \quad \mathcal{N}(0; I + t W_{t,\epsilon,1}^{1/2} B_{t,\epsilon,\pm C_0} W_{t,\epsilon,1}^{1/2}; C_0^\infty(U_{t,\epsilon,j})) \leq C t^{n/m-\omega},$$

$$(4.12) \quad \mathcal{N}(0; I + t W_{t,\epsilon,1}^{1/2} (H - E)^{-1} W_{t,\epsilon,1}^{1/2}; C_0^\infty(U_{t,\epsilon,j})) \leq C t^{n/m-\omega}.$$

Let P_1 be a spectral projection $P_{(-\infty, E)}(H)$, and set $P_2 = I - P_1$. If t_0 is sufficiently large and $t \geq t_0$, then $P_2 B_{t,\epsilon,\pm C_0} \geq 0$ due to (4.9), and of course $P_2(H - E)^{-1} \geq 0$. It follows that for any N there exists $C_N > 0$ such that

$$\begin{aligned}
 B_{t,\epsilon,\pm C_0} & \geq P_1 B_{t,\epsilon,\pm C_0} \geq (1 + C_1 t^{-\epsilon_1}) P_1 (H - E)^{-1} \geq -C_N (H - E_0)^{-N}, \\
 (H - E)^{-1} & \geq -C_N (H - E_0)^{-N}.
 \end{aligned}$$

Hence, we may prove (4.11) and (4.12) with $-C_N(H-E_0)^{-N}$ instead of $B_{t,\epsilon,-C_0}$ and $(H-E)^{-1}$, respectively.

Using Lemma 2.4, the Birman-Schwinger principle and Lemma 2.3, we see that it suffices to prove the estimates

$$(4.13) \quad \mathcal{N}(0; (H-E)^N - C_{1,N}t\langle x \rangle^{-m-\nu}; C_0^\infty(\mathbb{R}^n)) \leq Ct^{n/2-\omega},$$

$$(4.14) \quad \mathcal{N}(0; (H-E)^N - C_{1,N}t\chi_{U_{t,\epsilon,0}}; C_0^\infty(\mathbb{R}^n)) \leq Ct^{n/2-\omega},$$

where $\chi_{U_{t,\epsilon,0}}$ is the characteristic function of $U_{t,\epsilon,0}$. Upper bounds for the LHS in (4.13) and (4.14) are well-known (see e.g. [19]); one can also apply Lemma 2.3 and the general Theorems 6.1 and 6.2, and obtain

$$\begin{aligned} & \mathcal{N}(0; (H-E)^N - C_{1,N}t\langle x \rangle^{-m-\nu}; C_0^\infty(\mathbb{R}^n)) \\ & \leq C \text{meas}\{(x, \xi) \in \mathbb{R}^{2n} \mid (|\xi| + 1)^{2N} < C_{2,N}t\langle x \rangle^{-m-\nu}\} \leq C_{3,N}t^{n/2N+n/(m+\nu)}, \\ & \mathcal{N}(0; (H-E)^N - C_{1,N}t\chi_{U_{t,\epsilon,0}}; C_0^\infty(\mathbb{R}^n)) \\ & \leq C_{3,N}t^{n/2N} \text{meas } U_{t,\epsilon,0} \leq C_{4,N}t^{n/2N+n\epsilon}. \end{aligned}$$

Since $\nu > 0$, we can choose $\epsilon > 0$ and N so that (4.13) and (4.14) hold.

4.5. Thus, the first statement of Theorem 0.9, i.e. an analogue of Theorem 0.3 for non-negative W , has been proved, and analogues of Theorems 0.7 and 0.8 are proved similarly. The proof of the fact that the $j = 1$ terms in (4.5) and (4.6) give the principal terms in the asymptotic formulae remains the same, and to finish the proof, one has to obtain the bounds

$$(4.15) \quad \mathcal{N}(0; I - tW_{t,\epsilon,1}^{1/2}B_{t,\epsilon,\pm C_0}W_{t,\epsilon,1}^{1/2}; C_0^\infty(U_{t,\epsilon,j})) \leq Ct^{n/m-\omega},$$

$$(4.16) \quad \mathcal{N}(0; I - tW_{t,\epsilon,1}^{1/2}(H-E)^{-1}W_{t,\epsilon,1}^{1/2}; C_0^\infty(U_{t,\epsilon,j})) \leq Ct^{n/m-\omega},$$

for $j = -1, 0$.

Clearly, there exists $C > 0$ such that

$$-B_{t,\epsilon,\pm C_0} \geq -C(-\Delta + 1)^{-1}, \quad -(H-E)^{-1} \geq -C(-\Delta + 1)^{-1};$$

therefore we may prove (4.15) and (4.16) with $(-\Delta + 1)^{-1}$ instead of $B_{t,\epsilon,\pm C_0}$ and $(H-E)^{-1}$, respectively. Now we use Lemma 2.4, the Birman-Schwinger principle and Lemma 2.3, which show that it suffices to obtain the estimates

$$(4.17) \quad \mathcal{N}(0; -\Delta + 1 - Ct\langle x \rangle^{-m-\nu}; C_0^\infty(\mathbb{R}^n)) \leq C_1t^{n/m-\omega},$$

$$(4.18) \quad \mathcal{N}(0; -\Delta + 1 - Ct\chi_{U_{t,\epsilon,0}}; C_0^\infty(\mathbb{R}^n)) \leq C_1t^{n/m-\omega}.$$

Upper bounds for the LHS in (4.17) and (4.18) are well-known (see e.g. [19]). One can also apply the general Theorems 6.1 and 6.2, and obtain

$$\begin{aligned} & \mathcal{N}(0; -\Delta + 1 - Ct\langle x \rangle^{-m-\nu}; C_0^\infty(\mathbb{R}^n)) \\ & \leq C \text{meas}\{(x, \xi) \in \mathbb{R}^{2n} \mid (|\xi| + 1)^2 < C_{2,t}\langle x \rangle^{-m-\nu}\} \leq C_3t^{n/(m+\nu)}, \\ & \mathcal{N}(0; -\Delta + 1 - Ct\chi_{U_{t,\epsilon,0}}; C_0^\infty(\mathbb{R}^n)) \leq C_3t^{n/2} \text{meas } U_{t,\epsilon,0} \leq C_4t^{n/2+n\epsilon}. \end{aligned}$$

Since $m \in (0, 2)$ and $\nu > 0$, we can choose $\epsilon > 0$ such that (4.17) and (4.18) hold.

Thus, the proof of analogues of Theorems 0.7 and 0.8 for $W \geq 0$ satisfying (0.30) is completed.

4.6. To obtain an analogue of Theorem 0.6, one has to be more careful because an estimate (4.19) is of no use now: recall that the principal term of the asymptotics is of order $t^{n/2} \ln t$.

So, we have to repeat all the constructions of this section using $\ln \ln t$ instead of t^ϵ . Then in the RHS of (4.19) we have

$$C_4 t^{n/2} (\ln \ln t)^{n\epsilon} = o(t^{n/2} \ln t),$$

and all the errors in other formulae will also be of order $o(t^{n/2} \ln t)$.

4.7. **Singular perturbations.** To prove Theorem 0.10, it is sufficient to note that due to the condition (0.31), $\mathcal{W}_1 \in L_{\rho,0}(W\Lambda^{-\epsilon}; \mathbb{R}^{n*}/\Gamma^*; \text{End } L_2(\Omega))$; therefore $\mathcal{W} + \mathcal{W}_1$ and \mathcal{W} have exactly the same properties as far as the proofs above are concerned.

5. SOME FACTS OF THE THEORY OF PSEUDO DIFFERENTIAL OPERATORS

5.1. **Pdo on \mathbb{R}^n .** We need several basic definitions of a general calculus of pdo [10, Section 18] (see also [2]).

Let σ be the standard symplectic form on $\mathbb{R}_X^{2n} = \mathbb{R}_x^n \times \mathbb{R}_\xi^n$.

A metric g on \mathbb{R}^{2n} is called σ -temperate if and only if there exist $C, c > 0$ such that

- a) $g_X(X - Y) < c$ implies $g_Y \leq Cg_X$;
- b) $g_Y(\cdot) \leq Cg_X(\cdot)(1 + g_Y^\sigma(X - Y))^C, \quad \forall X, Y$.

Here

$$g_X^\sigma(Y) = \sup_{Z \neq 0} \frac{\sigma(Y, Z)^2}{g_X(Z)}.$$

We also require

$$h(X) := \sup_{Y \neq 0} \frac{g_X(Y)}{g_X^\sigma(Y)} \leq 1, \quad \forall X.$$

A function $p : \mathbb{R}^{2n} \rightarrow \mathbb{R}_+$ is called σ, g -temperate if and only if there exist $C, c > 0$ such that

- a) $g_X(X - Y) \leq c$ implies $p(X) \leq Cp(Y)$;
- b) $p(X) \leq Cp(Y)(1 + g_Y^\sigma(X - Y))^C, \quad \forall X, Y$.

Below, H_j stands for a Hilbert space, and we set $B_{jk} = \text{Hom}(H_j, H_k)$.

A function $a \in C^\infty(\mathbb{R}^{2n}; B_{12})$ is said to belong to $S(p; g; B_{12})$, if and only if for any k ,

$$\begin{aligned} & n_k(a; p; g) \\ & := \sup_X \sup_{Y_j \neq 0} |(d^k a)(X; Y_1, \dots, Y_k)| p(X)^{-1} g_X(Y_1)^{-1/2} \dots g_X(Y_k)^{-1/2} < \infty, \end{aligned}$$

where $d^k a$ is the k -th differential.

$S(p; g; B_{12})$ is a Fréchet space with $n_k(\cdot; p; g)$ as seminorms.

If $H_1 = H_2 = \mathbb{C}$, we omit B_{12} in the notation and write $S(p; g)$.

For our purposes, it suffices to use metrics which split:

$$g_{x,\xi}(y, \eta) = g'_{x,\xi}(y) + g''_{x,\xi}(\eta),$$

and some of the results formulated below hold for such metrics only.

Given $a \in S(p; g; B_{12})$, one can define a pdo A with the (left) symbol a as follows:

$$(Au)(x) = (2\pi)^{-n} \int_{\mathbb{R}^n} d\xi \int_{\mathbb{R}^n} dy \exp\{i(x - y, \xi)\} a(x, \xi) u(y).$$

Example 5.1. Let $\rho \in (0, 1]$, and let f satisfy (1.2) and (1.3). Then metric

$$g_{x,\xi}(y, \eta) = |y|^2 + \langle \xi \rangle^{-2\rho} |\eta|^2$$

is σ -temperate, f is a σ, g -temperate function, and the class $S(f; g)$ consists of functions satisfying the estimates

$$|D_x^\beta D_\xi^\alpha a(x, \xi)| \leq C_{\alpha, \beta} f(\xi) \langle \xi \rangle^{-\rho|\alpha|}, \quad \forall \alpha.$$

The first statement can be easily verified (see e.g. [10, Section 18]), and the second one was, in fact, established when we proved Lemma 1.1. The last statement is evident.

Thus, the class $L_{\rho, 0}(f; \mathbb{R}^n; B)$ introduced in Section 2, is a Hörmander class.

In particular, $L_{\rho, 0}(\Lambda^m; \mathbb{R}^n; B)$ is nothing but Hörmander class $L_{\rho, 0}^m(\mathbb{R}^n; B)$.

Here are the main theorems of the calculus of pdo on \mathbb{R}^n .

Theorem 5.1. *A acts continuously from $S(\mathbb{R}^n; H_1)$ into $S(\mathbb{R}^n; H_2)$ and admits a unique continuous extension to an operator from $S'(\mathbb{R}^n; H_1)$ into $S'(\mathbb{R}^n; H_2)$ which is also denoted by A.*

We write $A \in \Psi(p; g; B_{12})$.

If $H_1 = H_2 = \mathbb{C}$, we omit B_{12} in the notation.

Theorem 5.2. *Let $A \in \Psi(p; g; B_{23})$ and $B \in \Psi(q; g; B_{12})$ be pdo with the symbols a, b respectively.*

Then $AB \in \Psi(pq; g; B_{13})$, and for any N its symbol c admits the decomposition

$$c(x, \xi) = \sum_{|\alpha| \leq N-1} (\alpha!)^{-1} (\partial_\xi^\alpha a(x, \xi)) (D_x^\alpha b(x, \xi)) + r_N(x, \xi),$$

where $r_N \in S(pqh^N; g; B_{13})$, and the map

$$S(p; g; B_{23}) \times S(q; g; B_{12}) \ni (a, b) \mapsto r_N \in S(pqh^N; g; B_{13})$$

is bounded.

Theorem 5.3. *Let $A \in \Psi(p; g; B_{12})$ be a pdo with the symbol a .*

Then its formal adjoint $A^ \in \Psi(p; g; B_{21})$, and for any N its symbol admits the decomposition*

$$a^*(x, \xi) = \sum_{|\alpha| \leq N-1} (\alpha!)^{-1} i^{|\alpha|} D_x^\alpha D_\xi^\alpha a(x, \xi)^* + r_N(x, \xi),$$

where $r_N \in S(ph^N; g; B_{21})$.

Theorem 5.4. *$A \in \Psi(1; g; B_{12})$ maps $L_2(\mathbb{R}^n; H_1)$ into $L_2(\mathbb{R}^n; H_2)$, and there exist C, N , which depend only on n and the constants characterizing p and g but not on a , such that*

$$\|A\|_{L_2(\mathbb{R}^n; H_1) \rightarrow L_2(\mathbb{R}^n; H_2)} \leq C \max_{k \leq N} n_k(a; p; g).$$

Theorem 5.5. *Let $m_0 < m_1 < \dots < m_k < \dots$, where $m_k \rightarrow +\infty$ as $k \rightarrow +\infty$, and let $a_j \in S(ph^{m_j}; g; B_{12})$.*

Then there exists $a \in S(ph^{m_0}; g; B_{12})$ such that

$$a \sim \sum_{j \geq 0} a_j$$

in the sense that for any N

$$a - \sum_{m_j < N} a_j \in S(ph^N; g; B_{12}).$$

In Section 2, we also used the following trivial generalization of Theorem 5.5.

Theorem 5.6. *Let $m_{j0} < m_{j1} < \dots < m_{jk} < \dots$, where $m_{jk} \rightarrow +\infty$ as $k \rightarrow +\infty$, and let $m_{00} < m_{10} < \dots < m_{k0} < \dots$, where $m_{k0} \rightarrow +\infty$ as $k \rightarrow +\infty$.*

Let $a \sim \sum_{j \geq 0} a_j$ with $a_j \in S(ph^{m_{jk}}; g; B_{12})$ ($j = 0, 1, \dots$) admitting decompositions $a_{jk} \sim \sum_{k \geq 0} a_{jk}$, where $a_{jk} \in S(ph^{m_{jk}}; g; B_{12})$.

Then $a \sim \sum_{j,k \geq 0} a_{jk}$ in the sense that for any N

$$a - \sum_{m_{jk} < N} a_{jk} \in S(ph^N; g; B_{12}).$$

In Section 2, we also use the following construction which is justified by Theorem 5.5.

Let $a \in S_{\rho,0}(f; \mathbb{R}^{n^*}/\Gamma^*; B_{12})$, let

$$L(s, t) = \sum_{\alpha, \beta} L_{\alpha, \beta} s^\alpha t^\beta \in \mathbb{C}[[s, t]] = \mathbb{C}[[s_1, \dots, s_n; t_1, \dots, t_n]]$$

be a formal power series such that for any N , only finitely many $L_{\alpha, \beta}$ with $|\alpha| < N$ are non-zero, and let P be an operator acting continuously in each $S_{\rho,0}(f\Lambda^k; \mathbb{R}^n; B_{12})$, $k \in \mathbb{R}$.

Then Theorem 5.5 allows one to consider the symbol

$$a_L = L(D_\xi, P)a,$$

which is understood in the sense that

$$a_L - \sum_{|\alpha| < N, \beta} L_{\alpha, \beta} D_\xi^\alpha P^\beta a \in S_{\rho,0}(f\Lambda^{-N\rho}; \mathbb{R}^n; B_{12}).$$

Using theorems for the classes $L_{\rho,0}^m(\mathbb{R}^n; B)$, one deduces theorems for pdo on a compact manifold M without boundary, of the classes $L_{\rho,0}^m(M; B)$ (see e.g. [10, Section 16]).

In just the same manner one can deduce from theorems for $L_{\rho,0}(f; \mathbb{R}^n; B)$ theorems for the classes $L_{\rho,0}(f; \mathbb{R}^{n^*}/\Gamma^*; B)$ introduced in Section 1.

In the following subsection, we list theorems which are used in the paper.

5.2. Pdo on a torus.

Theorem 5.7. *$A \in L_{\rho,0}(f; \mathbb{R}^{n^*}/\Gamma^*; B_{12})$ acts continuously from $C^\infty(\mathbb{R}^{n^*}/\Gamma^*; H_1)$ into $C^\infty(\mathbb{R}^{n^*}/\Gamma^*; H_2)$, and admits a unique continuous extension to an operator from $\mathcal{D}'(\mathbb{R}^{n^*}/\Gamma^*; H_1)$ to $\mathcal{D}'(\mathbb{R}^{n^*}/\Gamma^*; H_2)$.*

Theorem 5.8. *Let $A \in L_{\rho,0}(f_1; \mathbb{R}^{n^*}/\Gamma^*; B_{23})$ and $B \in L_{\rho,0}(f_2; \mathbb{R}^{n^*}/\Gamma^*; B_{12})$.*

Then $AB \in L_{\rho,0}(f_1 f_2; \mathbb{R}^{n^}/\Gamma^*; B_{13})$, and its local symbols are expressed via the local symbols of A, B by means of Theorem 5.2.*

Theorem 5.9. *Let $A \in L_{\rho,0}(f; \mathbb{R}^{n^*}/\Gamma^*; B_{12})$.*

Then its formal adjoint $A^ \in L_{\rho,0}(f; \mathbb{R}^{n^*}/\Gamma^*; B_{21})$, and its local symbols are expressed via the local symbols of A by means of Theorem 5.3.*

Theorem 5.10. $A \in L_{\rho,0}(1; \mathbb{R}^{n^*}/\Gamma^*; B_{12}) : L_2(\mathbb{R}^{n^*}/\Gamma^*; H_1) \rightarrow L_2(\mathbb{R}^{n^*}/\Gamma^*; H_2)$ is bounded.

Theorem 5.11. Let the local symbols of $A \in L_{\rho,0}(f; \mathbb{R}^{n^*}/\Gamma^*; B_{12})$ satisfy the following condition: there exist C, C_1 such that

$$\|a(\theta, \xi)^{-1}\| \leq Cf(-\xi), \quad \forall |\xi| > C, \theta,$$

and let the A be invertible.

Then $A^{-1} \in L_{\rho,0}(f^{-1}; \mathbb{R}^{n^*}/\Gamma^*; B_{12})$.

Theorem 5.12. Let f satisfy (1.2) and (1.3) with $-m_- > 0$, and let local symbols of $A \in L_{\rho,0}(f; \mathbb{R}^{n^*}/\Gamma^*; \text{End } \mathbb{C}^m)$ satisfy

$$a(x, \xi) + a(x, \xi)^* > cf(-\xi), \quad \forall |\xi| > C, \theta.$$

Let $A : C^\infty(\mathbb{R}^{n^*}/\Gamma^*; \mathbb{C}^m) \rightarrow C^\infty(\mathbb{R}^{n^*}/\Gamma^*; \mathbb{C}^m)$ be symmetric.

Then A is essentially self-adjoint, and its closure is a semibounded (from below) operator with discrete spectrum.

Remark 5.1. Theorem 5.12 is also valid for pdo acting in sections of a finite-dimensional fibering over M .

Theorem 5.13. Let $A \in L_{\rho,0}(f; \mathbb{R}^{n^*}/\Gamma^*)$ be a positive definite operator satisfying the conditions of Theorem 5.12, and let \bar{A} be its closure in $L_2(\mathbb{R}^{n^*}/\Gamma^*)$.

Then $(\bar{A})^c \in L_{\rho,0}(f^c; \mathbb{R}^{n^*}/\Gamma^*)$ for any $c \in \mathbb{R}$.

Remark 5.2. Theorem 5.13 is also valid for pdo on \mathbb{R}^n [2]; in fact, it is proved similarly to the corresponding theorem in [2].

Theorem 5.14. Let f satisfies (1.2) and (1.3).

Then there exists an invertible pdo $A \in L_{\rho,0}(f; \mathbb{R}^{n^*}/\Gamma^*)$ with

$$A^{-1} \in L_{\rho,0}(f^{-1}; \mathbb{R}^{n^*}/\Gamma^*).$$

Remark 5.2. Once again this theorem is valid for pdo on \mathbb{R}^n as well, and is deduced from its \mathbb{R}^n -counterpart.

6. SOME GENERAL THEOREMS ON SPECTRAL ASYMPTOTICS

6.1. In Chapter 7 of [16] several general bounds for $\mathcal{N}(0; A_t; C_0^\infty(\Omega_t; \mathbb{C}^m))$ were obtained. We shall use only two of them, and not in their full generality.

In both, A_t is a symmetric pdo with the symbol $a_t \in S(p_t; g_t; \text{End } \mathbb{C}^m)$, g_t is a σ -temperate splitting metric, and p_t is a σ, g_t -temperate function (all the conditions are satisfied uniformly in $t \geq 1$).

Also in both theorems, use was made of the Weyl symbol of A_t ; one can use $a_t^0 := (a_t + a_t^*)/2$ instead (this follows from the more detailed formulation in [16] and a formula expressing the Weyl symbol via the left one).

In the formulations of the theorems, one uses the following functions, depending on constants $c > 0, \delta > 0$, and a set $U \subset \mathbb{R}^{2n}$:

$$U_t(g_t; c; \delta) = \{(x, \xi) \mid \inf_{(y, \eta) \in U} g_{t; x, \xi}(x - y, \xi - \eta) < ch_t(x, \xi)^\delta\},$$

$$V_{\pm c, \delta}(a_t^0; U) = (2\pi)^{-n} \sum_{j=1}^m \text{meas}\{(x, \xi) \in U \mid \lambda_j(x, \xi) \leq \pm ch_t(x, \xi)^\delta p_t(x, \xi)\},$$

where the λ_j are the eigenvalues of a_t^0 .

Theorem 6.1. *Let A_t be a symmetric pdo with the symbol $a_t \in S(p_t; g_t; \text{End } \mathbb{C}^m)$, g_t a σ -temperate splitting metric, and p_t a σ, g_t -temperate function (all the conditions are satisfied uniformly in $t \geq 1$). Let*

$$\max_{\mathbb{R}^{2n}} h_t(x, \xi) \rightarrow 0 \quad \text{as } t \rightarrow +\infty,$$

and let there exist $c > 0$ and $R(t)$ such that

$$(6.1) \quad a_t^0(x, \xi) := (a_t(x, \xi) + a_t(x, \xi)^*)/2 \geq cp_t(x, \xi), \quad \forall |x| + |\xi| > R(t).$$

Then for any $\delta \in (0, 1/3)$, there exist $t_0 > 0, c > 0, C$ such that

$$(6.2) \quad \begin{aligned} & |\mathcal{N}(0; A_t; C_0^\infty(\Omega; \mathbb{C}^m)) - V_{0,0}(a_t^0; \Omega \times \mathbb{R}^n)| \\ & \leq C(V_{c,\delta}(a_t^0; \Omega \times \mathbb{R}^n) - V_{-c,\delta}(a_t^0; \Omega \times \mathbb{R}^n)) + CV_{c,\delta}(a_t^0; (\partial\Omega \times \mathbb{R}^n)_t(g_t; c; \delta)) \\ & \quad + C \int \int h_t(x, \xi)^\delta dx d\xi, \end{aligned}$$

with the integral over $(x, \xi) \in (\partial\Omega \times \mathbb{R}^n)_t(g_t; c; \delta)$ satisfying

$$\lambda_1(x, \xi) \leq ch_t(x, \xi)^\delta p_t(x, \xi).$$

If $m = 1$, then the last term in (6.2) may be dropped and one may take any $\delta \in (0, 2/3)$.

Theorem 6.2. *Let A_t be a symmetric pdo with the symbol $a_t \in S(p_t; g_t; \text{End } \mathbb{C}^m)$, g_t a σ -temperate splitting metric, and p_t a σ, g_t -temperate function (all the conditions are satisfied uniformly in $t \geq 1$). Let there exist $C, \epsilon > 0$ such that*

$$h_t(x, \xi) \leq C(1 + |x| + |\xi|)^{-\epsilon}, \quad \forall x, \xi,$$

and let (6.1) hold.

Then for any $\delta \in (0, 1/3)$, there exist $t_0 > 0, c > 0, C$ such that

$$(6.3) \quad \begin{aligned} & |\mathcal{N}(0; A_t; C_0^\infty(\Omega; \mathbb{C}^m)) - V_{0,0}(a_t^0; \Omega \times \mathbb{R}^n)| \\ & \leq C(V_{c,\delta}(a_t^0; \Omega \times \mathbb{R}^n) - V_{-c,\delta}(a_t^0; \Omega \times \mathbb{R}^n)) + CV_{c,\delta}(a_t^0; (\partial\Omega \times \mathbb{R}^n)_t(g_t; c; \delta)) \\ & \quad + C \int \int h_t(x, \xi)^\delta dx d\xi + C, \end{aligned}$$

with the integral over $(x, \xi) \in (\partial\Omega \times \mathbb{R}^n)_t(g_t; c; \delta)$ satisfying

$$\lambda_1(x, \xi) \leq ch_t(x, \xi)^\delta p_t(x, \xi).$$

If $m = 1$, then the integral in (6.3) may be dropped and one may take any $\delta \in (0, 2/3)$.

6.2. The statements of Theorems 6.1 and 6.2 can be reformulated for pdo on a compact manifold without boundary, and the proofs remain essentially the same, being essentially local (even microlocal). The basic ingredient—a construction of an approximate spectral projection \mathcal{E} —is a microlocal procedure, as the proof that the symbol of \mathcal{E} satisfies necessary estimates; these estimates mean that \mathcal{E} belongs to a certain class of pdo. Now it is a class of pdo on M ; once this fact is established, one has to replace the calculus of pdo on \mathbb{R}^n by the one on M and use the same microlocal considerations as in [16].

One can obtain theorems of varying generality; we state one which suffices for our needs.

Let $\rho \in (0, 1]$, and let $f : \mathbb{R}^n \rightarrow \mathbb{R}_+$ satisfy the following two conditions.

a) There exists $C > 0$ such that

$$(6.4) \quad |(\nabla f)(x)| \leq C f(x) \langle x \rangle^{-\rho}.$$

b) There exist $d_1 > 0, d, m > 0, m_1$ such that

$$(6.5) \quad d_1 \langle x \rangle^{m_1} \leq f(x) \leq d_1 \langle x \rangle^m.$$

Let \mathcal{L} be a p -dimensional Hermitian fibering over a torus M , and let A be a symmetric pdo acting in $C^\infty(M; \mathcal{L})$, with local symbols

$$(6.6) \quad a_{U, \chi} \in S_{\rho, 0}(f; \mathbb{R}^n; \text{End } \mathbb{C}^p).$$

Let the eigenvalues λ_j of the local symbols satisfy the estimate

$$(6.7) \quad \text{Re } \lambda_j(x, \xi) \geq c f(-\xi) \quad \text{for } |\xi| > C,$$

where $C, c > 0$ depend on a chart and a trivialization (one can show that the condition itself is independent of this choice).

Then an analogue of Theorem 5.12 for pdo acting in sections of a finite-dimensional fibering over a torus states that A is essentially self-adjoint, and its closure is a semibounded (from below) operator with discrete spectrum.

Let there also exist functions $\mu_j : T^*M \rightarrow \mathbb{R}$, $1 \leq j \leq p$, such that for any local chart and any trivialization, the eigenvalues of the local symbol satisfy

$$(6.8) \quad |\lambda_j(x, \xi) - \mu_j(x, \xi)| \leq C f(-\xi) \langle \xi \rangle^{-\omega},$$

where $C, \omega > 0$.

Let $dx d\xi$ be a volume form on T^*M , and set

$$\begin{aligned} V(t) &= (2\pi)^{-n} \sum_{1 \leq j \leq p} \int \int_{\mu_j(x, \xi) < t} dx d\xi, \\ V_{c, \kappa}(t) &= \sum_{1 \leq j \leq p} \int \int_{\Omega_{j, t, c, \kappa}} dx d\xi, \\ J_{c, \kappa}(t) &= \sum_{1 \leq j \leq p} \int \int_{\Omega_{j, t, c, \kappa}} \langle \xi \rangle^{-\kappa} dx d\xi, \end{aligned}$$

where

$$\Omega_{j, t, c, \kappa} = \{(x, \xi) \mid \mu_j(x, \xi) - t \leq c(t + f(\xi)) \langle \xi \rangle^{-\kappa}\}.$$

Note that if we use local charts $\kappa_j : (\mathbb{R}^n \supset) U_j \rightarrow U_j^1(\subset M)$ such that $\kappa_r^{-1} \circ \kappa_s$ are restrictions of orthogonal operators, then all the functions above are well-defined.

Theorem 6.3. *Let (6.4)–(6.8) hold.*

Then for any $\delta \in (0, 2/3)$ there exist $c > 0, C$ such that

$$(6.9) \quad |N(t; \bar{A}) - V(t)| \leq C(V_{c,\kappa}(t) - V_{-c,\kappa}(t) + J_{c,\kappa}(t)),$$

where $\kappa = \min\{\omega, \delta\rho\}$.

6.3. Under additional conditions one can deduce from Theorem 6.3 an asymptotic formula with a remainder estimate.

Theorem 6.4. *Let (6.4)–(6.8) hold, and let $V(t)$ satisfy the Tauberian condition (0.8).*

Then there exists $\omega > 0$ such that

$$(6.10) \quad N(t; \bar{A}) = V(t)(1 + O(t^{-\omega})).$$

Proof. Due to (6.5) and (6.8), there exist $c_1 > 0, C_1$ such that

$$c_1 t^{n/m_1} \leq V(t) \leq C_1 t^{n/m}.$$

Therefore it suffices to prove that, with some $C_2, \omega > 0$.

$$(6.11) \quad V_{c,\kappa}(t) - V_{-c,\kappa}(t) \leq C_2 t^{n/m_1 - \omega} + C_2 V(t) t^{-\omega},$$

$$(6.12) \quad J_{c,\kappa}(t) \leq C_2 t^{n/m_1 - \omega} + C_2 V(t) t^{-\omega}.$$

Fix $\epsilon \in (0, 1/m_1)$ and set

$$\Omega_{j,t,c,\kappa}^- = \{(x, \xi) \in \Omega_{j,t,c,\kappa} \mid |\xi| < t^\epsilon\},$$

$$\Omega_{j,t,c,\kappa}^+ = \Omega_{j,t,c,\kappa} \setminus \Omega_{j,t,c,\kappa}^-,$$

$$V_{c,\kappa}^\pm(t) = \sum_{1 \leq j \leq p} \int \int_{\Omega_{j,t,c,\kappa}^\pm} dx d\xi,$$

$$J_{c,\kappa}^\pm(t) = \sum_{1 \leq j \leq p} \int \int_{\Omega_{j,t,c,\kappa}^\pm} \langle \xi \rangle^{-\kappa} dx d\xi.$$

Clearly,

$$(6.13) \quad V_{c,\kappa} = V_{c,\kappa}^+ + V_{c,\kappa}^-, \quad J_{c,\kappa} = J_{c,\kappa}^+ + J_{c,\kappa}^-,$$

$$(6.14) \quad J_{c,\kappa}^-(t) \leq C_3 t^{n\epsilon}, \quad V_{\pm c,\kappa}^-(t) \leq C_3 t^{n\epsilon},$$

$$(6.15) \quad J_{c,\kappa}^+ \leq C_3 V(t) t^{-\kappa\epsilon},$$

$$(6.16) \quad V_{c,\kappa}^+(t) \leq V(t + c_2 t^{1-\kappa\epsilon}), \quad V_{-c,\kappa}^+(t) \geq V(t - c_2 t^{1-\kappa\epsilon}) - C_4 t^{n\epsilon},$$

with some $C_3, C_4, c_2 > 0$.

Now, (6.11) follows from (6.13), (6.14), (6.16) and (0.8), while (6.12) follows from (6.13).

Theorem 6.4 is proved. \square

REFERENCES

1. Alama, S., Deift, P.A., Hempel, R.: Eigenvalue branches of the Schrödinger operator $H - \lambda W$ in a gap of $\sigma(H)$, *Commun. Math. Phys.*, 121, 291-321 (1989) MR **90c**:35046
2. Beals, R.: Weighted distribution spaces and pseudo-differential operators. *Journ. Anal. Math.*, 36, 131-189 (1981) MR **83a**:35105
3. Birman, M.Š.: Discrete spectrum in the gaps of a continuous one for perturbations with large coupling constant In: *Estimates and asymptotics for discrete spectra of integral and differential equations. Advances in Soviet mathematics*, vol.7, Providence, RI: AMS, 1991, pp. 57-74 MR **95h**:47009
4. Birman, M.Š.: On the discrete spectrum in the gaps of a perturbed periodic second order operator. *Funct. Anal. Appl.* 25, No 4, 158-161 (1991)
5. Birman, M.Š., Solomyak, M.Z.: Asymptotics of the spectrum of differential equations. *Itogi Nauki: Mat. Anal.*, vol. 14, VINITI, Moscow, 1977, pp. 5-58; English transl. in *J. Soviet Math.* 12 (1979), no. 3. MR **57**:6892
6. Birman, M.Š., Solomyak, M.Z.: *Spectral theory of self-adjoint operators in a Hilbert space.* Leningrad: LGU 1980; English transl., Reidel, Dordrecht, 1987. MR **82k**:47001; MR **93g**:47001
7. Deift, P.A., Hempel, R.: On the existence of eigenvalues of the Schrödinger operator $H - \lambda W$ in a gap of $\sigma(H)$. *Commun. Math. Phys.*, 103, 461-490 (1986) MR **87k**:35184
8. Gesztesy, F., Simon, B.: On a theorem of Deift and Hempel. *Commun. Math. Phys.*, 116, 503-505 (1988) MR **89g**:35080
9. Hempel, R.: On the asymptotic distribution of eigenvalue branches of the Schrödinger operator $H \pm \lambda W$ in a spectral gap of H . *J. Reine Angew. Math.* 399, 38-59 (1989) MR **90i**:35199
10. Hörmander, L.: *The analysis of linear partial differential operators.* 3. Berlin, New York, Heidelberg: Springer-Verlag 1985 MR **87d**:35002a
11. Ivrii, V.: Precise spectral asymptotics for elliptic operators. *Lecture Notes in Math.* 1100, 1984 MR **86h**:58139
12. Khryashchev, S.V.: Asymptotics of the discrete spectrum of the perturbed Hill operator. *Zap. LOMI*, 147, 188-189 (1985); English transl. in *J. Soviet Math.* 32 (1986), no. 2. CMP 18:07
13. Khryashchev, S.V.: On the discrete spectrum of the perturbed periodic Schrödinger operator. *Zap. LOMI*, 190, 157-162 (1991); English transl. in *J. Soviet Math.* 71 (1994), no. 1. MR **92h**:35168
14. Levendorskiĭ, S.Z.: The method of approximate spectral projection. *Izv. AN SSSR, Ser. Mat.* 49, 1177-1229 (1985); English transl. in *Math. USSR Izv.* 27 (1986), MR **87c**:35127
15. Levendorskiĭ, S.Z.: Non-classical spectral asymptotics. *Russian Math. Surveys*, 43, No 1, 149-192 (1988) MR **89h**:35249
16. Levendorskiĭ, S.Z.: *Asymptotic distribution of eigenvalues of differential operators.* Dordrecht: Kluwer Academic Publishers. 1990 MR **91m**:35176
17. Raikov, G.D.: Eigenvalue asymptotics for the Schrödinger operator with perturbed periodic potential. *Inv. Math.* 110, 75-93 (1992) MR **93k**:35200
18. Reed, M., Simon, B.: *Methods of modern mathematical physics.* 4. New York, San Francisco, London: Academic Press 1978 MR **58**:12429c
19. Rozenbljum, G.V., Solomyak, M.Z., Shubin, M.A. *Spectral theory of differential operators.* *Itogi Nauki i Tekniki: Sovremennye Problemy Mat.: Fundamental'nye Napravleniya*, vol. 64, VINITI, Moscow, 1989; English transl., *Encyclopaedia of Math. Sci.*, vol 64 (Partial Differential Equations, VII, Springer-Verlag, 1994.
20. Seeley, R.T.: Complex powers of an elliptic operator. *Singular Integrals, Proc. Symp. Pure Math.*, vol. 10, Amer. Math. Soc., 1969, pp. 288-307. MR **38**:6220
21. Sobolev, A.V.: Weyl asymptotics for the discrete spectrum of the perturbed Hill operator. In: *Estimates and asymptotics for discrete spectra of integral and differential equations. Adv. Sov. Math.* vol.7, Providence, RI: AMS, 1991, 159-178 MR **95i**:34158

ROSTOV INSTITUTE OF NATIONAL ECONOMY, ENGELS'A 69, 344798, ROSTOV-ON-DON, RUSSIA
E-mail address: leven@ns.rnd.runnet.ru