

Addendum to: The Infrared Spectrum of CS₂ in the ν_3 Band Region

F. Winther

Institut für Physikalische Chemie, Universität Kiel

Z. Naturforsch. **44a**, 594 (1989);
received April 1, 1989

In a recent paper by Wells et al. [1] the band origins of ν_3 of $^{12}\text{C}^{32}\text{S}_2$ and $^{13}\text{C}^{32}\text{S}_2$ have been determined with high accuracy from heterodyne frequency measurements. As mentioned in our paper on these bands [2] our spectra were calibrated with water vapour lines from [3]. This limited the accuracy of all band centers in the paper to $\pm 3 \cdot 10^{-4} \text{ cm}^{-1}$ (one standard error) – not allowing for any systematic error of the calibration lines, however.

Reprint requests to Dr. F. Winther, Institut für Physikalische Chemie der Christian-Albrechts-Universität zu Kiel, Ludwig-Meyn-Str. 8, D-2300 Kiel.

In a paper by Brown and Toth [4] it has been indicated that the frequencies of these calibration lines are $3.63 \cdot 10^{-4} \text{ cm}^{-1}$ too high (mean value for 71 lines). It should be noted that a corresponding correction has not been applied to the same frequencies in "Handbook of Infrared Standards" [5].

Indeed, a comparison of the band origins of [1] and [2] shows that our frequencies are 2.9(8) and $5.2(8) \cdot 10^{-4} \text{ cm}^{-1}$ too high for $^{12}\text{C}^{32}\text{S}_2$ and $^{13}\text{C}^{32}\text{S}_2$, respectively. This confirms the statement of [4] above about the water line frequencies.

In our spectra $^{13}\text{C}^{32}\text{S}_2$ was recorded in natural abundance so that the differences from the heterodyne measurements refer to the strongest and one of the weakest line series. There may have been a slight asymmetry in our line profiles (which we did not check), which would account for the incomplete numerical agreement for the two bands. However, a mean correction of $4 \cdot 10^{-4} \text{ cm}^{-1}$ should be subtracted from all band center frequencies given in our paper [2]. Their absolute accuracies should then all be about 10^{-4} cm^{-1} .

- [1] J. S. Wells, M. Schneider, and A. G. Maki, J. Mol. Spectr. **132**, 422 (1988).
- [2] F. Winther, U. Heyne, and A. Guarnieri, Z. Naturforsch. **43a**, 215 (1988).
- [3] G. Guelachvili, J. Opt. Soc. Amer. **73**, 137 (1983).

- [4] L. R. Brown and R. A. Toth, J. Opt. Soc. Amer. **B2**, 842 (1985).
- [5] G. Guelachvili and K. N. Rao, Handbook of Infrared Standards, Academic Press, NY 1986

0932-0784 / 89 / 0600-0594 \$ 01.30/0. – Please order a reprint rather than making your own copy.



Dieses Werk wurde im Jahr 2013 vom Verlag Zeitschrift für Naturforschung in Zusammenarbeit mit der Max-Planck-Gesellschaft zur Förderung der Wissenschaften e.V. digitalisiert und unter folgender Lizenz veröffentlicht: Creative Commons Namensnennung-Keine Bearbeitung 3.0 Deutschland Lizenz.

Zum 01.01.2015 ist eine Anpassung der Lizenzbedingungen (Entfall der Creative Commons Lizenzbedingung „Keine Bearbeitung“) beabsichtigt, um eine Nachnutzung auch im Rahmen zukünftiger wissenschaftlicher Nutzungsformen zu ermöglichen.

This work has been digitalized and published in 2013 by Verlag Zeitschrift für Naturforschung in cooperation with the Max Planck Society for the Advancement of Science under a Creative Commons Attribution-NoDerivs 3.0 Germany License.

On 01.01.2015 it is planned to change the License Conditions (the removal of the Creative Commons License condition "no derivative works"). This is to allow reuse in the area of future scientific usage.