

# Light-Harvesting Complexes of Aerobic Bacteriochlorophyll-Containing Bacteria *Roseococcus thiosulfatophilus*, RB3 and *Erythromicrobium ramosum*, E5 and the Transfer of Excitation Energy from Carotenoids to Bacteriochlorophyll

Vladimir Yurkov<sup>a,\*</sup>, Nasser Gad'on<sup>a</sup>, Alexander Angerhofer<sup>b</sup> and Gerhart Drews<sup>a</sup>

<sup>a</sup> Institut für Biologie II, Mikrobiologie, Albert-Ludwigs-Universität Freiburg, Schänzlestraße 1, D-79104 Freiburg i. Br., Bundesrepublik Deutschland

<sup>b</sup> Physikalisches Institut, Universität Stuttgart, Pfaffenwaldring 57, D-70550 Stuttgart, Bundesrepublik Deutschland

Z. Naturforsch. **49c**, 579–586 (1994); received June 6, 1994

Antenna Complexes, Carotenoids, Energy Transfer, Obligate Aerobic Photosynthetic Bacteria

The spectral analysis of membrane fractions and a fraction enriched in light-harvesting (LH) complex I + reaction center (RC) of *Roseococcus thiosulfatophilus*, RB3 and of the membrane fraction and of the isolated LH II and LH I + RC complexes of *Erythromicrobium ramosum*, E5 is reported. Quantum yields of singlet energy transfer between carotenoids and bacteriochlorophyll (light-harvesting function) were calculated from comparison of absorption and fluorescence excitation spectra of the respective preparations. The results indicate that the excess of carotenoids in the membrane does not contribute to the light-harvesting function of the LH complexes.

## Introduction

*Roseococcus thiosulfatophilus*, RB3 (Yurkov and Gorlenko, 1992) and *Erythromicrobium ramosum*, E5 (Yurkov *et al.*, 1991, 1992) are obligate aerobic chemotrophic bacteria which can synthesize a photosynthetic apparatus. The contribution of light energy to the formation of an electrochemical proton gradient across the cytoplasmic membrane in these bacteria is not known, but seems to be low (Harashima *et al.*, 1987; Yurkov and van Gemerden, 1993). The reasons why these bacteria can not grow photosynthetically under anaerobic conditions as the photosynthetic purple bacteria are unknown.

Many strains of these aerobic, bacteriochlorophyll (BChl)-containing bacteria belong taxonomically to the  $\alpha$ -subclass of *Proteobacteria* (Woese, 1987; Harashima *et al.*, 1988; Shiba *et al.*, 1991; Fuerst *et al.*, 1993; Yurkov *et al.*, 1994). Recently it was shown that the DNA sequence of the *puf* operon, encoding the genes for the proteins of

the light-harvesting complex I (LH I), the reaction center (RC) and the tetraheme subunit of *Roseobacter denitrificans*, is very similar to that of *Rhodobacter* species (Liebetanz *et al.*, 1991).

The organization of photosynthetic RC seems to be similar to that of purple bacteria. Two types of LH antenna systems are present (Harashima *et al.*, 1988). They were named LH I (B870) and LH II (B800–850) according to the terminology used for purple bacteria although some of them have slightly different absorption bands.

In a previous article we have shown that cells of *R. thiosulfatophilus* and *E. ramosum* contain a large number of unusual and mostly polar carotenoids (Crt) (Yurkov *et al.*, 1993).

The molar ratio of Crt to BChl was found to be 9:1 in whole cells but in LH complexes 0.1 to 1:1. Bacteriorubixanthinal was the major Crt in the pigment proteins of the LH and RC complexes of *E. ramosum*. In addition small amounts of spirilloxanthin (LH I + RC) and zeaxanthin (LH II) were present. In *R. thiosulfatophilus* C<sub>30</sub>-carotene-dioate(4,4'-diapocarotene-4,4'-dioate) (compound A) and the respective diglucosyl ester (compound B) are the major carotene compounds. The enriched LH I + RC fraction contains mainly compound B (di-( $\beta$ -D-glucopyranosyl)-4,4'-diapocar-

\* Present address: Centre de Cadarache, SBCIDPVE, F-13108 St. Paul lez Durance, France.

Reprint requests to Prof. Dr. G. Drews.  
Telefax: (49) 761-2032626.



tene-4,4'-dioate). The Crt's which are not bound to the BChl-protein complexes, were found in the cell envelope fraction (cytoplasmic membrane and outer membrane) (Yurkov *et al.*, 1993). The function of these carotenoids is unknown. They may play a role in scavenging free radicals and/or singlet oxygen, processes which have been observed for several Crt's in organic solvents (Krinsky, 1979, 1989; Oliveros *et al.*, 1992) or they screen the cells from high light intensities.

## Materials and Methods

### Strains, cultivation and membrane isolation

*Roseococcus thiosulfatophilus* strain, RB3, DSM N8511, and *Erythromicrobium ramosum*, E5, DSM N8510, were cultivated at 30 °C semi-aerobically in the dark in order to maximize the formation of the photosynthetic apparatus. The bacteria were grown on a yeast extract-peptone-acetate medium (Yurkov *et al.*, 1993) in a 12 l New Brunswick fermenter (stirring at 200 rpm) for 48 h. Cell-harvesting, disruption in a French pressure cell, and fractionation of cell membranes has been described previously (Yurkov *et al.*, 1993).

### Isolation of pigment-protein complexes

Membrane fractions I and II of *R. thiosulfatophilus* and fraction I of *E. ramosum*, isolated by sucrose density gradient centrifugation (Yurkov *et al.*, 1993), were used for spectroscopical studies and for isolation of pigment-protein complexes. These membrane fractions (about 17 µg BChl/ml) were resuspended in 20 mM Tris-HCl (pH 7.8) buffer and mixed (1:1) with 1.2% of lauryldimethylamine N-oxide (LDAO). The suspension was incubated at room temperature in the dark under gentle stirring. After 30 min the mixture was layered on a sucrose step gradient (0.3, 0.6 and 1.2 M sucrose in Tris-HCl buffer of 20 mM, pH 7.8) containing 0.05% LDAO. After centrifugation for 16 h at 90,000×g in a Beckman Ti60 rotor the pigmented protein bands were collected.

For further purification the LH II- and LH I-enriched fractions were directly applied to a DEAE-Sepharose column equilibrated with 20 mM NaPO<sub>4</sub> buffer (pH 8.0).

The complexes were purified by stepwise elution with increasing concentrations of NaCl in

20 mM NaPO<sub>4</sub> buffer (pH 8.0), containing 0.05% LDAO.

### Analytical methods

BChl and Crt were extracted from whole cells, membrane fractions and pigment-protein complexes with acetone:MeOH 7:2 (v/v). The total amount of BChl was determined according to Clayton (1966). The protein content was measured by the method of Lowry *et al.* (1951). Membrane proteins were separated by SDS-PAGE on an 11.5–16.5% linear gradient of acrylamide. Proteins were stained with Coomassie Brilliant Blue, G-250.

### Spectroscopic methods

Absorption spectra at room temperatures were recorded with a Kontron spectrophotometer UIVICON 860, and plotted by using the computer program "Diagramme" (Atari). Absorption, fluorescence and fluorescence excitation spectra at low temperatures were recorded using a home-built spectrometer, described in detail in Kaiser *et al.* (1981), Kaiser (1982) and Angerhofer *et al.* (1986).

The quantum yield of energy transfer from Crt's to BChl was calculated from the intensities of their respective absorption bands in the absorption and fluorescence excitation spectra according to the following formula (Goedheer, 1959):

$$\eta = \frac{I_{\text{Crt}}^{\text{exc}} \times I_{\text{BChl}}^{\text{abs}}}{I_{\text{Crt}}^{\text{abs}} \times I_{\text{BChl}}^{\text{exc}}}.$$

$\eta$  is the quantum yield of the energy transfer from the Crt excitation band ( $I^{\text{exc}}$  is the intensity of the excited Crt absorption band in the excitation spectrum,  $I^{\text{abs}}$  – the intensity of the same band as found in the absorption spectrum) to the fluorescing first excited singlet state of the BChl ( $I_{\text{BChl}}^{\text{exc,abs}}$  – are the intensities of the  $\text{S}_0 \Rightarrow \text{S}_1$  fluorescence excitation and absorption bands of BChl). The intensities may be taken just from the peak values of the spectra since differences in linewidth between corresponding peaks in the absorption and fluorescence excitation spectra were not observed. Additionally, we applied a multi-Gaussian fit to the spectra and extracted the area under the respective excitation bands for the calculation of the quantum yields. Both methods gave compa-

rable results to within  $\pm 5\%$  in all cases. In the following the average values gained by both methods will be given. Calculation of band intensities was done with spectra corrected for the instrument sensitivities and plotted on a linear photon energy scale (*i.e.* in  $\text{cm}^{-1}$ ).

## Results

### *Roseococcus thiosulfatophilus*, RB 3, pigment-protein complexes

From detergent-treated membranes of *R. thiosulfatophilus* a BChl-protein complex was isolated (see Methods) and eluted from DEAE-Sepharose columns at 350 mm NaCl. The complex showed a

main absorption band at 856 nm (at room temperature; Fig. 1 a). Since the absorption spectrum of this complex in the near-infrared (NIR) region was almost identical to that of the membrane (Fig. 1 in Yurkov *et al.*, 1993) and little BChl was lost during the isolation procedure, most if not all of the BChl molecules in strain RB 3 seems to be organized in this complex.

The NIR absorption maximum of the membrane fraction appearing at 864 nm at 6 K (see Fig. 3a, b-II) is unusual for purple bacteria because LH I complexes normally show maxima between 870 and 890 nm. The ratio of Crt to BChl absorption is relatively high (see Fig. 1b in Yurkov *et al.*, 1993, and Fig. 3a, b-II). The peaks at 484 nm, 514 nm, and 557 nm are tentatively assigned to




Fig. 1. Absorption spectra of isolated pigment-protein complexes recorded at room temperature. a) LH I + RC complex isolated from *Roseococcus thiosulfatophilus*; b) LH I + RC complex isolated from *Erythromicrobium ramosum*; c) LH II complex isolated from *E. ramosum*.




Fig. 2. Protein patterns of membranes or pigment-protein complexes separated by SDS polyacrylamide gel electrophoresis. a) Isolated from *Roseococcus thiosulfatophilus*, lane A: protein markers in kDa; lane B: membrane fraction from *Rhodobacter capsulatus* 37b4, from top to the bottom LH II $\gamma$ , LH I $\alpha$ , LH II $\alpha$ , LH II $\beta$  and LH I $\beta$ , double band; lane C: membrane fraction I of *R. thiosulfatophilus*; lane D: membrane fraction II of *R. thiosulfatophilus*; lane E: LH I-enriched fraction from membranes of *R. thiosulfatophilus*. b) *Erythromicrobium ramosum*, lane A: membrane fraction; lane B: enriched LH I complex; lane C: LH II complex; lane D: molecular weight markers in kDa.

compound B (see Introduction). They appeared with lower intensity in fraction II of the membrane (Fig. 3b) compared with fraction I (Fig. 3a). Fraction II showed a lower Crt/BChl ratio than fraction I (10:1).

The LH I + RC complex showed the same Crt absorption maxima as the membrane but a lower Crt/BChl ratio of about 1.4:1.0 (Fig. 3c-II), with approximately three times less Crt's per BChl (mol: mol) than membrane fraction I.

The polypeptide patterns of the isolated LH I complex was analyzed by SDS-PAGE (Fig. 2a). Four polypeptides with an apparent  $M_r$  of about 8000 (two upper) and 7000 (two lower) were observed. These polypeptides were also present in membrane fractions of *R. thiosulfatophilus* (Fig. 2a). The complex is named LH I.

#### *Fluorescence and quantum yield of Car $\Rightarrow$ BChl singlet energy transfer*

The quantum yield determination of excitation energy transfer are based on the total Crt's present in the respective fractions since we could not obtain highly purified RC and LH I preparations. The fluorescence emission spectrum of membrane fraction I showed one peak at 876 nm, originating from the LH I emission (see Fig. 3a-I). The main peak of the absorption and fluorescence excitation spectra (Fig. 3a-III) was at 864 nm. In the visible region the relative intensities differ considerably. The strong Crt absorption bands at 484, 514 and 557 nm (Fig. 3a-II) contributed only weakly to the fluorescence excitation spectrum 499, 511 nm; Fig. 3a-III). These data support the idea that only a subpopulation from the bulk of the Crt's in the membranes is efficient in energy transfer to LH I. The quantum yield calculated for the total Crt content was 7.7%. If we assume a Crt/BChl ratio of 1:1 in the LH I complex and neglect energy transfer from the membrane-bound secondary Crt's (Crt/BChl ratio 9:1) for membrane fraction I), one can estimate the quantum yield of Crt  $\Rightarrow$  BChl energy transfer in the native LH I complex to 77%, well in the range of values observed in purple photosynthetic bacteria (Noguchi *et al.*, 1990).

The spectral location of the fluorescence emission and excitation and absorption bands in fraction II are the same as in fraction I (Fig. 3b), but




Fig. 3. Absorption (II), fluorescence emission (I) and fluorescence excitation (III) spectra of fractions isolated from *Roseococcus thiosulfatophilus* recorded at 6 K. a) Membrane fraction I, b) membrane fraction II, c) the LH + RC-enriched fraction. Absorption and fluorescence excitation spectra are displayed normalized to equal height of their NIR band. Fluorescence emission and fluorescence excitation spectra were corrected to the instrumental response function.

the Crt absorption bands are weaker than in fraction I. A quantum yield for energy transfer of 11.2% was derived. This, however, reflects a lower concentration of bulk carotenoids (inactive in energy transfer to BChl) in this membrane preparation, rather than a higher energy transfer efficiency for the protein-bound carotenoids.

The isolated LH I + RC complex of *R. thiosulfatophilus* displayed the same fluorescence emission peak at 876 nm as in membrane fractions I and II (Fig. 3c-I). The fluorescence excitation and absorption spectra have a NIR band at 864 nm. At 594 nm the Qx band of the BChl is visible in the excitation spectrum. The Crt bands appeared in the excitation spectrum at 495, 528 and 569 nm which are somewhat red-shifted compared to the

peak maxima in the absorption spectrum. From the intensities a quantum efficiency for energy transfer of 29% was calculated. The fact that the Crt excitation bands were not found at the same spectral positions in both spectra may indicate that RC and LH complexes have different carotenoids. The quantum efficiency calculated that way was then obscured by the RC Crt's which do not contribute to the LH fluorescence.

#### *Erythromicrobium ramosum, E5, pigment-protein complexes*

From membrane fraction I of *E. ramosum* (Yurkov *et al.*, 1993) two subfractions were obtained after treatment with LDAO and sucrose density gradient centrifugation. The upper band was enriched in LH II and the lower band in LH I + RC. Both fractions were abundant in Crt's. The bands were separately applied to DEAE-Sephadex chromatography. The LH I + RC complex was eluted at 225 mM NaCl, the LH II at 100 mM NaCl. At room temperature the purified LH I + RC complex showed a main absorption band at 871 nm and a smaller one at 801 nm (Fig. 1b). LH II absorbed at 798 nm and 832 nm (Fig. 1c). At 6 K absorption maxima at 440 nm, 467 nm and 497 nm in the Crt absorption region, and at 589 nm, 797 nm, 838 nm and 878 nm, belonging to BChl were observed in the membrane fractions, indicating the presence of two LH complexes (Fig. 4a).

The isolated LH II complex (Fig. 4b) showed BChl maxima at 796 and 840 nm (BChl) and Crt maxima at 503 nm and 535 nm indicating a change in the Crt composition compared with the whole membranes. For LH I + RC fraction showed a LH I maximum at 884 nm (Fig. 4c) and a very weak Crt band around 500 nm.

Both LH II and LH I + RC fractions contained bacteriorubixanthinal as the main Crt compound and additionally zeaxanthin in the LH II fraction and spirilloxanthin in the LH I + RC fraction (Yurkov *et al.*, 1993).

The LH I complex contained two polypeptides with an apparent molecular weight of about 8000 and 6000 and the LH II preparation three polypeptides of about 16,000, 9000 and 8000  $M_r$ . The RC bands have not been identified.




Fig. 4. Absorption (II), fluorescence emission (I) and fluorescence excitation (III) spectra of fraction isolated from *Erythromicrobium ramosum* recorded at 6 K. a) Membrane fraction; b) light-harvesting complex II (LH II); c) LH I + RC fraction. Absorption and fluorescence excitation spectra are displayed normalized to equal height of their NIR band. The respective fluorescence spectra appear normalized to the height of this band. Fluorescence and fluorescence excitation spectra were corrected to the instrumental response function.

#### *Fluorescence and quantum yield of Car $\Rightarrow$ BChl singlet energy transfer*

The fluorescence emission bands of the membrane fraction of *E. ramosum* (Fig. 4a-I) at 896 nm and a weaker one at 859 nm, at 6 K, are indicative of LH I and LH II fluorescence, respectively. The fluorescence excitation spectrum taken at a detection wavelength of 897 nm showed the same spectral features as the absorption spectrum. Both spectra were normalized to the same intensity of the NIR band at 878 nm of the LH I complex. The two bands at 797 nm and 838 nm belong to the LH II complex. The intensity of these bands was about 30% lower than in the corresponding absorption spectrum, indicating that energy from

LH II to LH I was transferred at a low rate. This also explains why a LH II fluorescence emission band at 859 nm was observed. In the visible region the BChl Qx band was visible at 589 nm and the Crt absorption bands at 440, 467, 497 nm and a shoulder at 532 nm.

For analysis of the quantum yield of energy transfer in the membrane fraction of *E. ramosum* the 538 nm and the 497 nm bands of the Crt's were selected which gave yields of 25.5% and 33.0%, respectively. This difference may be explained by differences in Crt pigment composition of the LH I and LH II complexes and the Crt pool not associated with the photosystem. It also indicated that the Crt's not associated with the LH proteins are inactive in energy transfer to the BChl bound to these proteins.

The fluorescence emission spectrum of the LH II complex showed only one band with its maximum at 867 nm (Fig. 4b). The absorption and fluorescence excitation spectra revealed peaks at 503, 535, 586, 796, and 840 nm. The quantum yield of Crt to BChl energy transfer was calculated to be 94%. The fluorescence emission spectrum of the LH I + RC preparation (Fig. 4c) was characterized by one main peak at 915 nm, and two weak peaks at 861 nm and 790 nm. The main band was ascribed to the LH I emission. The weaker maxima at shorter wavelengths were probably due to residual LH II emission and free BChl. The fluorescence excitation and absorption spectra were again very similar with respect to the position of their peaks. They differ in intensity only in those bands which must be ascribed to the RC. The main band in the NIR appeared at 884 nm with a small shoulder at 838 nm, probably due to the residual LH II complex. The bands at 763 and 796 nm, belonging to the RC, were weaker by at least 50% in the fluorescence excitation spectrum. These results indicate some energy transfer from RC's to the LH I pool. In the visible region the BChl Qx band at 590 nm and weak Crt bands at 498 nm and 533 nm were observed. From their intensities a quantum yield of 60% was calculated.

## Discussion

The species described in this article are not closely related to *Rhodobacter capsulatus* or *Rhodobacter sphaeroides* (Yurkov *et al.*, 1994). The LH

complexes of these bacteria differ from those of many purple bacteria. The major NIR absorption band of the LH I + RC-enriched fraction of *R. thiosulfatophilus* was found at 856 nm (room temperature; 864 nm at 6 K). This and the polypeptide pattern (Fig. 2a) indicate an uncommon protein environment of BChl.

Unusual spectral properties were also observed for the LH II complex of *E. ramosum*. Besides the 798 nm peak the major NIR peak was at 832 nm (room temperature) and 840 nm (at 6 K), respectively. It has to be studied whether *E. ramosum* is able to express variable LH II complexes dependent from temperature or light intensity as *Rhodopseudomonas acidophila* and *R. palustris* (Bissig *et al.*, 1990; Evans *et al.*, 1990; Gardiner *et al.*, 1992; Tadros *et al.*, 1993). The isolated LH II complex contained three proteins like *R. capsulatus*.

In previous studies a RC-B865 complex from *Erythrobacter longus*, OCh101, and RC-B870 and B806 complexes from *Roseobacter denitrificans*, OCh114 were isolated and described (Shimada *et al.*, 1985). The isolation of new types of LH complexes, described in this article, underlines that these aerobic bacteria are heterogenous in their physiology, taxonomic position and composition of the photosynthetic apparatus.

The emission and excitation spectra of the membrane fractions of both *R. thiosulfatophilus* and *E. ramosum* demonstrate that most of the membrane-bound Crt's are not associated with the LH or RC complexes, and that these separately localized Crt's do not transfer excitation energy to the RC. The quantum yields for excitation energy transfer are thus only lower estimates to the true quantum yields in the LH complexes. This is also true for the isolated LH I + RC complexes of *R. thiosulfatophilus* and *E. ramosus* since the Crt's of the RC do not transfer their excitation energy to the LH I complex. The quantum yields of 29% and 60%, respectively, are thus only lower bounds of the true quantum yields in these LH I complexes.

The quantum yield of energy transfer was more exactly determined for the LH II complex of *E. ramosum*. A quantum yield of 94% is comparable to what has been observed for the B800–850 complex of the purple bacterium *Rhodobacter sphaeroides* strain 2.4.1 (Cogdell *et al.*, 1981; van Grondelle *et al.*, 1982; Hayashi *et al.*, 1987; Nogu-

chi *et al.*, 1990), but higher than the yields observed for the LH II complexes of other purple bacteria (Angerhofer *et al.*, 1986; Noguchi *et al.*, 1990; Cogdell *et al.*, 1992).

High concentrations of Crt's not active in photosynthetic energy transfer have been observed in membranes of *Erythrobacter longus*, OCh 101 (Noguchi *et al.*, 1992). The main Crt was (3S, 2'R, 3'R)-3,2',3'-trihydroxy- $\beta,\beta$ -carotene-4-one-3-sulphate (erythroxanthin sulphate), a highly polar molecule (Takaichi *et al.*, 1991). It was suggested that its possible function is photoprotection by quenching harmful singlet oxygen (Noguchi *et al.*, 1992). High concentrations of "extrinsic" carotenoids have also been observed in algae. It has recently been suggested that high extrachloroplastic concen-

trations of astaxanthin ester-containing lipid globules function as a sunscreen to minimize photodamage during periods of exposure to intense solar radiation of snow algae (e.g. *Chlamydomonas nivalis*; Bidigara *et al.*, 1993). In addition to the "sunscreen" effect these secondary Crt's may act as inhibitors of photodynamically induced damage as indicated by a recent study of Hagen *et al.* (1993) on *Haematococcus lacustris*.

### Acknowledgements

The work was supported by grants of the Deutsche Forschungsgemeinschaft and by Fonds der Chemischen Industrie (G. D.). V. Y. was supported by an Alexander von Humboldt fellowship.

Angerhofer A., Cogdell R. J. and Hipkins M. F. (1986), A spectral characterization of the light-harvesting pigment-protein complexes from *Rhodopseudomonas acidophila*. *Biochim. Biophys. Acta* **848**, 333–341.

Bidigare R. R., Ondrussek M. E., Kennicutt M. C., Iturriaga R., Harvey H. R., Honam R. W. and Macko S. A. (1993), Evidence for a photoprotective function for secondary carotenoids of snow algae. *J. Phycol.* **29**, 427–434.

Bissig I., Wagner-Huber R. V., Brunisholz B. A. and Zuber H. (1990), Multiple antenna complexes in various purple photosynthetic bacteria, in: *Molecular Biology of Membrane-Bound Complexes in Phototrophic Bacteria* (G. Drews and E. A. Dawes, eds.). Plenum Press, New York, pp. 199–210.

Clayton R. K. (1966), Spectroscopic analysis of bacteriochlorophyll *in vivo* and *in vitro*. *Photochem. Photobiol.* **5**, 669–677.

Cogdell R. J., Hipkins M. F., MacDonald W. and Truscott T. G. (1981), Energy transfer between the carotenoid and bacteriochlorophyll within the B800–850 light-harvesting pigment-protein complex of *Rhodobacter sphaeroides*. *Biochim. Biophys. Acta* **634**, 191–202.

Cogdell R. J., Anderson P. O. and Gillbro T. (1992), Carotenoid singlet states and their involvement in photosynthetic light-harvesting pigments. *J. Photochem. Photobiol. B: Biol.* **15**, 105–112.

Evans W. R., Fleischman D. E., Cavert H. E., Pyati P. V., Alter G. M. and Rao N. S. S. (1990), Bacteriochlorophyll and photosynthetic reaction centers in *Rhizobium* strain BTA:1. *Appl. Environm. Microbiol.* **56**, 3445–3449.

Fuerst J. A., Hawkins J. A., Holmes A., Sly L. I., Moore C. J. and Stackebrandt E. (1993), *Porphyrobacter neustonensis* gen. nov., spec. nov., an aerobic bacteriochlorophyll-synthesizing budding bacterium from fresh water. *Int. J. System. Bacteriol.* **43**, 125–134.

Gardiner A. T., MacKenzie R. C., Barrett S. J., Kaiser K. and Cogdell R. J. (1992), The genes for the peripheral antenna complex apoproteins from *Rhodopseudomonas acidophila* 7050 form a multigene family, in: *Research in Photosynthesis, Vol. I* (N. Murata, ed.). Kluwer Academic Publishers, Dordrecht, pp. 77–80.

Goedheer J. C. (1959), Energy transfer between carotenoids and bacteriochlorophyll in chromatophores of purple bacteria. *Biochim. Biophys. Acta* **35**, 1–8.

Hagen C., Braune W. and Greulich F. (1993), Functional aspects of secondary carotenoids in *Haematococcus lacustris*, IV. Protection from photodynamic damage. *J. Photochem. Photobiol. B: Biol.* **20**, 153–160.

Harashima H., Kawazoe K., Yoshida I. and Kamata H. (1987), Light-stimulated aerobic growth of *Erythrobacter* spec. OCh 114. *Plant Cell Physiol.* **28**, 365–374.

Harashima K., Shiba T. and Murata N. (eds.) (1988), *Aerobic Photosynthetic Bacteria*. Japan Sci. Soc. Press, Tokyo; Springer Verlag, Heidelberg.

Hayashi H., Iwata K., Noguchi T. and Tasumi M. (1987), Correlation between the efficiency of energy transfer and the polyene chain structure of carotenoids in purple photosynthetic bacteria, in: *Progr. Photosynthesis Res., Vol. I* (J. Biggins, ed.). Martinus Nijhoff Publ., Dordrecht, pp. 33–36.

Kaiser G. H. (1982), Emissions- und Anregungsspektren am photosynthetisierenden Bakterium *Rhodopseudomonas sphaeroides*. Ph.D. thesis, Stuttgart.

Kaiser G. H., Beck J., von Schütz J. U. and Wolf H. C. (1981), Low temperature excitation and emission spectroscopy of the photosynthetic bacteria *Rhodopseudomonas sphaeroides* wild type strain ATCC 17023. *Biochim. Biophys. Acta* **634**, 153–164.

Krinsky N. I. (1979), Carotenoid protection against oxidation. *Pure Appl. Chem.* **51**, 649–660.

Krinsky N. I. (1989), Antioxidant functions of carotenoids. *Free Radical Biol. Med.* **7**, 617–635.

Liebetanz R., Hornberger U. and Drews G. (1991), Organization of the genes coding for the reaction centre L and M subunits and B870 antenna polypeptides  $\alpha$  and  $\beta$  from the aerobic photosynthetic bacterium *Erythrobacter* spec. OCh 114. *Mol. Microbiol.* **5**, 1459–1468.

Lowry O. H., Rosebrough U., Farr A. L. and Randal R. J. (1951), A protein measurement with the Folin phenol reagent. *J. Biol. Chem.* **193**, 265–275.

Noguchi T., Hayashi H. and Tasumi M. (1990), Factors controlling the efficiency of energy transfer from carotenoids to bacteriochlorophyll in purple photosynthetic bacteria. *Biochim. Biophys. Acta* **1017**, 280–290.

Noguchi T., Hayashi H., Shimada K., Takaichi S. and Tasumi M. (1992), *In vivo* states and functions of carotenoids in an aerobic photosynthetic bacterium *Erythrobacter longus*. *Photosynth. Res.* **31**, 21–30.

Oliveros E., Murasecco-Suardi P., Braun A. M. and Hansen H. J. (1992), Efficiency of singlet oxygen quenching by carotenoids measured by near-infrared steady-state luminescence, in: *Methods in Enzymology* **Vol. 213**, (L. Packer, ed.), pp. 420–429.

Shiba T., Shioi Y., Takamiya K., Sutton D. and Wilkinson C. (1991), Distribution and physiology of aerobic bacteria containing bacteriochlorophyll *a* on the east and west coast of Australia. *Appl. Environ. Microbiol.* **57**, 295–300.

Shimada K., Hayashi H. and Tasumi M. (1985), Bacteriochlorophyll-protein complexes of aerobic bacteria, *Erythrobacter longus* and *Erythrobacter* spec. OCh114. *Arch. Microbiol.* **143**, 244–247.

Tadros M. H., Katsiou E., Hoon M., Yurkova N. and Ramji D. P. (1993), Cloning of a new antenna gene cluster and expression analysis of the antenna gene family of *Rhodopseudomonas palustris*. *Eur. J. Biochem.* **217**, 867–875.

Takaichi S., Furihara K., Ishidsu J. I. and Shimada K. (1991), Carotenoid sulphates from the aerobic photosynthetic bacterium *Erythrobacter longus*. *Phytochemistry* **30**, 3411–3415.

van Grondelle R., Kramer H. J. M. and Rijgersberg C. P. (1982), Energy transfer in the B800–850 carotenoid light-harvesting complex of various mutants of *Rhodopseudomonas sphaeroides* and of *R. capsulata*. *Biochim. Biophys. Acta* **682**, 108–215.

Woese C. R. (1987), Bacterial evolution. *Microbiol. Rev.* **51**, 221–271.

Yurkov V. and Gorlenko V. M. (1992), A new genus of freshwater aerobic, bacteriochlorophyll *a*-containing bacteria, *Roseococcus* gen. nov. *Mikrobiologiya* **60**, 628–632, engl. transl.

Yurkov V., Lysenko A. M. and Gorlenko V. M. (1991), Hybridization analysis of the classification of bacteriochlorophyll *a*-containing freshwater aerobic bacteria. *Mikrobiologiya* **60**, 362–366, engl. transl.

Yurkov V., Gorlenko V. M. and Kompentseva E. I. (1992), A new type of freshwater aerobic orange-colored bacterium *Erythromicrobium* gen. nov. containing bacteriochlorophyll *a*. *Mikrobiologiya* **61**, 169–172, engl. transl.

Yurkov V. and van Gemerden H. (1993), Impact of light/dark regimen on growth rate, biomass formation and bacteriochlorophyll synthesis in *Erythromicrobium hydrolyticum*. *Arch. Microbiol.* **159**, 84–89.

Yurkov V., Gad'on N. and Drews G. (1993), The major part of polar carotenoids of the aerobic bacteria *Roseococcus thiosulfatophilus* RB3 and *Erythromicrobium ramosum* E5 is not bound to the bacteriochlorophyll *a* complexes of the photosynthetic apparatus. *Arch. Microbiol.* **160**, 372–376.

Yurkov V., Stackebrandt E., Holmes A., Fuerst J. A., Hugenholtz P., Golecki J., Gad'on N., Gorlenko V. M., Kompentseva E. I. and Drews G. (1994), Phylogenetic positions of novel aerobic, bacteriochlorophyll *a*-containing bacteria and description of *Roseococcus thiosulfatophilus* gen. nov., spec. nov., *Erythromicrobium ramosum* gen. nov., spec. nov., and *Erythrobacter litoralis* spec. nov. *Int. J. System. Bacteriol.* **44**, 427–434.