

## Three Further Dammarane Type Triterpenes from the Frond Exudate of the Fern *Notholaena rigida*

Fco. Javier Arriaga<sup>a</sup>, Angel Rumbero<sup>b</sup> and Eckhard Wollenweber<sup>c</sup>

<sup>a</sup> Tabacalera S.A., Centro I + D, c/Embajadores 51, E-28012 Madrid, Spain

<sup>b</sup> Departamento de Química Orgánica, Universidad Autónoma de Madrid, Cantoblanco, E-28049 Madrid, Spain

<sup>c</sup> Institut für Botanik der Technischen Hochschule, Schnittspahnstraße 3, D-64287 Darmstadt, Bundesrepublik Deutschland

Z. Naturforsch. **51c**, 750–752 (1996); received July 4, 1996

*Notholaena rigida*, Pteridaceae, Farinose Frond Exudate, Novel Triterpenes, Dammarane Derivatives

From the farinose frond exudate of the fern *Notholaena rigida* two new triterpenes have been identified. They were found to be the C-24 epimers of 3 $\beta$ -acetoxy-24, 25-diO-isopropylidene-12 $\beta$ , 20(S),24,25-tetrahydroxy-dammarane. A third triterpene is identical with the known compound pyxinol, (20S,24R)-20,24-epoxy-dammarane-3 $\beta$ ,12 $\beta$ ,25-triol.

### Introduction

The lipophilic farinose coating on the lower frond surface of the Mexican fern *Notholaena rigida* Dav. is composed of terpenoids and flavonoids. In a first paper we dealt with the identification of flavone aglycones (Scheele *et al.*, 1987). Later we reported several triterpenes, including two epoxy-dammarane derivatives (Arriaga-Giner *et al.*, 1991; Arriaga-Giner *et al.*, 1992), an isopropylidene-cyclolanostanol (Arriaga-Giner *et al.*, 1991), and two epimeric isopropylidene-dammarane derivatives (Arriaga *et al.*, 1996). From remaining fractions we have now isolated three further triterpenes which were identified by detailed spectroscopic studies.

### Material and Methods

Dry fronds of *Notholaena rigida* were collected near Cd. Victoria in Edo. Tamaulipas, México in

May, 1983. Collection data, isolation and analysis procedures have been reported previously (Arriaga-Giner *et al.*, 1991). As in the preceding paper (Arriaga *et al.*, 1996), one of those fractions was subjected to "flash" chromatography on Si-gel using  $\text{CH}_2\text{Cl}_2$ -MeOH 15:1 as eluent. Mass spectra were measured on a VG Autospec at 70 eV *via* solid probe. NMR spectra were recorded on a Bruker AC-300 spectrometer at 300 MHz (for  $^1\text{H}$ ) and 75.4 MHz (for  $^{13}\text{C}$ ) in  $\text{CDCl}_3$ . Multiplicities were assigned through DEPT experiments. Mps are uncorrected.

Three pure products were obtained, which all appeared as colourless oils. They exhibit the following spectral properties:

Compound **1A** ( $R_f$  0.64): EI-MS  $m/z$  (rel. int.): 576 ( $M^+$ , -), 561 ( $M^+$ -Me, 0.3), 543 ( $M^+$ -Me- $\text{H}_2\text{O}$ , 2), 525 ( $M^+$ -Me- $2\text{H}_2\text{O}$ , 1) 500 (1), 482 (6), 415 (5), 413 (5), 341 (5), 249 (5), 201 (10), 191 (10), 189 (22), 161 (11), 147 (21), 143 (44), 129 (100), 121 (23), 107 (23), 95 (23), 81 (30), 71 (39) and 59 (85).  $^1\text{H}$  NMR  $\delta$  ppm (J, Hz): 0.85, 0.85, 0.85, 0.90, 0.98, 1.11, 1.22, 1.24, 1.33, 1.43, 2.04 (3H each, all s), 3.58 (H-12, ddd, 10.1, 10.1, 4.9), 3.72 (H-24, brd, 8.6) and 4.49 (H-3, dd, 11.8, 7.4).  $^{13}\text{C}$  NMR: see Table I.

Compound **1B** ( $R_f$  0.60): EI-MS  $m/z$  (rel. int.): 576 ( $M^+$ , -), 561 ( $M^+$ -Me, 0.5), 543 ( $M^+$ -Me- $\text{H}_2\text{O}$ , 2), 525 ( $M^+$ -Me- $2\text{H}_2\text{O}$ , 0.5) 500 (1), 482 (5), 415 (5), 413 (5), 341 (6), 249 (5), 201 (12), 191 (14), 189 (22), 161 (13), 147 (22), 143 (64), 129 (100), 121 (28), 107 (28), 95 (29), 81 (36), 71 (46) and 59 (91).  $^1\text{H}$  NMR  $\delta$  ppm (J, Hz): 0.85, 0.85, 0.85, 0.91, 1.00, 1.11, 1.19, 1.24, 1.34, 1.42, 2.04 (3H each, all s), 3.63 (H-12, m), 3.68 (H-24, m) and 4.45 (H-3, dd, 11.8, 7.4).  $^{13}\text{C}$  NMR: see Table I.

Compound **2** ( $R_f$  0.52): EI-MS  $m/z$  (rel. int.): 476 ( $M^+$ , -), 461 ( $M^+$ -Me, 2), 443 ( $M^+$ -Me- $\text{H}_2\text{O}$ , 3), 425 ( $M^+$ -Me- $2\text{H}_2\text{O}$ , 2), 400 (31), 381 (12), 357 (6), 207 (22), 191 (16), 143 (100), 125 (30), 107 (27), 85 (46) and 59 (28).  $^1\text{H}$  NMR  $\delta$  ppm (J, Hz): 0.76, 0.84, 0.89, 0.96, 0.97, 1.09, 1.26, 1.27 (3H each, all s), 3.18 (H-3, dd, 10.5, 5.2), 3.50 (H-12, ddd, 10.4, 10.4, 4.5) and 3.84 (H-24, dd, 8.3, 6.5).  $^{13}\text{C}$  NMR: see Table I.

Reprint requests to E. Wollenweber. Telefax # 06151/166878.

0939–5075/96/0900–0750 \$ 06.00 © 1996 Verlag der Zeitschrift für Naturforschung. All rights reserved.

D



Dieses Werk wurde im Jahr 2013 vom Verlag Zeitschrift für Naturforschung in Zusammenarbeit mit der Max-Planck-Gesellschaft zur Förderung der Wissenschaften e.V. digitalisiert und unter folgender Lizenz veröffentlicht: Creative Commons Namensnennung-Keine Bearbeitung 3.0 Deutschland Lizenz.

Zum 01.01.2015 ist eine Anpassung der Lizenzbedingungen (Entfall der Creative Commons Lizenzbedingung „Keine Bearbeitung“) beabsichtigt, um eine Nachnutzung auch im Rahmen zukünftiger wissenschaftlicher Nutzungsformen zu ermöglichen.

This work has been digitized and published in 2013 by Verlag Zeitschrift für Naturforschung in cooperation with the Max Planck Society for the Advancement of Science under a Creative Commons Attribution-NoDerivs 3.0 Germany License.

On 01.01.2015 it is planned to change the License Conditions (the removal of the Creative Commons License condition "no derivative works"). This is to allow reuse in the area of future scientific usage.

## Results and Discussion

The  $^1\text{H}$  NMR spectrum of compound **1A** exhibits eleven methyl signals (all singlets) between 0.85 and 2.04 ppm. Three sets of one-proton resonances were observed at 3.58, 3.72 and 4.49 ppm which correspond to methine hydrogens adjacent to an oxygen substituent. The one-proton resonance at 4.49 ppm was assigned to H-3 $\alpha$  on the basis of its splitting pattern and followed from the MS data ( $m/z$  249/189) and methyl singlet at 2.04 ppm which suggested a 3 $\beta$ -acetoxy-dammarane skeleton.

The  $^{13}\text{C}$  NMR spectrum of compound **1A** displays 35 carbon atom signals (see Table I), show-

Table I.  $^{13}\text{C}$  NMR Data for the *Notholaena* triterpenoids **1A** and **1B**, and **2** (and its derivative **2a**).

a,b,c,d: Signals are interchangeable within the same column.

Carbon (multiplicity)	<b>1A</b>	<b>1B</b>	<b>2</b>	<b>2a</b>
1 t	38.5	38.6	38.9	38.6
2 t	23.6a	23.6a	27.4	23.9
3 d	80.8	80.7	78.8	80.6
4 s	37.8	37.7	38.9	38.0
5 d	55.9	55.9	55.9	56.1
6 t	18.1	18.1	18.2	18.3
7 t	34.7	34.6	34.8	34.9
8 s	39.8	39.7	39.7	39.9
9 d	49.8	49.9	50.5	50.6
10 s	37.0	37.0	37.1	37.1
11 t	30.5b	30.8b	31.1a	32.2
12 d	70.2	71.0	70.9	71.0
13 d	48.1	47.6	47.9	48.3
14 s	51.3	51.5	52.0	52.1
15 t	30.9b	31.3b	31.3a	31.6
16 t	26.2	26.3	24.9	25.4
17 d	52.9	53.9	49.3	49.4
18 q	16.1c	16.1c	15.2b	16.6a
19 q	16.4c	16.4c	15.3b	16.4a
20 s	73.5	73.7	86.4	86.6
21 q	26.7d	26.9d	26.1	26.9b
22 t	32.9	32.0	32.6	32.7
23 t	23.5a	23.4a	28.5	28.6
24 d	84.4	84.1	85.4	85.5
25 s	80.8	80.4	70.0	70.3
26 q	22.6	22.9	27.5c	27.1b
27 q	25.5	26.1	27.9c	27.6b
28 q	27.9	27.9	27.9c	28.0b
29 q	15.6	15.7	16.3b	15.4a
30 q	16.7	16.8	18.1	18.3
Subst. s	106.8	106.5		
q	28.3	28.6		
q	26.9 d	27.0 d		
$\text{CH}_3-\text{C=O}$ s	170.9	171.0		
$\text{CH}_3-\text{C=O}$ q	21.2	21.2		




Fig. 1. Structural formulae for the *Notholaena* triterpenoids **1A** and **1B**, the 24*R* and 24*S*-epimers of 3 $\beta$ -acetoxy-12 $\beta$ , 20(S)-dihydroxy-24,25-di-O-isopropylidene-dammarane, and triterpenoid **2**, 3 $\beta$ ,12 $\beta$ ,25-trihydroxy-20 (*S*), 24(*R*)-epoxy-dammarane (and its derivative **2a**).

ing five deshielded carbons between 70.2–84.4 ppm, a carbon signal at 106.8 ppm that suggested the existence of an acetonide, and a carbonyl at 170.9 ppm.

The  $^1\text{H}$  NMR spectrum of compound **1B** shows substantially the same signals as compound **1A**, but two one-proton multiplets at 3.63 and 3.68 ppm are partially overlapped. No significant differences in methyl signals are observed.

Comparison of the  $^{13}\text{C}$  NMR spectra of compounds **1A** and **1B** (see Table I) showed no differences with previously isolated triterpene-pentol epimers (Arriaga *et al.*, 1996) other than affecting the neighbourhood of the acetate (carbons 2–4). The chemical shifts for the signals attributed to C-17, C-21 and C-22 suggested the same stereochemistry at C-20 (Asakawa *et al.*, 1977). Thus, both compounds seem to be epimers at C-24.

The mass spectrometric fragmentation pattern of both compounds is also identical. The molecular ion corresponding to the formula of  $\text{C}_{35}\text{H}_{60}\text{O}_6$  is not observed because losses of water and methyl are highly favoured. The structure of these new natural products, **1A** and **1B**, is fully in accordance with the structure of 3 $\beta$ -acetoxy-12 $\beta$ ,20(*S*)-dihydroxy-24,25-di-O-isopropylidene-dammarane, but assignment of the two epimers at C-24 was not possible.

Compound **2** shows a  $^1\text{H}$  NMR spectrum that differs in the methyl number and lack of the ace-

tate. A comparison with related triterpenes isolated from this plant lead us to assume that the side chain is different, the acetonide being absent. Reviewing the literature we conclude that the spectroscopic data are fully in agreement with the structure of pyxinol,  $3\beta,12\beta,25$ -trihydroxy- $20(S),24(R)$ -epoxy-dammarane, previously isolated from the lichen *Pyxine endochrysina* (Yoshioka *et al.*, 1972). Its  $^{13}\text{C}$  NMR spectrum, reported here for the first time, is closely related (see Table I) to that of its 3-acetyl derivative, **2a**, previously used in the structural elucidation of *Panax* saponins (Tanaka and Yahara, 1978). Again, the chemical shifts for C-21 and C-22 at ca. 26 and 33 ppm agree for  $20S$  stereochemistry (21 and 39 ppm, respectively, for  $20R$ ) (Francisco *et al.*, 1984).

Arriaga-Giner F. J., Rumbero A. and Wollenweber E. (1996), Two new triterpenes from the frond exudate of the fern *Notholaena rigida*. *Z. Naturforsch.* **51c**, 423–425.

Arriaga-Giner F. J., Rumbero A. and Wollenweber E. (1992), New triterpenes from the frond exudates of some *Notholaena* species. *Z. Naturforsch.* **47c**, 508–511.

Arriaga-Giner F. J., Rullkötter J., Peakman T. M. and Wollenweber E. (1991), New triterpenes from the frond exudate of the fern *Notholaena rigida*. *Z. Naturforsch.* **46c**, 507–512.

Asakawa J., Kasai R., Yamasaki K. and Tanaka O. (1977),  $^{13}\text{C}$  NMR study of ginseng saponins and their related dammarane type triterpenes. *Tetrahedron* **33**, 1935–1939.

Denisenko V. A., Novikov V. A., Malinovskaya G. V. and Elyakov G. B. (1983), Effects of side-chains at C-17

The pair of epimers **1A/1B** are the 3-O-acetyl derivatives of the parent compounds recently reported from the same plant (Arriaga *et al.*, 1996). Acetonides as natural products are rather unusual, but we have found five derivatives from the dammarane and cycloartane skeletons in this species (Arriaga *et al.*, 1996; Arriaga-Giner *et al.*, 1991). Pyxinol, compound **2**, is reported for the second time only as a natural product. As other dammarane-triterpenoids isolated from this plant, it has the side chain cyclized as tetrahydrofuran ring but the stereochemistry of C-20 differs from the other reported components. An isomer with reverse configuration at C-3 and C-20 has also been isolated as a natural product (Denisenko *et al.*, 1983).

on  $^{13}\text{C}$  chemical shifts of dammarane-type tetracyclic triterpenoids. *Izv. Akad. Nauk. SSR Ser. Khim.* **32**, 2727–2734.

Francisco C. G., Freire R., Hernández R., Salazar J. A. and Suárez E. (1984), Carbon-13 NMR study of (20,24)-epoxydammarane triterpenes. *Org. Mag. Res.* **22**, 34–38.

Scheele C., Wollenweber E. and Arriaga-Giner F. J. (1987), New flavonoids from cheilanthes ferns. *J. Nat. Prod.* **50**, 181–187.

Tanaka O. and Yahara S. (1978), Dammarane saponins of leaves of *Panax pseudo-ginseng* subsp. *himalaicus*. *Phytochemistry* **17**, 1353–1358.

Yoshioka I., Yamauchi H. and Kitagawa I. (1972), Lichen Triterpenoids. V. On the neutral triterpenoids of *Pyxine endochrysina* Nyl. *Chem. Pharm. Bull.* **20**, 502–513.