

# Bioactive Constituents from *Dracocephalum subcapitatum* (O. Kuntze) Lipsky

Soodabeh Saeidnia<sup>a,\*</sup>, Ahmad Reza Gohari<sup>a</sup>, Michiho Ito<sup>b</sup>, Fumiyuki Kiuchi<sup>c</sup>,  
and Gisho Honda<sup>b</sup>

<sup>a</sup> Department of Pharmacognosy, Faculty of Pharmacy, Mazandaran University of  
Medical Sciences, PO Box 48175-861, Sari, Iran. Fax: +98-1 51-3 25 40 60.  
E-mail: soodabehsaeidnia@hotmail.com

<sup>b</sup> Graduate School of Pharmaceutical Sciences, Kyoto University, Kyoto 606-8501, Japan

<sup>c</sup> Tsukuba Medicinal Plant Research Station, National Institute of Health Sciences,  
Ministry of Health, Labour and Welfare, Tsukuba, Ibaraki, 305-0843, Japan

\* Author for correspondence and reprint requests

Z. Naturforsch. **60c**, 22–24 (2005); received August 4/September 14, 2004

From an EtOAc extract of *Dracocephalum subcapitatum*, five flavonoids, calycopterin, xanthomicrol, isokaempferide, luteolin and apigenin, together with five terpenoids, oleanolic acid, ursolic acid, geranial, neral and limonene-10-al, were isolated. Among them, citral and limonene-10-al were the most effective components against epimastigotes of *Trypanosoma cruzi*, the parasitic agent of Chagas disease.

*Key words:* *Dracocephalum subcapitatum*, Labiatae, Limonene-10-al