

**stichting
mathematisch
centrum**



AFDELING ZUIVERE WISKUNDE
(DEPARTMENT OF PURE MATHEMATICS)

ZW 165/81

SEPTEMBER

A.M. COHEN

A CHARACTERIZATION OF SUBSPACES OF GIVEN RANK
IN A PROJECTIVE SPACE

Preprint

kruislaan 413 1098 SJ amsterdam

Printed at the Mathematical Centre, 413 Kruislaan, Amsterdam.

The Mathematical Centre, founded the 11-th of February 1946, is a non-profit institution aiming at the promotion of pure mathematics and its applications. It is sponsored by the Netherlands Government through the Netherlands Organization for the Advancement of Pure Research (Z.W.O.).

A characterization of subspaces of given rank in a projective space^{*)}

by

Arjeh M. Cohen

ABSTRACT

A theorem by Cooperstein that partially characterizes the natural geometry $A_{n,d}(F)$ of subspaces of rank $d-1$ in a projective space of rank n over a finite field F , is somewhat strengthened and generalized to the case of an arbitrary division ring F .

Moreover, this theorem is used to provide characterizations of $A_{n,2}(F)$ and $A_{5,3}(F)$ which will be of use to characterizations of other (exceptional) Lie group geometries.

KEY WORDS & PHRASES: *projective geometry, Grassmann varieties*

^{*)} This report will be submitted for publication elsewhere.

1. INTRODUCTION

Theorem A by Cooperstein in [2] provides a partial characterization of the geometry $A_{a,d}(F)$ on all subspaces of rank (= projective dimension) $d-1$ of a projective space of rank a over a finite field F . Though there are more (partial) characterizations, cf. [5], [6], this one has the advantage of being ready-made for characterizations of geometries corresponding to groups of Lie type, see for instance Theorem B of [2]. This note deals with a generalization of Theorem A to the case of a projective space of finite rank over an arbitrary division ring F . The present version is stronger than the original theorem in that it describes more specifically what happens in 'case (iii)'. In fact, it shows that case (iii) does not occur at all if the geometry is finite.

However, many steps in the proof are taken from or inspired by Cooperstein's proof of Theorem A. The infinite case (i.e. where the geometry and hence F is infinite) depends on the classification of polar spaces of rank 3 (used in 4.2) as given in [7].

Two applications of the theorem are given: a characterization of the lines in a projective space of finite rank, and a characterization of the planes in a projective space of rank 5. Precise formulation of the results will be given in Section 2 after some notation and terminology has been introduced.

2. TERMINOLOGY, NOTATION AND MAIN RESULT.

An *incidence system* (P, L) is a set P of *points* together with a collection L of subsets of cardinality > 1 , called *lines*. If (P, L) is an incidence system then the point graph or collinearity graph of (P, L) is the graph (P, Γ) whose vertex set is P and whose edges consist of the pairs of collinear points. The incidence system is called *connected* whenever its collinearity graph is connected. Likewise terms such as *(co)cliques*, *paths* will be applied freely to (P, L) when in fact they are meant for (P, Γ) . We let $d(x, y)$ for $x, y \in P$ denote the ordinary distance in (P, Γ) and write

$$\Gamma_i(x) = \{y \in P \mid d(x, y) = i\}.$$

Also $\Gamma(x) = \Gamma_1(x)$ and $x^\perp = \{x\} \cup \Gamma(x)$.

For a subset X of P and $y \in P$ we write $d(y, X) = \min_{x \in X} d(y, x)$,

$$X^\perp = \bigcap_{x \in X} x^\perp \text{ and } \Gamma(X) = \bigcup_{x \in X} \Gamma(x).$$

(P, L) is called *nondegenerate* if $P^\perp = \emptyset$.

A subset X of P is called a *subspace* of (P, L) whenever each point of P on a line bearing two distinct points of X is itself in X . A subspace X is called *singular* whenever it induces a clique in (P, Γ) . The length i of a longest chain $X_0 \subsetneq X_1 \subsetneq \dots \subsetneq X_i = X$ of nonempty singular subspaces X_j of X is called the *rank* of X and denoted by $\text{rk}(X)$.

For a subset X of P , the subspace generated by X is denoted $\langle X \rangle$. Instead of $\langle X \rangle$ we also write $\langle x_1, Y \rangle$ if $X = \{x_1\} \cup Y$, and so on.

If F is a family of subsets of P and X is a subset of P , then $F(X)$ denotes the family of members of F contained in X , while F_X denotes the family of members of F containing X . If $X = \{x\}$ for some $x \in P$, we often write F_x instead of $F_{\{x\}}$. Furthermore, if H is another family of subsets of P , then $F(H)$ denotes $\{F(H) \mid H \in H\}$.

If G is a group of automorphisms of (P, L) such that $L \not\subseteq x^G$ for any $x \in P$ and $L \in L$, then $(P, L)/G$ denotes the quotient of (P, L) by G , i.e. the incidence system whose points are the orbits in P of G and whose lines are of the form $\{x^G \mid x \in L\}$ for $L \in L$. The incidence system (P, L) is called *linear* if any two distinct points are on at most one line. If x, y are collinear points of a linear incidence system, then xy denotes the unique line through them; thus $xy = \langle x, y \rangle$.

A line is called *thick* if there are at least three points on it, otherwise it is called *thin*. Recall (from [2]) that (P, L) is a *polar space* if $|x^\perp \cap L| \neq 1$ implies $L \subseteq x^\perp$ for any $x \in P$ and $L \in L$ that the rank of a polar space is the maximal number $k \geq 1$ such that there exists a chain $\emptyset = V_0 \subsetneq V_1 \subsetneq \dots \subsetneq V_k$ of singular subspaces in (P, L) and that a *generalized quadrangle* is a polar space of rank 2. The objects under study here are incidence systems (P, L) in which the following four axioms hold:

(P1) for any $x \in P$ and $L \in L$ with $|x^\perp \cap L| > 1$ the line L is entirely contained in x^\perp (this means (P, L) is a Gamma space in D.G. Higman's terminology)

- (P2) the connected components of (P, L) are not complete.
- (P3) For any two $x, y \in P$ with $d(x, y) = 2$, the subsets $x^\perp \cap y^\perp$ forms a subspace isomorphic to a nondegenerate generalized quadrangle.
- (P4) For $x \in P$, $L \in L$ such that $x^\perp \cap L = \emptyset$ but $x^\perp \cap L^\perp \neq \emptyset$ the subset $x^\perp \cap L^\perp$ is a line.

For ease of reference and with the result below in mind, an incidence system (P, L) satisfying (P1), (P2), (P3), (P4) (but not necessarily connected) will be called a *Grassmann space*. The incidence structure whose points are the subspaces of rank d of a projective space over a division ring F of rank n and whose lines are the subspaces incident to an incident pair x, y of a subspace x of rank $d+1$ and a subspace y of rank $d+1$, is denoted by $A_{n, d+1}(F)$.

MAIN THEOREM. (P, L) is a connected Grassmann space with thick lines all whose singular subspaces have finite ranks iff one of the following holds

- (i) (P, L) is a nondegenerate polar space of rank 3 with thick lines.
- (ii) There are $a \geq 4$, $d \leq (a+1)/2$ and a division ring F such that $(P, L) \cong A_{a, d}(F)$.
- (iii) There are $d \geq 5$, an infinite division ring F and an involutory automorphism σ of $A_{2d-1, d}(F)$ induced by a polarity of the underlying projective space over F of rank $2d-1$, with $d(x, x^\sigma) \geq 5$ for all points x of $A_{2d-1, d}(F)$, such that $(P, L) \cong A_{2d-1, d}(F) / \langle \sigma \rangle$.

This theorem is proved in Section 6.

APPLICATIONS. Suppose (P, L) is an incidence system with thick lines.

- (i) (P, L) is a Grassmann space all whose singular subspaces have finite ranks and in which $x^\perp \cap L^\perp \neq \emptyset$ for any $x \in P$ and $L \in L$ iff (P, L) is either a nondegenerate polar space of rank 3 or isomorphic to $A_{a, 2}(F)$ for some $a > 4$ and some division ring F .
- (ii) (P, L) is a Grassmann space in which for any two intersecting lines $L_1, L_2 \in L$ and any point $z \in P$ there exists $u \in z^\perp$ with $u^\perp \cap L_1 \neq \emptyset$ and $u^\perp \cap L_2 \neq \emptyset$ iff (P, L) is either a nondegenerate polar space of rank 3 or isomorphic to one of $A_{4, 2}(F)$, $A_{5, 3}(F)$ for some division ring F .

These applications are treated in Section 7.

3. PRELIMINARY RESULTS

Throughout this section, (P, L) will be a Grassmann space.

The definitions of generalized quadrangles and polar spaces and some of their properties can be found in [2]. We shall first recall some facts from [2] whose proofs do not depend on any finiteness assumptions.

LEMMA 3.1. *Let (P, L) be a Grassmann space. Then (P, L) is linear and is determined by its collinearity graph in the sense that for any two distinct collinear $x, y \in P$, $\{x, y\}^{\perp\perp}$ is the unique line on x, y . Moreover, we have*

- (i) *maximal cliques are singular subspaces;*
- (ii) *for any clique X of P , the subspace $\langle X \rangle$ is singular;*
- (iii) *if X is a subset of P , then X^\perp is a subspace;*
- (iv) *if x, y, z form a clique of P not contained in a line, then $\{x, y, z\}^\perp$ is a maximal singular subspace.*

PROPOSITION 3.2. (Cooperstein) *Let (P, L) be a Grassmann space. For any $x, y \in P$ with $d(x, y) = 2$, the subset $S(x, y)$ defined by $S(x, y) = \{z \in P \mid (\forall L \in L)(L \subseteq \{x, y\}^\perp \Rightarrow z^\perp \cap L \neq \emptyset)\}$ is a subspace isomorphic to a polar space of rank 3 with the property that $z^\perp \cap S$ is a singular subspace for any $z \in P \setminus S$.*

As a matter of fact, (P_4) is not needed for the lemma and the proposition. The proof of Proposition 3.2 can be found in [2] though some care has to be taken to relax the condition that lines are thick (cf. [1]). The family of all $S(x, y)$ obtained as described above will be denoted by S , and the family of all maximal cliques will be denoted by M . A member of S will be called a *symp* or a *hyperline*; a maximal singular subspace will often be called *max space* for short.

COROLLARY 3.3.

- (i) *Each singular subspace of rank ≤ 2 is contained in a symp. Hence, it is a point, a line or a projective plane;*
- (ii) *If M is a singular subspace and M properly contains a line, then M is a projective space.*

We shall denote the family of singular subspaces of rank 2 by V and call its members *planes*.

REMARK 3.4. Axiom (P4) can be replaced by

$$(P4)' \quad (\forall S \in \mathcal{S}) (\forall x \in P \setminus S) (|x^\perp \cap S| > 1 \Rightarrow (x^\perp \cap S) \in V)$$

PROOF. $(P4) \Rightarrow (P4)'$. Let $|x^\perp \cap S| > 1$ for $S \in \mathcal{S}$ and $x \in P \setminus S$. By the above proposition, $x^\perp \cap S$ is a singular subspace of S and hence of rank 1 or 2. Take $z \in x^\perp \cap S$ and $y \in S \setminus (x^\perp \cup z^\perp)$. Apply (P4) to the point y and the line $L = xz$. Since $y^\perp \cap (x^\perp \cap S) \neq \emptyset$, as S is a polar space and $x^\perp \cap S$ contains a line, we have $y^\perp \cap L^\perp \neq \emptyset$. Moreover, $u \in y^\perp \cap L$ would yield $u \in y^\perp \cap z^\perp$; hence $u \in S \setminus \{z\}$ and $x \in uz$, so $x \in S$, which is absurd. Therefore $y^\perp \cap L = \emptyset$, so that $y^\perp \cap L^\perp$ is a line contained in $x^\perp \cap S$ but not on z . It follows that $x^\perp \cap S$ is a plane.

$(P4) \Leftarrow (P4)'$. Suppose $x \in P$ and $L \in \mathcal{L}$ are such that $x^\perp \cap L = \emptyset$ and $x^\perp \cap L^\perp \neq \emptyset$. Take $y \in L$ and consider $S = S(x, y)$. Since $\langle y, x^\perp \cap L^\perp \rangle$ is a singular subspace of S of rank ≥ 1 , it is a plane by (P4)'. It follows that $x^\perp \cap L^\perp$ is a line, as wanted \square .

LEMMA 3.5. If S is a symp and $x, y \in P \setminus S$ are collinear, while $x^\perp \cap S \in V$ and $y^\perp \cap S \neq \emptyset$, then either $y^\perp \cap S \subseteq x^\perp \cap S$ or $y^\perp \cap S \in V$ and $x^\perp \cap y^\perp \cap S$ is a singleton.

PROOF. Suppose $z \in y^\perp \cap S \setminus x^\perp$. First of all we show that $y^\perp \cap S$ is a plane, too. As $x^\perp \cap S \in V(S)$ and S is a polar space, $z^\perp \cap x^\perp \cap S$ is a line in S . Now both $z^\perp \cap x^\perp \cap S$ and y are in the generalized quadrangle $x^\perp \cap z^\perp$, so there is $u \in x^\perp \cap S$ with $\{u\} = x^\perp \cap z^\perp \cap S \cap y^\perp$. Since $uz \subseteq y^\perp \cap S$, Remark 3.4 implies that $y^\perp \cap S$ is a plane. Finally, $x^\perp \cap y^\perp \cap S = z^\perp \cap x^\perp \cap y^\perp \cap S = \{u\}$ \square

COROLLARY 3.6. If $S \in \mathcal{S}$ and $M \in \mathcal{M}$ satisfy $|M \cap S| > 1$, then $M \cap S \in V(S)$.

PROOF. For any $w \in M \setminus S$, we have $w^\perp \cap S \in V(S)$ by Remark 3.4. If $z, w \in M \setminus S$, then $z^\perp \cap S = w^\perp \cap S$ by Lemma 3.5. If $M \subseteq S$, there is nothing to prove; so assume $M \setminus S \neq \emptyset$. Taking $z \in M \setminus S$, we get $z^\perp \cap S = \bigcap_{w \in M \setminus S} w^\perp \cap S = \bigcap_{w \in M} w^\perp \cap S =$

$= M^\perp \cap S = M \cap S$. In particular, $M \cap S = z^\perp \cap S \in V(S)$. \square

Let S be a symp. On the set of planes $V(S)$ a graph $(V(S), \sim)$ is defined by $V_1 \sim V_2$ iff $\text{rk}(V_1 \cap V_2) = 0$ ($V_1, V_2 \in V(S)$). It is well known that $(V(S), \sim)$ has either one or two connected components. In the latter case, each line is in precisely two members of $V(S)$, one of each connected component, and the connected components are complete graphs.

COROLLARY 3.7. *Let $S \in \mathcal{S}$ and let K be a union of connected components of $(V(S), \sim)$. Then*

$$H(K, S) = \bigcup_{K \in K} K^\perp$$

is a subspace on S .

PROOF. As $S = \bigcup_{K \in K} K$, the subset $H(K, S)$ clearly contains S . We need only show that if $x, y \in P \setminus S$ are collinear and $x^\perp \cap S, y^\perp \cap S \in K$, then any $z \in xy$ is contained in $H(K, S)$. If $x^\perp \cap y^\perp \cap S = x^\perp \cap S$, then clearly $z^\perp \cap S = x^\perp \cap S \in K$, so we are done. Therefore, we may assume $x^\perp \cap y^\perp \cap S = \{u\}$ for some $u \in P$. Consequently, $z \in P \setminus S$. Take $v \in x^\perp \cap S \setminus \{u\}$ and $w \in v^\perp \cap y^\perp \cap S \setminus \{u\}$ (note that w exists because $v, y^\perp \cap S$ are in the polar space S). Now x, y, w, v is a 4-circuit and $z \in xy$, so that there is $z_1 \in z^\perp \cap vw$. Note that $z_1 \neq u$, for otherwise $v \in uw$, whence $v \in y^\perp \cap S$ conflicting $v \neq u$. Thus $|z^\perp \cap S| > 1$ as $z_1, u \in z^\perp \cap S$, and we are done by Remark 3.4 and Lemma 3.5. \square

LEMMA 3.8. (i) *If $M \in \mathcal{M}$ and $x \in P \setminus M$ satisfy $x^\perp \cap M \neq \emptyset$, then $x^\perp \cap M \in \mathcal{L}$.*
(ii) *If $M \in \mathcal{M}$ and $L \in \mathcal{L}$ with $\text{rk}(L \cap M) = 0$, then there is a unique $N \in \mathcal{M}$ with $M \cap N \in \mathcal{L}$.*

PROOF. (i) Suppose $z \in x^\perp \cap M$. Take $y \in M \setminus x^\perp$ and consider $S = S(x, y)$. If $M \subseteq S$, there is nothing to prove. Otherwise, $M \cap S$ contains z and y , so $M \cap S \in V(S)$ by Corollary 3.6. It results that $x^\perp \cap M = x^\perp \cap (M \cap S)$ is a line.

(ii) By (i), $L^\perp \cap M$ is a line. Thus $N = \langle L, L^\perp \cap M \rangle^\perp$ is the unique max space containing L with $M \cap N \in \mathcal{L}$ \square

Notice that Lemma 3.8(ii) can be reformulated as (L_x, M_x) is a general-

ized quadrangle for each $x \in P$.

LEMMA 3.9. *The graph (V, \approx) defined by $V_1 \approx V_2$ iff $V_1 \subseteq V_2^\perp$ and $V_1 \cap V_2 \in L$ for $V_1, V_2 \in V$ is connected. In particular, any plane V is contained in a symp.*

PROOF. Note that the subgraph induced on $V(S)$ is connected for any $S \in S$. Let $V \in V$. By connectedness of (P, L) , it suffices to prove that any plane W with $V \cap W \neq \emptyset$ is joined to V by a path in (V, \approx) . Let $W \in V \setminus \{V\}$ with $V \cap W \neq \emptyset$. Take $v \in V \setminus W$ and $w \in W \setminus V$. If $v \notin w^\perp$, consider $S(v, w)$. There are planes M, N in $S(v, w)$ such that $\langle v, V \cap W \rangle \subseteq M$ and $\langle w, V \cap W \rangle \subseteq N$. Now $\text{rk}(M \cap V) > \text{rk}(V \cap W)$ and $\text{rk}(N \cap W) > \text{rk}(V \cap W)$, so by induction we are reduced to the case where $V \subseteq W^\perp$. It suffices to treat the case where $V \cap W \in L$.

Since symps exist we may assume $V \subseteq S$ for some $S \in S$. Let U be a plane in S with $V \cap U = V \cap W$. Again, take $v \in V \setminus W$, $w \in W \setminus V$ and $u \in U \setminus V$. Then $u \notin v^\perp$ and $w \in v^\perp$. If $w \in u^\perp$, then $w \in u^\perp \cap v^\perp \subseteq S(u, v) = S$, and $W = \langle w, V \cap W \rangle \subseteq \langle u^\perp \cap v^\perp, U \cap V \rangle \subseteq S$. So we may assume $w \notin u^\perp$. But then $W \subseteq S(u, v)$, finishing the proof of the Lemma. \square

COROLLARY 3.10. *The graph (M, \approx) defined by $M_1 \approx M_2$ iff $\text{rk}(M_1 \cap M_2) = 1$, is connected.*

PROOF. Note that M_1 and M_2 are adjacent in (M, \approx) iff there are planes $V \subseteq M_1$ and $W \subseteq M_2$ with $V \not\subseteq W^\perp$ and $V \cap W \in L$. Thus there is a surjective morphism $(V, \approx) \rightarrow (M, \approx)$ of graphs given by $V \mapsto V^\perp$ (cf. Lemma 3.1 (iv)). The desired result is therefore a consequence of the above lemma. \square

FROM NOW ON WE ASSUME THAT THE LINES OF (P, L)
ARE THICK

LEMMA 3.11. *The graph (L, \sim) defined by $L_1 \sim L_2$ iff $\text{rk}(L_1 \cap L_2) = 0$ and $L_1 \not\subseteq L_2^\perp$, is connected.*

PROOF. As before, the proof comes down to the case where $L_1 \subset L_2^\perp$ and $\text{rk}(L_1 \cap L_2) = 0$. But then $\langle L_1, L_2 \rangle \in V$, so the lemma results from the analogous statement for polar spaces with thick lines. \square

LEMMA 3.12. *Let $L_1, L_2 \in L$. There is a bijection between*

$$M(L_1) \text{ and } M(L_2).$$

PROOF. By connectedness of (L, \sim) as defined in Lemma 3.11, we need only prove the lemma for $L_1, L_2 \in L$ with $L_1 \not\subseteq L_2^\perp$ and $L_1 \cap L_2$ is a point. Take $x \in L_1 \setminus L_2$ and $y \in L_2 \setminus L_1$ and let $u: M(L_1) \rightarrow M(L_2)$ be given by $u(M) = \langle y, M \cap y^\perp \rangle^\perp$. It is not hard to verify that u is a bijection. \square

LEMMA 3.13. *Let $M, N \in M$ satisfy $\text{rk}(M \cap N) = 0$. Then $\text{rk}(M) = \text{rk}(N)$.*

PROOF. $M \cap N = \{u\}$ for some $u \in P$. In view of 3.8, the map $\phi: L_u(M) \rightarrow L_u(N)$ given by $\phi(X) = X^\perp \cap N$ is well defined. Moreover, it is an isomorphism of projective spaces. Hence the result. \square

Consider the graph (M, \approx) defined by $M_1 \approx M_2$ iff $\text{rk}(M_1 \cap M_2) = 0$. The above lemma states that the members of a connected component of (M, \approx) all have the same rank. Lemma 3.8(ii) and connectedness of (P, L) yield that for any line L and each connected component K of (M, \approx) there is a member of K on L . The following lemma shows that in fact (M, \approx) cannot have more than two connected components.

LEMMA 3.14. *Suppose a line is contained in at least three max spaces. Then (M, \approx) is connected. In particular, all max spaces have the same rank.*

PROOF. By Lemma 3.12, any line is contained in at least three max spaces. Let M, N be two max spaces with $M \cap N \in L$. We claim the existence of $K \in M$ with $K \cap M = K \cap N$ a singleton.

In view of Corollary 3.10 it follows that (M, \approx) is connected. The last statement is then a direct consequence of 3.13. To show the existence of K as described choose $x \in M \cap N$ and $y \in (M \cap N)^\perp \setminus (M \cap N)$. Note that y exists because of the assumption that $M \cap N$ is in at least three members of M . By Lemma 3.9, $\langle M \cap N, y \rangle$ is contained in a symp, so there is $z \in P$ with $z^\perp \cap \langle M \cap N, y \rangle = \langle x, y \rangle$. Now $K = \langle x, y, z \rangle^\perp \in M$ and $\{x\} \subseteq K \cap M = z^\perp \cap (y^\perp \cap M) = z^\perp \cap (M \cap N) = \{x\}$ by Lemma 3.8. So $K \cap M = \{x\}$. Similarly, $K \cap N = \{x\}$, so the claim holds. \square

LEMMA 3.15. *If $\text{rk}(M) = 2$ for some $M \in M$, then for any $x \in P$ and $L \in L$ we have $x^\perp \cap L^\perp \neq \emptyset$. In particular, the diameter of (P, L) is 2.*

PROOF. We may assume that $x^\perp \cap L = \emptyset$. By induction with respect to $d(x, L)$, it suffices to prove the first statement in the case where $d(x, L) = 2$. Let $y, z \in P$ be such that $x \in y^\perp$ and $z \in y^\perp \cap L$, and take $w \in L \setminus \{z\}$. The hypothesis implies that there is a max space N of rank 2 on yz . Since $x^\perp \cap N$ and $w^\perp \cap N$ are lines in N , they intersect in a point, say u . Since $u \in x^\perp \cap w^\perp \cap N \subseteq x^\perp \cap w^\perp \cap z^\perp = x^\perp \cap L^\perp$, we have shown $x^\perp \cap L^\perp \neq \emptyset$ as wanted. \square

COROLLARY 3.16. *If all max spaces have rank 2, then (P, L) is a polar space of rank 3.*

PROOF. Let $x \in P$ and $L \in L$. We prove the Buekenhout-Shult axiom $x^\perp \cap L \neq \emptyset$. Suppose the contrary. Then, since the above lemma yields $x^\perp \cap L^\perp \neq \emptyset$, axiom (P4) implies that $x^\perp \cap L^\perp$ is a line disjoint from L . Thus $\text{rk}(\langle L, x^\perp \cap L^\perp \rangle) = 3$, conflicting the hypothesis. \square

LEMMA 3.17. *If $S \in S$ and $x \in P$ satisfy $L_x \subseteq L(S)$, then (P, L) is a polar space of rank 3.*

PROOF. We prove that $P = S$. In view of the connectedness of (P, L) it suffices to show that for any $y \in x^\perp$ all $z \in y^\perp$ are contained in S . Let y, z be as described. If $z \in x^\perp \setminus \{x\}$ we must have $zx \in L(S)$, so $z \in S$. Suppose $z \notin x^\perp$. Then $S(x, z)$ is a symp on x . But since symps are geodesically closed, S is the only symp on x . We obtain $S(x, z) = S$, and $z \in S$ as wanted. \square

4. A PROPERTY OF CLASSICAL GENERALIZED QUADRANGLES

Throughout this section, (P, L) is a generalized quadrangle with thick lines (P, L) called *classical* whenever it occurs as the residue of a point in a nondegenerate polar space of rank 3 whose lines are thick. Since polar spaces of this rank are classified [7] the list of all classical generalized quadrangles is known. The result is quoted in Theorem 4.1. For the duration of this section, we shall adopt terminology from [7], without recalling all definitions. The aim of this section is to prove Proposition 4.2.

THEOREM 4.1. (Buekenhout-Shult, Veldkamp, Tits). *Let (P, L) be a classical generalized quadrangle. Then (P, L) is one of the following:*

- (i) *A polar space $Q(\pi)$ of a projective space over a division ring F where π is a polarity determined by a nondegenerate trace-valued (σ, ϵ) -hermitian form of Witt index 2 for some antiautomorphism σ of F with $\sigma^2 = 1$ and some $\epsilon \in \{1, -1\}$.*
- (ii) *A polar space $Q(\kappa)$ of a projective space over a division ring F where κ is a projective pseudo-quadratic form represented by a nondegenerate σ -quadratic form of Witt index 2 for some antiautomorphism σ of F with $\sigma^2 = 1$.*
- (iii) *The dual of the generalized quadrangle $Q(\kappa_0)$ in a projective space over the field F defined in (ii) where κ_0 is represented by the quadratic form $q: E \times F^4 \rightarrow F$ over F defined by*

$$(x_0, x_1, x_2, x_3, x_4) \rightarrow N(x_0) - x_1x_3 + x_2x_4$$

for E a Cayley division algebra over the field F and $N: E \rightarrow F$ the quadratic norm form of this algebra.

- (iv) $\{x, y\}^\perp$ *for two noncollinear points x, y of $A_{3,2}(F)$.*

A grid is by definition a generalized quadrangle in which each point is precisely two lines. Clearly the generalized quadrangles in (iv) are grids. In Lemma 4.5 we shall find all grids occurring in the list. But first, the main result of this section will be stated.

We recall that a family R of lines in (P, L) is called a *spread* in (P, L) if the members of R partition P (i.e. $P = \bigcup_{L \in R} L$ and for any two distinct $L_1, L_2 \in R$ we have $L_1 \cap L_2 = \emptyset$).

A grid has precisely two spreads, they are also called the *parallel classes* of the grid. If L_1, L_2 are disjoint lines of (P, L) such that the subspace $\langle L_1, L_2 \rangle$ is a grid, then $L_1 L_2$ denotes the parallel class of the grid containing L_1 and L_2 .

PROPOSITION 4.2. *Let (P, L) be a nondegenerate generalized quadrangle with thick lines which is either finite or classical. Suppose it admits a spread R in which for any two distinct $L_1, L_2 \in R$ the subspace $\langle L_1, L_2 \rangle$ is a grid and the family $L_1 L_2$ is contained in R such that*

$(R, \{L_1, L_2 \mid L_1, L_2 \in R; L_1 \neq L_2\})$ is a projective space. Then the rank of R as a projective space is 1 and (P, L) is a grid.

The remainder of this section is devoted to the proof of this proposition. Thus, from now on until the end of this section we assume that R is a spread of the generalized quadrangle (P, L) . In the next lemma, the finite case is dealt with by a straightforward computational argument.

LEMMA 4.3. *If (P, L) is finite; then $\text{rk}(R) = 1$.*

PROOF. Suppose $\text{rk}(R) > 1$. Then (P, L) is not a grid. In particular, it is then a regular generalized quadrangle, i.e. there is a constant number, say $1+t$, of lines through each point, and a constant number of points, say $1+s$, on each line. By well-known theory [3], we have $t \leq s^2$. On the other hand, $1+st = |R| = \frac{s^{m+1}-1}{s-1}$ if the rank of R is m . It follows that $t = 1+s$ (and $m = 2$). A straightforward computation on multiplicities of eigenvalues of the adjacency matrix of the collinearity graph (cf. [3]) leads to integrality conditions which are only satisfied if $s = 1$. But this is excluded by the requirement that the lines be thick. \square

The assumption that lines are thick is necessary, since the regular complete bipartite graph on 6 points provides a counterexample.

The classical case depends on the classification of classical generalized quadrangles as stated in Theorem 4.1. If (P, L) is as in (iv) of this theorem, there is nothing to prove.

LEMMA 4.4. *(P, L) is not isomorphic to a generalized quadrangle as described in 4.1 (iii).*

PROOF. If (P, L) satisfies (iii) of 4.1, then the dual of $\langle L_1, L_2 \rangle$ is a bipartite graph in the dual of (P, L) . On the other hand, according to 10.7 of [7], the dual of $\langle L_1, L_2 \rangle$ is the dual of (P, L) itself. This yields the absurdity that $Q(\kappa_0)$ of 4.1 (iii) is a bipartite graph. \square

If X is a subset of a projective space we denote by $[X]$ the projective subspace of this projective space spanned by X .

LEMMA 4.5. *Let F be a division ring, let σ be an antiautomorphism of F such that $\sigma^2 = 1$ and let $\epsilon \in \{1, -1\}$. Suppose ξ is either a polarity π determined by a nondegenerate trace valued (σ, ϵ) -hermitian form f of Witt index 2 or a projective pseudo-quadratic form κ represented by a nondegenerate σ -quadratic form q of Witt index 2. If L_1, L_2 are lines of $Q(\xi)$ with $L_1 \cap L_2 = \emptyset$ such that $\langle L_1, L_2 \rangle$ is a grid, then $\sigma = 1$ and (if $\xi = \pi$) $\epsilon = 1$. Thus, F is a field. Furthermore, $\langle L_1 \cup L_2 \rangle = [L_1 \cup L_2] \cap Q(\xi)$ unless $\xi = \pi$ and F has characteristic 2.*

PROOF. Take distinct points e_1, e_3 in L_1 and e_2, e_4 in L_2 such that $\{e_2\} = e_3^\perp \cap L_2$ and $\{e_4\} = e_1^\perp \cap L_2$. Put $F_{\sigma, \epsilon} = \{t - t^\sigma \epsilon \mid t \in F\}$. As in 8.10 of [7], choose E_1, E_2, E_3, E_4 , points of the vector space underlying the projective space in which $Q(\xi)$ is defined, such that E_i represents e_i (i.e. such that the ray through E_i is e_i for $i = 1, 2, 3, 4$) and such that

$$f\left(\sum_{i=1}^4 E_i x_i, \sum_{i=1}^4 E_i y_i\right) = x_1^\sigma y_2 + \epsilon x_2^\sigma y_1 + x_3^\sigma y_4 + \epsilon x_4^\sigma y_3 \quad \text{if } \xi = \pi$$

and

$$q\left(\sum_{i=1}^4 E_i x_i\right) = x_1^\sigma x_2 + x_3^\sigma x_4 + F_{\sigma, \epsilon} \quad \text{if } \xi = \kappa.$$

Now take $a \in F_{\sigma, \epsilon}$ (where $\epsilon = 1$ if $\xi = \kappa$). Then the calculation performed in 8.10 of [7] shows that the projective point $p(a)$ represented by $(1, a, 0, 0)$ on the basis E_1, E_2, E_3, E_4 is in $\langle L_1, L_2 \rangle$. But $p(a)$ is collinear with both e_3 and e_4 and hence in $\{e_1, e_2\}$ as $\langle L_1, L_2 \rangle$ is a grid. It follows that $a = 0$, and the conclusion is that $F_{\sigma, \epsilon} = \{0\}$.

If $\epsilon = -1$, this reads $t + t^\sigma = 0$ for all $t \in F$, so that F has characteristic 2 and $\epsilon = 1$.

It results that $\epsilon = 1$ and $t - t^\sigma = 0$ for all $t \in F$, whence $\sigma = 1$. Since σ is an anti-automorphism, F must be commutative and therefore a field.

The final statement of the lemma now results from (8.10) of [7]. \square

LEMMA 4.6. *Let F be a field and let ξ be either a polarity π determined by a nondegenerate symmetric form f of Witt index 2 or a projective quadratic form κ represented by a nondegenerate quadratic form q of Witt index 2.*

Suppose L_1, L_2, L_3 are distinct lines of $Q(\xi)$ such that for each $i \in \{1, 2, 3\}$ the subspace $\langle L_i \cup L_{i+1} \rangle$ of $Q(\xi)$ is a grid and $L_i \cap \langle L_{i-1} \cup L_{i+1} \rangle = \emptyset$ (indices modulo 3). Then there are lines $N_1 \in L_1 L_2 \setminus \{L_1, L_2\}$ and $N_2 \in L_1 L_3 \setminus \{L_1, L_3\}$ such that if $\langle N_1 \cup N_2 \rangle$ is a grid, the intersection $N_1 N_2 \cap L_2 L_3$ does not contain a line of $Q(\xi)$ which is disjoint with L_1 .

PROOF. Let e_1, e_2, e_3, e_4 and E_1, E_2, E_3, E_4 be as in the proof of 4.5. Thus $e_1, e_3 \in L_1$; $e_1 \neq e_3$; $e_2, e_4 \in L_2$ and $e_1 \in e_4^\perp$, $e_2 \in e_3^\perp$; furthermore the vector E_i represents e_i for $i = 1, 2, 3, 4$ and

$$\left. \begin{aligned} f\left(\sum_{i=1}^4 E_i x_i, \sum_{i=1}^4 E_i y_i\right) &= x_1 y_2 + x_2 y_1 + x_3 y_4 + x_4 y_3 & \text{if } \xi = \pi \\ \text{and} \\ q\left(\sum_{i=1}^4 E_i x_i\right) &= x_1 x_2 + x_3 x_4 & \text{if } \xi = \pi \end{aligned} \right\}$$

Next, take $e'_5 \in L_3$ with $\{e'_5\} = e_1^\perp \cap L_3$ and $e_5 \in e_1 e'_5$ with $\{e_5\} = e_2^\perp \cap e_1 e'_5$. Then $e_5 \in \langle L_1 \cup L_3 \rangle$ so there is a line $L'_3 \in R$ on e_5 contained in $\langle L_1 \cup L_3 \rangle$. Since $e_1 \neq e_5$, we may replace L_3 by L'_3 without harming generality, so as to obtain $e_5 \in e_1^\perp \cap e_2^\perp \cap L_3$. Let $e_6 \in L_3$ be such that $\{e_6\} = e_3^\perp \cap L_3$ and let $e'_4 \in L_2$ be such that $\{e'_4\} = e_6^\perp \cap L_2$. The projective space $A = [L_1 \cup L_2 \cup L_3]$ has rank 3, 4 or 5.

Let us first consider the case where $\text{rk}(A) = 5$. If $\xi = \pi$, then $\text{char}(F) \neq 2$ as otherwise the Witt index would be strictly larger than 2. So we may assume that $\xi = \kappa$. Consider $q|_{[L_1 \cup L_3]}$. Let $\gamma \in F$ and E'_4 a vector representing e'_4 be such that $E'_4 = E_4 + E_2 \gamma$ (note that $e'_4 \neq e_2$). It is easily derived that there are vectors E_5, E_6 representing e_5, e_6 such that

$$q(E_2 x_2 + E_4 x_4 + E_5 x_5 + E_6 x_6) = x_2 x_6 + x_4 x_5 + \gamma x_4 x_6.$$

Considering $q|_{[L_1 \cup L_3]}$, we obtain $\alpha, \beta \in F \setminus \{0\}$ such that

$$q(E_1 x_1 + E_3 x_3 + E_5 x_5 + E_6 x_6) = \alpha x_1 x_6 + \beta x_3 x_5. \quad (x_i \in F).$$

The foregoing restrictions describe $q|_A$ fully:

$$q\left(\sum_{i=1}^6 E_i x_i\right) = x_1 x_2 + x_3 x_4 + x_2 x_6 + x_4 x_5 + \alpha x_1 x_6 + \beta x_3 x_5 + \gamma x_4 x_6 \quad (x_i \in F)$$

Now let $n_1(n_2, n_3, n_4$ resp.) be the point of $Q(k) \cap A$ whose homogeneous coordinates with respect to E_1, E_2, \dots, E_6 are $(1, 0, 0, 0, 1, 0)$ $((0, 0, -\alpha, 0, 0, \beta)$, $(1, 0, 0, 1, 0, 0)$, $(0, -1, 1, 0, 0, 0)$ resp.). Then $N_1 = n_1 n_2$ is a line of $L_1 L_3$ and $N_2 = n_3 n_4$ is a line of $L_1 L_2$.

Note that $N_1 \cap N_2 = \emptyset$ as $L_1 L_2 \cap L_1 L_3 = \{L_1\}$. Now suppose $\langle N_1 \cup N_2 \rangle$ is a grid with $\{N\} = N_1 N_2 \cap L_2 L_3$ for a line N of $Q(k)$. Then clearly $N \neq N_1, N_2$. Moreover $e_2 e_5$ is a line of $\langle L_2 \cup L_3 \rangle$ not parallel to L_2 , so $e_2 e_5 \cap N \neq \emptyset$. But a point of $N \setminus \langle L_1 \cup L_2 \rangle$ has homogeneous coordinates of the form $v + \lambda u$ for $\lambda \in F$, where $v = (\zeta, 0, -\alpha, 0, \zeta, \beta)$ and $u = (1, -\eta, \eta, 1, 0, 0)$ for $\zeta, \eta \in F$ are homogeneous coordinates of a point in N_1, N_2 respectively. Thus $e_2 e_5 \cap N \neq \emptyset$ implies the existence of $\zeta, \eta, \lambda, \mu, v \in F$ such that

$$(1 + \lambda \zeta, -\eta, \eta - \lambda \alpha, 1, \lambda \zeta, \lambda \beta) = (0, \mu, 0, 0, v, 0)$$

The equation leads to a classical contradiction in the fourth coordinate. This proves the lemma in the case where $\text{rk}(A) = 5$.

Next, assume that $\text{rk}(A) \leq 4$. Then $L_3 \cap [L_1 \cup L_2]$ so $L_3 \cap ([L_1 \cup L_2] \cap Q(\xi) \setminus \langle L_1 \cup L_2 \rangle) \neq \emptyset$. According to Lemma 4.3, this implies that F has characteristic 2 and that $\xi = \pi$. In particular, π is a symplectic form.

If $\text{rk}(A) = 4$, then π is degenerate and has a kernel consisting of a single (projective) point z . Clearly $z \notin A \cap Q(\xi)$, so we may consider the quotient by $[z]$ so as to reduce the proof to the case where $\text{rk}(A) = 3$.

Thus, for the rest of the proof, we have that F has characteristic 2, that $\text{rk}(A) = 3$ and that $\xi = \pi$ is a polarity determined by the symplectic form whose restriction to A is given by

$$f\left(\sum_{i=1}^4 E_i x_i, \sum_{i=1}^4 E_i y_i\right) = x_1 y_2 + x_2 y_1 + x_3 y_4 + x_4 y_3 \quad (x_i, y_i \in F)$$

A straightforward computation using $e_5 \in \{e_1, e_2\}^\perp$ yields the existence of $\alpha \in F \setminus \{0\}$ such that E_5 given by $(0, 0, \alpha, 1)$ on the basis E_1, E_2, E_3, E_4 represents e_5 .

Also, $e_6 \in \{e_3, e_5\}^\perp$ leads to the existence of $\beta \in F \setminus \{0\}$ such that the vector E_6 given by $(1, \beta, 0, 0)$ on the same basis, represents e_6 .

Now let $n_1(n_2, n_3, n_4$ resp.) be the point of $Q(\pi)$ whose homogeneous coordinates with respect to E_1, E_2, E_3, E_4 are $(1, 0, \alpha, 1)((1, \beta, \beta, 0), (1, 0, 0, 1), (0, 1, 1, 0)$ resp.). Then $N_1 = n_1 n_2$ is a line in $L_1 L_3$ and $N_2 = n_3 n_4$ is a line in $L_1 L_2$. Now $\langle N_1 \cup N_2 \rangle$ is a grid. Put $N = \langle N_1 \cup N_2 \rangle \cap \langle L_2 \cup L_3 \rangle$. Let x, y be the point of $Q(\pi)$ whose homogeneous coordinates with respect to E_1, E_2, E_3, E_4 are

$$X = \begin{cases} (0, \alpha, \alpha, 1) & \text{if } \alpha = \beta \\ (0, (\zeta+1)\alpha, \zeta\alpha, \zeta) \text{ where } \zeta^2 = \alpha(\alpha+\beta), & \text{if } \alpha \neq \beta \end{cases}$$

and

$$Y = (\alpha, \alpha\beta, 0, \alpha + \eta\beta), \text{ where } \eta^2 = \alpha(\alpha+\beta)/\beta^2$$

respectively. Then x, y are distinct collinear points of N and $X\eta\beta^2 + Y(\alpha+\eta\beta) = (\alpha^2 + \eta\alpha\beta, 0, \alpha^2\beta, 0) (= X\alpha^2 + Y\alpha \text{ if } \alpha = \beta)$ represents a point of xy on L_1 .

It follows that $\{xy\} = N_1 N_2 \cap L_2 L_3$, so that $N_1 N_2 \cap L_2 L_3$ does not contain a line of $Q(\pi)$ which is disjoint with L_1 . This settles the lemma. \square

The classical case of Proposition 4.2 is dealt with by the following lemma.

LEMMA 4.7. *If (P, L) is classical, then $\text{rk}(R) = 1$.*

PROOF. In view of 4.4 and the observation, made before, that (P, L) is a grid in case (iv) of 4.1, we need only consider cases (i) and (ii). Let L_1, L_2 be two lines from R . Then $L_1 \cap L_2 = \emptyset$ and $\langle L_1 \cup L_2 \rangle$ is a grid, so by Lemma 4.5 we may assume that $(P, L) = Q(\xi)$ for ξ as described in the hypotheses of Lemma 4.6. Suppose we have $L_3 \in R \setminus L_1 L_2$. Then $L_i \cap L_{i+1} = \emptyset$ and $\langle L_i \cup L_{i+1} \rangle = \emptyset$ for each $i \in \{1, 2, 3\}$ (indices taken modulo 3). By Lemma 4.6, however, there are $N_1 \in L_1 L_2 \setminus \{L_1, L_2\}$ and $N_2 \in L_1 L_3 \setminus \{L_1, L_3\}$ such that $N_1 N_2 \cap L_2 L_3$ does not contain a member of R . This means that Pasch's axiom is not satisfied, contradicting that $(R, \{L_1 L_2 \mid L_1, L_2 \in R; L_1 \neq L_2\})$ is a projective space. The conclusion is that $R = L_1 L_2$, in other words, that $\text{rk}(R) = 1$. \square

5. THE POINT RESIDUE OF A GRASSMANN SPACE

We continue the study of Grassmann spaces. In this section (P, L) is a connected Grassmann space whose lines are thick. Furthermore, ∞ is a fixed point of P and $P^\infty, L^\infty, S^\infty, M^\infty$ stand for $L_\infty, L_\infty(V_\infty), L_\infty(S_\infty), L_\infty(M_\infty)$ respectively. Moreover, if $V \in V_\infty$, then V^∞ denotes $L_\infty(V)$. Similarly for members of L, S and M .

It is straightforward to check that (P^∞, L^∞) is a connected incidence system of diameter 2 satisfying axioms (P1) and (P2). By 3.1, the members of M^∞ are maximal singular subspaces of (P^∞, L^∞) isomorphic to projective spaces and of the form L^\perp for any line L contained in them. Moreover, (P^∞, M^∞) is a generalized quadrangle by the remark following 3.8, which is easily seen to be nondegenerate. Members of S lead to generalized quadrangles in (P^∞, L^∞) . We shall call them *quads*. Any two noncollinear points are in a unique quad. Also, if $S \in S^\infty$ and $x \in P^\infty - S$, then $x^\perp \cap S$ is either empty or a line of (P^∞, L^∞) . This is immediate from (P4)'.

We recall from 3.7 that for $S \in S$ and $M \in M$, the subset $S \cup \{z \in M \setminus S \mid z^\perp \cap S \in V\}$ is denoted by $H(V(S), S)$. We shall also write $H(S)$ instead of $H(V(S), S)$.

LEMMA 5.1. *Suppose there are $M \in M$ and $S \in S$ with $M \cap S = \{\infty\}$. Then $M \cap H(S)$ is a subspace of M of rank at most 2.*

PROOF. Set $V = M \cap H(S)$. It follows from 3.7 that V is a subspace of M . Recall that $M^\infty, S^\infty, V^\infty$ denote the subspaces of (P^∞, L^∞) induced by M, S, V respectively. Let R be the subfamily of L^∞ whose members occur as $z^\perp \cap S^\infty$ for some $z \in V^\infty$. Then R is a spread of the quad S^∞ , for any two members of R are disjoint (in P^∞) by 3.5 and if $x \in S^\infty$, then $x^\perp \cap M^\infty = \{y\}$ for some $y \in V^\infty$ by Lemma 3.8(i), whence $y^\perp \cap S^\infty$ is a member of R on x . Now let L be a line of (P^∞, L^∞) in V^∞ . Then $U = \bigcup_{x \in L} x^\perp \cap S^\infty$ is a grid in S^∞ . For suppose there are $x_1, y_1 \in U$ with $x_1 \in y_1^\perp \setminus \{y_1\}$. Then there are unique $x, y \in L$ with $x_1 \in x^\perp \cap S^\infty$ and $y_1 \in y^\perp \cap S^\infty$. If $z_1 \in x_1 y_1$, then either $x = y$ and $z_1 \in x^\perp \cap S^\infty$ or $x \neq y$. In the latter case x, y, y_1, x_1 is a 4-circuit, so there is $z \in xy$ with $z_1 \in z^\perp \cap S^\infty$. So U is a subspace. Proceeding with $x \neq y$, we see that $x_1 y_1$ and $x^\perp \cap S^\infty$ are the only two lines on x_1 in U , so U is indeed a grid in S^∞ . Moreover, one parallel class of lines in U is entirely contain-

ed in \mathcal{R} . Denoting by $L_1 L_2$ for $L_1, L_2 \in \mathcal{R}$ the parallel class of lines in $\langle L_1, L_2 \rangle$ belonging to \mathcal{R} , we obtain a surjective morphism

$$u: (V^\infty, L^\infty(V^\infty)) \rightarrow (\mathcal{R}, \{L_1 L_2 \mid L_1 L_2 \in \mathcal{R}; L_1 \neq L_2\})$$

of projective spaces given by $u(x) = x^\perp \cap S^\infty$ ($x \in V^\infty$). If $x_1, x_2 \in V^\infty$ satisfy $x_1^\perp \cap S^\infty = x_2^\perp \cap S^\infty$, then $N = \langle x_1, x_2, x_1^\perp \cap S^\infty \rangle$ is a singular subspace with $\text{rk}(M^\infty \cap N) \geq \text{rk}(\langle x_1, x_2 \rangle)$. Since $N \cap S^\infty = x_1^\perp \cap S^\infty \neq \emptyset$, we have $M^\infty \neq N$, whence $\text{rk}(M^\infty \cap N) \leq 0$. It results that $\text{rk}(\langle x_1, x_2 \rangle) = 0$, i.e. $x_1 = x_2$. This shows that u is bijective, so that $\text{rk}(V) = \text{rk}(V^\infty) + 1 = \text{rk}(\mathcal{R}) + 1 \leq 2$ by Proposition 4.2. \square

LEMMA 5.2. *Suppose $M \in M^\infty$ and $S, T \in S^\infty$ satisfy $S \cap T \neq \emptyset$, $M \cap S \neq \emptyset$ and $M \cap T \neq \emptyset$. Then $M \cap S \cap T \neq \emptyset$.*

PROOF. Let $x \in S \cap T$ and $u \in M \cap S$, $w \in M \cap T$. If $x \in M$ or $u = w$, we are done. So assume $x \notin M$ and $u \neq w$. Now $x \in w^\perp$ would imply $w \in u^\perp \cap x^\perp \subseteq S$ if $u \notin x^\perp$ and $x \in (uw)^\perp = M$ otherwise; similarly $x \in v^\perp$ can be settled. Assume $x \notin u^\perp \cup w^\perp$. Then there is a unique point y in $x^\perp \cap M$. We have $y \in \{x, u\}^\perp \cap \{x, w\}^\perp \subseteq S \cap T$, so $y \in M \cap S \cap T$. \square

LEMMA 5.3. *Assume that for any $M \in M^\infty$ and $S \in S^\infty$, we have $M \cap S \neq \emptyset$. Then $M^\infty = L^\infty$ and $|S^\infty| = 1$, so that (P, L) is a polar space of rank 3.*

PROOF. Fix $x \in P^\infty$. Suppose S, T are distinct quads on x . Write $L = S \cap T$. We shall first show that L is a line. Indeed, it is a singular subspace on x , so L is either a point or a line. Choose $M \in M^\infty$ not on x . By Lemma 5.2, there must be a point y in $M \cap S \cap T$, so that $xy \subseteq L$. It follows that $L = xy$ is a line. If $N \in M^\infty$ is disjoint from L^\perp , we get a contradiction with $N \cap L = \emptyset$. Since such N exist, it follows that S is the only quad on x . Therefore, S contains all points in P^∞ noncollinear with x . But for each point $z \in x^\perp \setminus \{x\}$, there is a point $u \in z^\perp \setminus x^\perp$, so that $z \in x^\perp \cap u^\perp \subseteq S$. This shows that $P^\infty = S$. Thus the maximal cliques are members of L^∞ , i.e. $M^\infty = L^\infty$. Finally, by Lemma 3.17, the Grassmann space (P, L) must be a polar space of rank 3. \square

LEMMA 5.4. *If $\text{rk}(M_0) = 2$ for some $M_0 \in M$, then $x^\perp \cap M \neq \emptyset$ for any $x \in P$ and any $M \in M$ of rank > 2 .*

PROOF. Suppose $M \in M$ is of rank > 2 and $x \in P \setminus M$. In view of the connectedness of (P, L) , we may restrict attention to the case where there are $z \in P$ and $y \in M$ such that $z \in x^\perp \cap y^\perp$. As $y \in z^\perp \cap M$, we have $z^\perp \cap M \in L$ by Lemma 3.8(i). If $x^\perp \cap L \neq \emptyset$, we are done. So assume $x^\perp \cap L = \emptyset$. Now $z \in x^\perp \cap L^\perp$, so $x^\perp \cap L^\perp \in L$ by (P4), so $\langle x^\perp \cap L^\perp, L \rangle$ is a projective space of rank 3 on L . But M is the unique space on L of rank > 2 by 3.13 and 3.14, so $z \in x^\perp \cap L^\perp \subseteq M$; in particular $z \in x^\perp \cap M$, terminating the proof. \square

LEMMA 5.5. *Suppose $\text{rk}(M_0) = 2$ for some $M_0 \in M$. If both $M_1, M_2 \in M$ have rank > 2 , then $|M_1 \cap M_2| = 1$.*

PROOF. We only need to establish $M_1 \cap M_2 \neq \emptyset$ in view of 3.14. Suppose $M_1 \cap M_2 = \emptyset$. Take $x \in M_1$. By the previous lemma and Lemma 3.8 (i), $L = x^\perp \cap M_2$ is a line. Take $v, w \in L$ with $v \neq w$ and consider $B = v^\perp \cap M_1$ and $C = w^\perp \cap M_1$. If $B = C$, then $\langle B, L \rangle$ is a projective space of rank 3 on L so is contained in M_2 , which conflicts $M_1 \cap M_2 = \emptyset$. Thus $B \neq C$. Now B, C are lines on x in M_1 , so $\text{rk}(\langle B, C \rangle) = 2$ and there is $y \in M_1 \setminus \langle B, C \rangle$. But $y^\perp \cap L = \emptyset$ so $A = y^\perp \cap L^\perp \in L$ as $x \in A$ by (P4). Consequently, $\langle A, L \rangle$ has rank 3 and contains L , so is in M_2 . It results that A is in M_2 , whence $x \in M_1 \cap M_2$. \square

COROLLARY 5.6. *Suppose there are $M_1, M_2 \in M$ with $\text{rk}(M_1) = 2$ and $\text{rk}(M_2) = m > 2$. Let M^+ (M^- resp.) be the connected component of (M, \sim) whose members have rank $m(2$ resp.). Then (M^+, P^{M^+}) , where $P^{M^+} = \{M_x^+ | x \in P\}$, is a projective space of rank $m+1$ such that the points and lines of (P, L) correspond to the lines and pencils of (M^+, P^{M^+}) respectively. In other words, (P, L) is isomorphic to $A_{m+1,2}(F)$ for some division ring F .*

PROOF. We verify Tallini's axioms in [6]. First of all, it is obvious that no line is a maximal singular subspace.

(I) Any two members of M^+ meet in exactly one point.

This is the content of Lemma 5.5.

(II) If $M \in M^+$ and $M_1 \in M^-$ then $M \cap M_1$ is either empty or a line.

This follows from Lemmas 3.13 and 3.1 (iv).

(III) For any line L there is exactly one $M \in M^+$ and one $M_1 \in M^-$ such that $L = M \cap M_1$.

This results from the remarks preceding Lemma 3.14.

The Corollary now follows from Proposition I in [6].

Instead of referring to [6], a direct proof could have been given, but this would have lengthened the paper by another few pages.

LEMMA 5.7. Assume that each line is in at least three max spaces. If $M \cap S$ is empty for $M \in M^\infty$ and $S \in S^\infty$, then $\{x \in M \mid x^\perp \cap S \in L^\infty\}$ contains a subspace which is a projective plane.

PROOF. Take $x \in S$. It has a unique neighbor y in M . As $L_1 = y^\perp \cap S$ contains x , it must be a line on x . Let L be another line in S on x , and take $x_2 \in L \setminus \{x\}$. There is $y_2 \in x_2^\perp \cap M_1$. Note that $y \neq y_2$ for $y^\perp \cap S$ is a clique and $x_2 \notin L_1^\perp$. Write $L_2 = y_2^\perp \cap S_1$. This is a line disjoint from L_1 (cf. 3.5). Suppose L^1 is a third line on x , not in $L_1^\perp \cup L^\perp$ (such a line exists by assumption). Take $w \in L^1 \setminus \{x\}$. If $w \notin S$, then $w^\perp \cap S$ contains x , so must be a line in S distinct from L_1 and L . Therefore there is a point $x_3 \in x^\perp \cap S \setminus (L_1 \cup L)$. Again, take $y_3 \in x_3^\perp \cap M$ and consider $y_3^\perp \cap S$. It is a line on x not in $\langle L_1, L_2 \rangle$. Thus $y_3 \notin yy_2$ and $\langle y, y_2, y_3 \rangle \in V$ is the subspace of the desired kind. \square

COROLLARY 5.8. Each line of (P, L) is in precisely two max spaces, unless (P, L) is a polar space of rank 3.

PROOF. Suppose there is a line in strictly more than two max spaces. Let $S \in S$ and $M \in M$ satisfy $M \cap S = \{\infty\}$ and consider $V = M \cap H(S)$ (cf. 3.7 and 5.1). By 5.1, $\text{rk}(V) \leq 2$ and by 5.7, $\text{rk}(V) \geq 3$, contradiction. It results that the conditions of Lemma 5.3 are satisfied, so that (P, L) is a polar space of rank 3. \square

We summarize the results obtained in this section.

PROPOSITION 5.9. Let (P, L) be a connected Grassmann space with thick lines. Assume (P, L) is not isomorphic to a polar space (of rank 3) or $A_{n,2}(F)$ for some $n \geq 4$ and some division ring F . Then for each point $x \in P$, the residue

(P_x, M_x) is a grid. In particular, each line is in precisely two max spaces. Moreover, the rank of any max space is > 2 .

6. COOPERSTEIN'S THEOREM A.

Throughout this section, (P, L) is a connected Grassmann space whose lines are thick such that any line is in precisely two max spaces, each of them of rank > 2 . We fix a point ∞ of P and maintain the notation of Section 5 concerning residues on ∞ .

LEMMA 6.1. *Let $M, N \in M$ and $S \in S$.*

- (i) *If $M \cap S \neq \emptyset$, $N \cap S \neq \emptyset$ and $M \cap N \neq \emptyset$, then $M \cap N \cap S$ is a singular subspace.*
- (ii) *If $M \cap S, N \cap S \in V$ and $\text{rk}(M \cap N \cap S) = 0$, then $M \approx N$.*

PROOF. Both (i) and (ii) follow from the fact that $x^\perp \cap S$ is a singular subspace for any $x \in P \setminus S$. \square

We supply the graph (M, \approx) with the natural family of lines that turns M into a Gamma space whose collinearity graph is (M, \approx) . To avoid confusion, we denote by M^T for $M \in M$ (rather than M^\perp which has a distinct interpretation) the set of vertices in (M, \approx) of distance at most 1 to M . For $M_1, M_2 \in M$ with $M_1 \approx M_2$, the line $M_1 M_2$ is defined by $M_1 M_2 = \{M_1, M_2\}^{TT}$. A priori, it is not clear that this turns M into a linear incidence system, but it will follow from 6.3 that it is. By C we denote the family of all such lines, i.e.

$$C = \{M_1 M_2 \mid M_1, M_2 \in M, M_1 \approx M_2\}.$$

We need some more notation. For $x \in P$, $L \in L$, $V \in V$ and $M \in M$ with $x \in L \subseteq V \subseteq M$, denote by $p(L, M)$ the unique member of M containing L and distinct from M . Furthermore, put $\ell(x, V) = \{p(L^1, V^\perp) \mid L^1 \in L(V)_x\}$, $m(x, M) = \{p(L^1, M) \mid L^1 \in L(M)_x\}$ and

$$n(x, V) = \left\{ p(L', W^\perp) \mid \begin{array}{l} W \in V_x; W \cap p(L'', V^\perp) \in L \\ \text{for each } L'' \in L(V)_x; L' \in L(W) \end{array} \right\}.$$

LEMMA 6.2. *Two distinct max spaces M_1, M_2 are of distance 2 in (M, \approx) iff $M_1 \cap M_2 = \emptyset$ and there is $M \in M$ with $M \cap M_1, M \cap M_2 \in L$. In particular, (M, \approx) is not complete.*

PROOF. Suppose M_1, M_2 are of distance 2 in (M, \approx) . Then $M_1 \cap M_2 = \emptyset$, for if $M_1 \cap M_2 \in L$ then $\{M_1, M_2\}^\top = \emptyset$ by 3.8(ii). Let $H \in \{M_1, M_2\}^\top$. There are $x_i \in P$ with $H \cap M_i = \{x_i\}$ for $i = 1, 2$. Consider $L_1 = x_2^\perp \cap M_1$ and $L_2 = x_1^\perp \cap M_2$. By 3.8 we know $L_i \in L$. Now $\langle x_1, x_2, L_i \rangle^\perp$ ($i = 1, 2$) and H are three max spaces on $x_1 x_2$, while H differs from the first two as it intersects M_1 and M_2 in a point. So $M = \langle x_1, x_2, L_1, L_2 \rangle^\perp$ is a max space with $M \cap M_i = L_i$.

Conversely, let M be a max space with $M \cap M_i \in L$ for $i = 1, 2$ and suppose $M_1 \cap M_2 = \emptyset$. Take $x_i \in M \cap M_i$ and consider $H = p(x_1 x_2, M)$. By 3.8, $H \cap M_i = \{x_i\}$ so that $H \in \{M_1, M_2\}^\top$. Thus M_1, M_2 have distance ≤ 2 ; in fact their distance is ≥ 2 as $M_1 \cap M_2 = \emptyset$. This establishes that M_1, M_2 are of distance 2.

Finally, let $M \in M$ and let $L_1, L_2 \in L$ be disjoint lines (they exist as $\text{rk}(M) \geq 3$). Then $p(L_1, M)$ and $p(L_2, M)$ are not joined as they have distance 2 by the above criterion. \square

LEMMA 6.3. *Suppose $M_1, M_2 \in M$ satisfy $M_1 \approx M_2$. Let $V \in V_\infty$ be such that $M_1 \cap M_2 = \{\infty\}$ and $V \cap M_1, V \cap M_2 \in L$. Then $\{M_1, M_2\}^\top = m \cup n$, where $m = m(\infty, V^\perp)$ and $n = n(\infty, V)$ are maximal cliques in (M, \approx) with $m \cap n = \ell(\infty, V)$. Moreover, if for $Y \in m$ and $Y' \in n$ we have $Y \approx Y'$, then at least one of Y, Y' is in $\ell(\infty, V)$. In particular, $M_1 M_2 = \ell(\infty, V)$.*

PROOF. Clearly $\ell(\infty, V) = \ell(\infty, V')$ for any $V' \in V_\infty$ with $V' \cap M_1, V' \cap M_2 \in L$, as V^∞ and V'^∞ are parallel lines of a grid in (P^∞, L^∞) on the 4-circuit $(V' \cap M_1)^\infty, (V \cap M_1)^\infty, (V \cap M_2)^\infty, (V' \cap M_2)^\infty$.

Let us now determine $\{M_1, M_2\}^\top$. By Lemma 6.2, any two distinct members X, X' of m satisfy $X \cap X' = \{\infty\}$, so m is a clique contained in $\{M_1, M_2\}^\top$.

If $M \in n$, there are $W \in V_\infty$ with $W \cap p(L^1, V^\perp) \in L$ for each $L' \in L(V)_\infty$ and $L \in L$ with $L = M \cap W$. Note that $M_1 \cap W \in L$ since $(W^\perp)^\infty$ is parallel to $(V^\perp)^\infty$ in (P^∞, M^∞) . As $\ell(\infty, V) = \ell(\infty, W)$, we may replace W by V without loss of generality. If $M \in m$, we have $M \in \ell(\infty, V)$ as before. Assume $\infty \notin L$, and let $i \in \{1, 2\}$. Since $V \cap M_i$ and L are lines in V , there is a point $x_i \in V$ such

that $\{x_i\} = L \cap M_i$. It follows by 3.8(ii) that $M \cap M_i = \{x_i\}$ (note that M and V^\perp meet in xx_i). Thus $M \in \{M_1, M_2\}^\perp$.

So far, we have shown $m \cup n \subseteq \{M_1, M_2\}$. The converse is straightforward: each member M of $\{M_1, M_2\}^\perp$ is in m whenever $M \cap M_i = \{\infty\}$ and in n otherwise. Let us now consider $m \cap n$. The inclusion $\ell(\infty, V) \subseteq m \cap n$ is obvious. To prove the opposite inclusion, let $M \in m \cap n$. Since $M \in n$, there is $W \in V_\infty$ with $L' \subseteq W$ and $W \cap M_i \in L$ for $i = 1, 2$ with $M = p(L', W^\perp)$, so that $M \cap W = L'$; but $\infty \in M$ as $M \in m$, so $\infty \in L'$ and $M \in \ell(\infty, W)$. By the first paragraph of this proof, this is equivalent to $M \in \ell(\infty, V)$. This proves $m \cap n = \ell(\infty, V)$. Next, let $Y \in m$ and $Y' \in n \setminus \ell(\infty, V)$ with $Y \approx Y'$. Write $L = Y \cap V^\perp$ and let x, y_1, y_2 be such that $\{x\} = Y \cap Y'$, $Y' \cap M_1 = \{y_1\}$, $Y' \cap M_2 = \{y_2\}$. Then $\{x, y_1, y_2\}$ is a clique (in Y'). Note that $y_1 \neq y_2$ as $Y' \notin \ell(\infty, V)$. If $x \notin y_1 y_2$, then $\langle x, y_1, y_2 \rangle \in V$ and $\langle x, y_1, y_2 \rangle^\perp = Y^\perp$. But $L \subseteq \langle y_1, y_2, x \rangle^\perp$, so $L \subseteq Y'$ contradicting that $Y \cap Y'$ is a singleton. Hence $x \in y_1 y_2$, so that $L = \infty x$ is in V and $Y \in \ell(\infty, V)$. This establishes the one but last claim of the lemma. The last one follows directly. \square

COROLLARY 6.4. (M, C) is a Gamma space with thick lines, where $C = \{\ell(x, V) \mid x \in P, V \in V_x\}$.

PROOF. It follows from the previous lemma that (M, C) is a Gamma space and that the definition of C coincides with the one given in the text preceding 6.2. Thickness of the lines is a consequence of the bijection $L \rightarrow \ell(x, V)$ for fixed $L \in L(V)$ with $x \notin L$ given by $y \rightarrow p(xy, V^\perp)$ ($y \in L$) \square

LEMMA 6.5. If $X_1, X_2 \in M$ are of distance 2 in (M, C) , then $\{X_1, X_2\}^\perp$ is a grid.

PROOF. Note that $X_1 \cap X_2 = \emptyset$ and there is $H \in M$ with $L_i = H \cap X_i \in L$ for each $i \in \{1, 2\}$. We claim that any $Y \in \{X_1, X_2\}^\perp$ meets X_i in a point of L_i . For let $y_1 \in X_1 \setminus L_1$. Then $y_1^\perp \cap L_2 = \emptyset$ as otherwise $y_1^\perp \cap K$ would contain more than the line L_1 , leading to $y_1 \in K$ and $K = X_1$, conflicting $X_1 \cap X_2 = \emptyset$.

Thus $L_1 = y_1^\perp \cap L_2^\perp$ by (P4), so that $y_1^\perp \cap X_2 \neq \emptyset$ whence $X_1 \cap X_2 \neq \emptyset$, which is absurd. We conclude that $y_1^\perp \cap X_2 = \emptyset$, so that no max space on y_1 is in X_2^\perp . This implies that any $Y \in \{X_1, X_2\}^\perp$ meets X_1 in a point of L_1 . Similarly the claim is proved for $i = 2$. The claim yields the following description of the subspace under study:

$$\{X_1, X_2\}^T = \{p(L, K) \mid L \in L(K), L \cap L_1 \neq \emptyset, L \cap L_2 \neq \emptyset\}.$$

The lines in $\{X_1, X_2\}^T$ are of the form $\ell(x, \langle x, L_i \rangle)$ for $x \in L_j$, where $\{i, j\} = \{1, 2\}$.

In particular, $\{X_1, X_2\}^T$ is isomorphic to the geometry on the lines intersecting two given lines L_1, L_2 in a projective space of rank 3, in which two members are collinear whenever they intersect. This geometry is well known and easily checked to be that of a grid (cf. Section 4). \square

LEMMA 6.6. *Suppose $\ell \in C$ and $M \in M$ satisfy $M^T \cap \ell = \emptyset$ and $M^T \cap \ell^T \neq \emptyset$. Then $M^T \cap \ell^T \in C$.*

PROOF. It suffices to show that $M^T \cap \ell^T$ contains at least two points.

Let $x \in P$ and $V \in V_x$ be such that $\ell = \ell(x, V)$. Suppose $H \in M^T \cap \ell^T$. Then by Lemma 6.3 we have $H \in m \cup n$, where $m = m(x, V^\perp)$ and $n = n(x, V)$. Let $y \in P$ be such that $H \cap M = \{y\}$. Suppose $H \in m$. Then $x \in H$. Write $L = x^\perp \cap M$. This is a line on y , so $W = \langle x, L \rangle$ is a plane. Take $L' \in L(W)_x$ with $y \notin L'$, then $p(L', W^\perp)$ is a member of $M^T \cap \ell^T$ distinct from H .

Suppose $H \in n \setminus m$. Then $d(x, y) = 2$. Consider $S = S(x, y)$. Note that $y^\perp \cap V$ is a line of $x^\perp \cap y^\perp$. Let L be a line of $x^\perp \cap y^\perp$ parallel (and distinct) to $y^\perp \cap V$, and define $H' = \langle y, L \rangle^\perp$. Then $H' = p(L', \langle x, L' \rangle^\perp) \in n$ so $H' \in M^T \cap n \subseteq M^T \cap \ell^T$. As $H' \neq H$, we are done. \square

COROLLARY 6.7. *(M, C) is a Grassmann space with thick lines.*

PROOF. Axiom (P1) is proved in 6.4 (where it is also stated that lines are thick), (P2) in 6.2, (P3) in 6.5 and (P4) in 6.6. \square

LEMMA 6.8. *Take $M \in M_\infty$ of rank i . Then $m(\infty, M)$ is a projective space in (M, C) of rank $i-1$. If $V \in V_\infty$ is such that $V \cap M \in L$, then $n(\infty, V)$ is a projective space of rank $i+1$.*

PROOF. Write $m = m(\infty, M)$ and $n = n(\infty, V)$. By Corollary 3.3, both m and n are projective spaces. By construction of m , there is a bijective map $\mu: M^\infty \rightarrow m$ given by $\mu(L^\infty) = p(L, M)$ for $L \in L_\infty$. Given $L' \in V_\infty$ with $V' \subseteq M$, we have that $\mu(\{L \in L_\infty \mid L \subseteq V'\}) = \{p(L, (V')^\perp) \mid L \subseteq V'\} = \ell(\infty, V')$, so that μ maps lines of (P^∞, L^∞) in M^∞ onto lines of (M, C) in m . As $\text{rk}(M^\infty) = i-1$, this shows that $\text{rk}(m) = i-1$.

Next, consider n . Choose $L_1, L_2 \in L(V)_\infty$ distinct and write $M_i = p(L_i, V^\perp)$. Furthermore let H_i be a hyperspace of M_i disjoint from ∞ . Given $x_1 \in H_1$, the line $x_1^\perp \cap M_2$ on ∞ intersects H_2 in a point x_2 .

This leads to a map $\psi: H_1 \rightarrow n$ given by $\psi(x_1) = p(x_1, x_2, \langle \infty, x_1 x_2 \rangle^\perp)$. This map is easily seen to be injective. Moreover, if L_1 is a line of H_1 , then $\psi(L_1)$ is a line of (M, C) .

Let $L_2 = \bigcup_{x \in L_1} x^\perp \cap H_2$ and take $x_1, y_1 \in L_1$, $x_1 \neq y_1$. Then there are unique $x_2, y_2 \in L_2$ collinear with x_1, y_1 respectively. Consider the generalized quadrangle $x_1^\perp \cap y_2^\perp$. It contains the lines ∞x_2 and ∞y_1 , so there is a point $\infty' \in x_1^\perp \cap y_2^\perp \cap x_2^\perp \cap y_1^\perp \setminus \{\infty\}$. Now $\psi(L_1) = \ell(\infty', \langle \infty', L_1 \rangle)$ is a line in N .

As a consequence, $\psi(H_1)$ is a singular subspace of n of rank $i-1$ (note that M_1 is of rank i). But $\ell(\infty, V)$ is a line of n completely disjoint from $\psi(H_1)$. We conclude that $\text{rk } n \geq i+1$. We finish by showing that any member N of n is on a line in (M, C) from a member of $\ell(\infty, V)$ to a member of $\psi(H_1)$. By analogous arguments to what we have seen before, we are easily led to the case where $N = p(y_1 y_2, V)$ for distinct y_i in $N \cap M_1 \setminus \{\infty\}$ ($i = 1, 2$). Let $x_i \in N \cap H_i$, so that $x_i^\perp = y_i^\perp$. If $x_1 = y_1$ and $x_2 = y_2$ then $N \in \psi(H_1)$, so we may assume that $x_1 x_2 \neq y_1 y_2$. Since both lines are in V , there is $z \in V$ with $x_1 x_2 \cap y_1 y_2 = \{z\}$. Now $N = p(y_1 y_2, V)$ is on the line $\ell(z, V)$ which has member $p(\infty z, V)$ in $\ell(\infty, V)$ and member $p(x_1 x_2, V)$ in $\psi(H_1)$.

We conclude that n is spanned by $\ell(\infty, V)$ and $\psi(H_1)$. Thus $\text{rk}(n) \leq i+1$, and equality holds. \square

Before stating the main theorem, we recall the notion of quotient. The quotient of the incidence system $A = (P_1, L_1)$ by the group G of automorphisms of A is meant to be the incidence system A/G whose point set is $P_1/G = \{x^G \mid x \in P_1\}$, the set of orbits of G in P_1 , and whose family of lines is $L_1/G = \{L^G \mid L \in L_1\}$, where $L^G = \{x^G \mid x \in L\}$ for $L \in L_1$. Note that this quotient is again an incidence system if $L \not\subseteq x^G$ for each $x \in P$, $L \in L$.

THEOREM 6.9. *Let (P, L) be a connected Grassmann space with thick lines, whose max spaces have finite ranks. Then one of the following holds:*

- (i) (P, L) is a nondegenerate polar space of rank 3 with thick lines.

- (ii) $(P, L) \cong A_{a,d}(F)$ for some $a \geq 4$, $d \leq (a+1)/2$ and some division ring F .
- (iii) There is a natural number $d \geq 5$, a division ring F and an involutory automorphism σ of $A = A_{2d-1,d}(F)$, interchanging the connected components of the graph (M, \sim) on the max spaces, with $d(x, x^\sigma) \geq 5$ for all points x of A such that $(P, L) \cong A/\langle \sigma \rangle$.

PROOF. By 3.13 and 3.14, $\text{rk}(M)$ for $M \in M$ attains at most two values. Let d be the minimal of these and let b be the other one if it exists, let $b = d$ otherwise. The proof runs by induction on d . The case $d = 2$ has been settled in Proposition 5.9.

Assume $d > 2$, and suppose (P, L) is not a polar space of rank 3. By 5.9 we have that each line is in exactly two max spaces. By 6.7 and 6.8, there is a connected component M^+ of (M, \sim) such that the induced subgraph (M^+, \sim) is the collinearity graph of the connected Grassmann space (M^+, C^+) where $C^+ = \{\ell \in C \mid \ell \cap M^+ \neq \emptyset\}$, whose max spaces have ranks $d-1$, $b+1$. The induction hypothesis then yields that (ii) occurs, so that $(M^+, C^+) \cong A_{d+b-1, d-1}(F)$ for some division ring F . Now $A_{d+b-1, d}(F)$ can be thought of as the incidence system obtained from $A_{d+b-1, d-1}(F)$ by taking the max spaces of rank $d-1$ from one connected component under \sim in $A_{d+b-1, d-1}(F)$ for points and the relation \approx (i.e. $M \approx N$ iff $M \cap N$ meet in a point) for collinearity. Remember that this determines $A_{d+b-1, d}$ as any Grassmann space is determined by its collinearity graph (cf. 3.1).

Let (P', L') be the incidence system that can be obtained from (M^+, C^+) in just the way $A_{d+b-1, d}(F)$ is obtained from $A_{d+b-1, d-1}(F)$ (note that this makes sense as $(M^+, C^+) \cong A_{d+b-1, d-1}(F)$).

If $m \in P'$, then m is a projective space in (M, C) of rank $d-1$, so $m = m(x, M)$ for a unique $x \in P$ and some $M \in M \setminus m$. Thus there is a map $\mu: P' \rightarrow P$ sending $m \in P'$ to the unique $x \in P$ for which there is $M \in M \setminus m$ with $m = m(x, M)$. This map is clearly surjective and is either 2:1 or 1:1 according as $M = M^+$ or not, i.e. according as (M, \sim) has one or two connected components. We claim that μ is a morphism of graphs. For if m, n are collinear in (P', L') , the points $\mu(m)$ and $\mu(n)$ are both contained in the max space M for which $m \cap n = \{M\}$. Consequently, $\mu(m)$ and $\mu(n)$ are collinear in (P, L) . Thus, if (M, \sim) is disconnected, we have $(P, L) \cong (P', L') \cong A_{d+b-1, d}(F)$.

Let from now on (M, \sim) be connected. Now μ is a surjective 2:1 morphism. Also $b = d$ in view of 3.13 and 3.14 so $(P', L') \cong A_{2d-1, d}(F)$ for some $d \geq 3$. Choose $m \in P'$. We shall show that μ is bijective when restricted to the neighborhood m^\perp of m in (P', L') . Let x, y be distinct collinear points of P and suppose $\mu(m) = x$ and let $m_1, m_2 \in P'$ both be collinear with m and such that $\mu(m_1) = \mu(m_2) = y$.

As before, we may assume that $m = m(x, M)$ for $M \in M$ with $xy \subseteq M$. Similarly we may take $M_i \in M$ with $xy \subseteq M_i$ such that $m_i = m(y, M_i)$, for each $i \in \{1, 2\}$. Suppose now that $M_1 \neq M_2$. Since each of M, M_1, M_2 contains xy , it follows that M coincides with M_1 or M_2 . Without loss of generality we may assume that $M = M_1$. Since m_2 is collinear with m , there is $Y \in M$ such that $Y \in m(x, M) \cap m(y, M_2)$. This means that $M \cap Y$ is a line on x and $M_2 \cap Y$ is a line on y . Thus Y contains xy , so either $Y = M$ or $Y = M_2$. But $Y = M$ conflicts $M \cap Y \in L$ and $Y = M_2$ conflicts $M_2 \cap Y \in L$. It results that $M_1 = M_2$, so that $m_1 = p(y, M_1) = p(y, M_2) = m_2$. We have established that the restriction of μ to the members of P' collinear with a given point is injective.

Our next step is to show that the restriction of μ to the subset m^\perp of P' of members collinear with m is an isomorphism of graphs. Thus for $m_1, m_2 \in P' \setminus \{m\}$ collinear with m such that $x_1 = \mu(m_1)$, $x_2 = \mu(m_2)$ are collinear in P , we have to derive that m_1 is collinear with m_2 in (P', L') . Let $V = \langle x, x_1, x_2 \rangle$, where $x = \mu(m)$.

Since $m \sim m_i$, there are $X_i \in m \cap m_i$ for $i = 1, 2$. Thus X_i contains xx_i . If $x_1 \in xx_2$, then $X_1 \cap X_2 = xx_2 \in L$, conflicting $X_1 \approx X_2$. It follows that V is a plane. Since $V \cap X_i = xx_i \in L$, we have $m_i = m(x_i, V^\perp)$ so that $p(x_1, x_2, V^\perp) \in m_1 \cap m_2$, whence $m_1 \sim m_2$.

Next, define $\sigma: (P', L') \rightarrow (P', L')$ to be the unique map such that $\mu^{-1}(\mu(m)) = \{m, m^\sigma\}$ for each $m \in P'$. Clearly, σ is an involution. Also, σ is an automorphism of (P', L') . For if $m \sim n$ for $m, n \in P'$, then $\mu(m) \sim \mu(n)$ and $m \neq n^\sigma$ since μ is bijective on m^\perp . But then $m^\sigma \sim n^\sigma$ since μ is bijective on $(n^\sigma)^\perp$. So indeed, σ is an automorphism of (P', L') and $d(m, m^\sigma) \geq 3$ for any $m \in P'$.

But if there is $m \in P'$ with $d(m, m^\sigma) = 3$, then there are $m_1, m_2 \in P'$ with $m \sim m_1 \sim m_2 \sim m^\sigma$, so that $m \sim m_2^\sigma \sim m_1^\sigma \sim m^\sigma$. Since μ is an isomorphism on the subgraph induced on m^\perp , and $\{\mu(m_1), \mu(m_2), \mu(m)\}$ is a clique, this

yields that $m_1 \sim m_2^\sigma$. This is in contradiction with $m_1 \sim m_2$.

We have shown that $d(m, m^\sigma) \geq 4$ for any $m \in P'$. Suppose $d(m, m^\sigma) = 4$ for some $m \in P'$. Then there is a minimal path $m \sim m_1 \sim m_2 \sim m_3 \sim m^\sigma$ with $m_i \in P'$ ($i = 1, 2, 3$), so that $m_1 \in m^\perp \cap m_2^\perp$ and $m_3^\sigma \in m^\perp \cap (m_2^\sigma)^\perp$. This leads to two connected components $\mu(m^\perp \cap m_2^\perp)$ and $\mu(m^\perp \cap (m_2^\sigma)^\perp)$ in $\mu(m)^\perp \cap \mu(m_2)^\perp$. Indeed, if there are $n_1 \in m^\perp \cap m_2^\perp$ and $n_2 \in m^\perp \cap (m_2^\sigma)^\perp$ with $\mu(n_1) \sim \mu(n_2)$, then $n_1 \sim n_2$ as $n_1, n_2 \in m^\perp$, so $m_2 \sim n_1 \sim n_2 \sim m_2^\sigma$ is a path of length 3 contradicting $d(m_2, m_2^\sigma) \geq 4$. But this contradicts the fact that $\mu(m)^\perp \cap \mu(m_2)^\perp$ is connected (as it is a generalized quadrangle by assumption). We conclude that $d(m, m^\sigma) \geq 5$ for all $m \in P'$. Finally, since $(P', L') \cong A_{2d-1, d}(F)$ has diameter d , the existence of σ implies that $d \geq 5$. This ends the proof of the theorem. \square

REMARK 6.10. The converse of 6.9 also holds: if (P, L) is as described in (i), (ii) or (iii) of the theorem, then (P, L) is a connected Grassmann space with thick lines whose max spaces have finite ranks.

In case (iii), σ is induced by a polarity of the projective space over F of rank $2d-1$ such that $x \cap x^\sigma$ has codimension at least 5 in x for any subspace x of rank $d-1$. By the classification of such polarities, cf. [3], it follows that F must be infinite.

The above remarks put together with 6.9, prove the main theorem stated in Section 2.

We conclude this section by mentioning that $A_{2d-1, d}(\mathbb{R}) / \langle \sigma \rangle$ for $d \geq 5$, where σ is the polarity associated with the quadratic form $\sum_{i=1}^{2d} x_i^2$ (or any other nondegenerate form of Witt index at most $d-5$), provides an example of a Grassmann space of the type occurring in (iii) of the main theorem.

7. APPLICATIONS.

In this section (P, L) is a connected Grassmann space with thick lines. Consider the following two axioms, each of them stronger than (P4).

- (Q4) If $x \in P$ and $L \in L$ with $x^\perp \cap L = \emptyset$, then $x^\perp \cap L^\perp \in L$.
- (R4) If $L_1, L_2 \in L$ with $L_1 \cap L_2 \neq \emptyset$ and $z \in P$, then there is $u \in z^\perp$ with $u^\perp \cap L_1 \neq \emptyset$ and $u^\perp \cap L_2 \neq \emptyset$.

It is an easy exercise to show that (Q4) holds for (P, L) iff (Q4)' holds, where

(Q4)' If $S \in \mathcal{S}$ and $x \in P \setminus S$, then $x^\perp \cap S$ is either empty or a maximal clique in S .

Also, (R4) is easily shown to be equivalent to (R4)':

(R4)' If $S \in \mathcal{S}$ and $x \in P \setminus S$, then $x^\perp \cap S$ is either a singleton or a maximal clique in S .

We note that (P, L) has diameter 2 if (Q4) holds and diameter at most 3 if (R4) holds.

LEMMA 7.1. *Suppose (P, L) satisfies (Q4). Let S, T be distinct symps on ∞ . If $S \cap T \supseteq \{\infty\}$, then $S \cap T \in \mathcal{V}$.*

PROOF. Consider the residue of ∞ . Suppose $x \in S^\infty \cap T^\infty$. Take $y \in T^\infty \setminus x^\perp$. Note that $y \notin S^\infty$ as $S^\infty \cap T^\infty$ is a clique. Since $L = y^\perp \cap S^\infty$ must be a line in $S^\infty \setminus \{x\}$, there is $z \in x^\perp \cap L \setminus \{x\}$. This implies $z \in x^\perp \cap y^\perp \subseteq T^\infty$, so that $xz \subseteq S \cap T$. \square

THEOREM 7.2. *If (P, L) is a connected Grassmann space with thick lines whose max spaces have finite ranks and in which (Q4) holds, then (P, L) is either a polar space or rank 3 or isomorphic to $A_{a,2}(F)$ for some $a \geq 4$ and some division ring F .*

PROOF. Suppose $M_1, M_2 \in \mathcal{M}$ have rank > 2 and $M_1 \cap M_2 \in L$. In order to apply Proposition 5.9, we verify that $L = M_1 \cap M_2$ is in at least three max spaces. The hypotheses on the ranks of M_1, M_2 imply the existence of points $x_1, x_2 \in M_1 \setminus M_2$, and $y_1, y_2 \in M_2 \setminus M_1$ such that $\text{rk}(\langle x_1, x_2, L \rangle) = \text{rk}(\langle y_1, y_2, L \rangle) = 3$. Clearly $x_i \notin y_i^\perp$. Consider $S_i = S(x_i, y_i)$ for $i = 1, 2$. As $M_1 \cap M_2 \in L(S_1 \cap S_2)$, Lemma 7.1 yields that $S_1 \cap S_2 \in \mathcal{V}$. Thus, if $S_1 \cap S_2 \subseteq M_1$, then $S_1 \cap S_2 = S_1 \cap S_2 \cap M_1 = S_1 \cap M_1 = S_2 \cap M_2$ by consideration of ranks, so $\langle x_1, x_2, L \rangle \subseteq S_1 \cap S_2$ and $3 = \text{rk}(\langle x_1, x_2, L \rangle) \leq \text{rk}(S_1 \cap S_2) = 2$, a contradiction. Hence $S_1 \cap S_2 \not\subseteq M_1$. Similarly, one can prove $S_1 \cap S_2 \not\subseteq M_2$. Now $(S_1 \cap S_2)^\perp$ is a third max space on L , and we can finish by Proposition 5.9. \square

FROM NOW ON WE ASSUME THAT (R4) HOLDS FOR (P, L)

LEMMA 7.3. Let x_1, x_2, x_3, x_4, x_5 be a minimal 5-circuit (i.e. $x_i^\perp \cap x_{i+2}x_{i+3} \neq \emptyset$ for all i , indices taken modulo 5). If $x_1^\perp \cap S(x_2, x_4) \in V$, then $x_i^\perp \cap S(x_{i+1}, x_{i+3})$ and $x_i^\perp \cap S(x_{i-1}, x_{i-3})$ are in V for all $i(1 \leq i \leq 5)$.

PROOF. Note that $x_1^\perp \cap S(x_2, x_4) \in V$ iff $\{x_1, x_2, x_3, x_4\}^\perp \neq \emptyset$. Thus $x_4^\perp \cap S(x_1, x_3) \in V$ follows. Also for $u \in \{x_1, x_2, x_3, x_4\}^\perp$, we have $ux_4 \subseteq x_3^\perp \cap S(x_1, x_4)$, so $x_3^\perp \cap S(x_1, x_4) \in V$. Similarly $x_2^\perp \cap S(x_1, x_4) \in V$. The argument is easily completed. \square

COROLLARY 7.4. Let $L \in L$ and $S \in S$ with $S \cap L = \emptyset$. If $x, y \in L$ and $x_1, y_1 \in S$ with $x^\perp \cap S = \{x_1\}$ and $y^\perp \cap L = \{y_1\}$ then $x_1 \in y_1^\perp$.

PROOF. We may assume $x_1 \neq y_1$, for else there is nothing to prove. Take $u \in x_1^\perp \cap y_1^\perp \setminus \{x_1, y_1\}$, and consider the 5-circuit u, x_1, x, y, y_1 . Since $x \notin y_1^\perp \cup u^\perp$ and $y \notin x_1^\perp \cup u^\perp$ and $x^\perp \cap S(x_1, y_1) = \{x_1\}$, the lemma implies that $x_1 \in y_1^\perp$. \square

LEMMA 7.5. Suppose $x_1 \sim x_2 \sim x_3 \sim x_4$ is a path in (P, L) with $x_1 \notin x_3^\perp$ and $x_2 \notin x_4^\perp$ such that $\{x_1, x_2, x_3, x_4\}^\perp = \emptyset$. Then for $M \in M$ on x_1x_2 , there is a unique max space M_1 on x_3x_4 of distance 2 to M in (M, \sim) .

PROOF. By Lemma 3.8 there is a (unique) max space M' on x_2x_3 collinear with M . Similarly there is a unique max space M_1 on x_3x_4 collinear with M' . Now $M_1 \cap M = \emptyset$ as $M_1 \cap M \subseteq \{x_1, x_2, x_3, x_4\}^\perp$, so M_1 has distance 2 to M in (M, \sim) . Suppose M_2 is also a max space on x_3x_4 of distance 2 to M . Then by 6.4 there are max spaces N_1, N_2 on x_2x_3 with $N_i \cap M, N_i \cap M_1 \in L$ for each $i \in \{1, 2\}$. But N_1, N_2, M are three max spaces on x_2 the intersection of any two of which contains a line. This implies $N_1 = N_2$. But then N_1, M_1, M_2 are three max spaces on x_3 , the intersection of any two of which contains a line. It results that $M_1 = M_2$. \square

LEMMA 7.6. Suppose (P, L) satisfies (R4). If x_1, x_2, \dots, x_5 is a minimal 5-circuit in P , then

- (i) $x_i^\perp \cap S(x_{i+1}, x_{i+3}) \in V$ for each i ($1 \leq i \leq 5$), indices taken modulo 5).
- (ii) $\{x_1, x_2, \dots, x_5\}^\perp = \emptyset$.

PROOF. (i) Suppose x_1, x_2, \dots, x_5 is a minimal 5-circuit which is a counter-example to the statement. By 7.3, it is a circuit with $x_i^\perp \cap S(x_{i+1}, x_{i+3})$ a singleton for each i . Let M be a max space on x_3x_4 and take $M_1 \in M$ on x_2x_3 with $M_1 \cap M = \{x_3\}$ and $M_2 \in M$ on x_4x_5 with $M_2 \cap M \in L$. Now $L_1 = x_1^\perp \cap M_1$, $L_2 = x_1^\perp \cap M_2$, $L_3 = x_3^\perp \cap M_2 = M \cap M_2$, $L_4 = x_4^\perp \cap M_2$ are lines on x_2, x_5, x_4, x_3 respectively.

Since $L_1 \cap L_4 \subseteq \{x_1, x_2, x_3, x_4\}^\perp$ and $L_2 \cap L_3 \subseteq \{x_1, x_5, x_4, x_3\}^\perp$ we have by the assumption that $L_1 \cap L_4 = L_2 \cap L_3 = \emptyset$. Take $u \in L_3 \setminus \{x_4\}$ and $v \in L_4 \setminus \{x_3\}$. Then $u \notin v^\perp$. For $u \in v^\perp$ would imply $L_3 \subseteq L_4^\perp$ and $\langle L_3, L_4 \rangle^\perp = M$ so that $M \cap M_1$ would contain the line L_4 , conflicting $M \cap M_1 = \{x_3\}$.

Consider $S = S(u, v)$. Note that $V = x_5^\perp \cap S$ contains L_3 and must therefore be a plane in S . Similarly for $W = x_2^\perp \cap S$. Note that $x_1 \notin S$, for else $x_1^\perp \cap x_3x_4 \neq \emptyset$.

Now $x_1^\perp \cap S \neq \emptyset$ by (R4). As $x_1 \in x_2^\perp \cup x_5^\perp$, Lemma 3.5 implies that $x_1^\perp \cap x_2^\perp \cap S$ and $x_1^\perp \cap x_5^\perp \cap S$ are nonempty. If $z \in \{x_1, x_2, x_5\}^\perp \cap S$, then $z \in \{x_1, x_2, x_3, x_4, x_5\}^\perp \cap S$, as $x_5^\perp \cap S$ is a clique on x_4 and $x_2^\perp \cap S$ is a clique on x_3 . So we may assume $\{x_1, x_2, x_5\}^\perp \cap S = \emptyset$. Thus $|x_1^\perp \cap S| \geq |x_1^\perp \cap x_2^\perp \cap S| + |x_1^\perp \cap x_5^\perp \cap S| \geq 2$, so that $x_1^\perp \cap S \in \mathcal{V}$. Write $U = x_1^\perp \cap S$. Since U, x_3, x_4 are in S , there is $w \in x_3^\perp \cap x_4^\perp \cap U$. But now zx_3 is a line in $x_4^\perp \cap S(x_1, x_3)$; this settles (i).

(ii) Assume $u \in \{x_1, x_2, \dots, x_5\}^\perp$. Put $L = x_1^\perp \cap (x_3x_4)^\perp$. Since $x_1^\perp \cap x_3x_4 = \emptyset$ by minimality of the circuit, $L \in L$. Now $\langle x_1, x_5, u \rangle^\perp, \langle x_1, x_2, u \rangle^\perp \in M_{xu}$, so $\langle x_1, L \rangle \leq \langle x_1, x_5, u \rangle^\perp$ or $\langle x_1, L \rangle \subseteq \langle x_1, x_2, u \rangle^\perp$.

Without loss of generality, assume $\langle x_1, L \rangle \subseteq \langle x_1, x_5, u \rangle^\perp$. Then $\langle x_4, L \rangle \subseteq x_3^\perp \cap x_5^\perp$ conflicting ranks. \square

LEMMA 7.7. *Let (P, L) satisfy (R4). If S, T are distinct symps, then $S \cap T$ is not a singleton.*

PROOF. Suppose S, T are symps such that $S \cap T = \{x\}$ for some $x \in P$. Take $z \in S \setminus x^\perp$. By axiom (R4), there is $y \in z^\perp \cap T$. Now $y \in T \setminus x^\perp$, for else $y \in x^\perp \cap z^\perp \subseteq S$, so $y \in S \cap T = \{x\}$ and $y = x$ conflicting $z \notin x^\perp$. Choose $v_1, v_2 \in x^\perp \cap y^\perp$ with $v_1 \notin v_2^\perp$, and take $u \in x^\perp \cap z^\perp$.

Let $i \in \{1,2\}$. Now u, x, v_i, y, z is a 5-circuit with $u \notin y^\perp$ (for else $u \in x^\perp \cap y^\perp \subseteq T$), $x \notin z^\perp \cup y^\perp$ and $v_i \notin z^\perp$ (for otherwise $v_i \in x^\perp \cap z^\perp \subseteq S$). So either $u \in v_i^\perp$ and $v_i^\perp \cap S \supseteq xu$, or $v_i^\perp \cap S \in V$. At any rate, $v_i^\perp \cap S \in V$ for each $i \in \{1,2\}$. Put $V_i = v_i^\perp \cap S$ and consider $W = y^\perp \cap S$. As $z \in W \setminus (V_1 \cup V_2)$, we must have $W \in V$ by Lemma 3.5. But then $x^\perp \cap W$ is a line (as both x, W are in S) contained in $x^\perp \cap y^\perp$, hence in T . \square

LEMMA 7.8. *Suppose (R4) holds for (P, L) . Then $\text{rk}(M) \leq 3$ for any $M \in \mathcal{M}$.*

PROOF. Suppose M is a singular subspace of rank 4. Pick $x \in M$ and $V, W \in V(M)$ with $V \cap W = \{x\}$, and let S, T be symps on V, W respectively. Since $x \in S \cap T$, we know by Lemma 7.6 that there is a line L on x in $S \cap T$. Now $V \subseteq L^\perp$ would imply $L \subseteq V^\perp \cap T = W$; but also $L \subseteq V$, as $\langle V, L \rangle$ is a singular subspace of S , so that $L \subseteq S \cap T = \{x\}$ which is absurd. Hence there is $z \in L \setminus \{x\}$ with $z \notin V^\perp$. Since z, V are in S , we obtain that $L_1 = z^\perp \cap V$ is a line on x . Similarly, $L_2 = z^\perp \cap W$ is a line on x . But now $z \in L_1^\perp \cap L_2^\perp = \langle L_1, L_2 \rangle^\perp = M^\perp$, so $L \subseteq M$ and $V \subseteq L^\perp$, which has just been excluded.

It follows that no max space of rank 4 exists. \square

THEOREM 7.9. *If (P, L) is a connected Grassmann space with thick lines in which (R4) holds then (P, L) is either a polar space of rank 3 or isomorphic to one of $A_{4,2}(F), A_{5,3}(F)$ for some division ring F .*

PROOF. This is a direct consequence of 6.9 and 7.8. \square

ACKNOWLEDGEMENTS

The problem dealt with in this paper was raised in a discussion with W.M. Kantor. Both his and E.E. Shult's interest were a great support. Also, H.A. Wilbrink's contributions to the proof of a property on generalized quadrangles (Proposition 4.2) are gratefully acknowledged.

REFERENCES

- [1] BUEKENHOUT, F., *Cooperstein's theory*, preprint.
- [2] COOPERSTEIN, B.N., *A characterization of some Lie incidence structures*, *Geometriae Dedicata*, 6 (1977) 205-258.
- [3] DIEUDONNÉ, J., *La géométrie des groupes classiques*, Springer, Berlin, 1963.
- [4] HIGMAN, D.G., *Invariant relations, coherent configurations and generalized polygons*, pp. 27-43 in 'Combinatorics, Part 3: Combinatorial group theory', Math Centre Tract 57, Amsterdam, 1974.
- [5] SPRAGUE, A., *Rasch's axiom and projective spaces*, *Discrete Mathematics* 33 (1981), 79-87.
- [6] TALLINI, G., *On a characterization of the Grassmann manifold representing the lines in a projective space*, pp. 354-358 in 'Finite Geometries and Designs', Proceedings of the 2nd Isle of Thorns Conference 1980, LMS Lecture Note 49, Cambridge University Press 1981.
- [7] TITS, J., *Buildings of spherical type and finite BN pairs*, Springer LNM 386, Berlin, 1974.
- [8] TITS, J., *A local approach to buildings*, preprint.

ONTVANGEN D 5 OKT. 1981