

**stichting
mathematisch
centrum**



AFDELING ZUIVERE WISKUNDE
(DEPARTMENT OF PURE MATHEMATICS)

ZW 130/79

NOVEMBER

R. TIJDEMAN & M. VOORHOEVE

BOUNDED DISCREPANCY SETS

Preprint

2e boerhaavestraat 49 amsterdam

Printed at the Mathematical Centre, 49, 2e Boerhaavestraat, Amsterdam.

The Mathematical Centre, founded the 11-th of February 1946, is a non-profit institution aiming at the promotion of pure mathematics and its applications. It is sponsored by the Netherlands Government through the Netherlands Organization for the Advancement of Pure Research (Z.W.O).

Bounded discrepancy sets^{*)}

by

R. Tijdeman & M. Voorhoeve

ABSTRACT

Let $\omega = \{\xi_j\}_{j=1}^{\infty}$ be a sequence in $[0,1)$. We define the discrepancy function D_n by $D_n(\omega, \alpha) = Z_n(\omega, \alpha) - n\alpha$, where $Z_n(\omega, \alpha)$ is the number of elements in $[0, \alpha)$ among the first n terms of ω . It is known that $\sup_{\alpha, n} D_n(\omega, \alpha) = \infty$ for every sequence ω . In this paper sets S are characterized for which an ω exists such that $\sup_n D_n(\omega, \alpha) < \infty$ for every $\alpha \in S$. Furthermore we investigate sets S such that $\sup_{\alpha \in S, n \in \mathbb{N}} D_n(\omega, \alpha) < \infty$ for some ω . In particular, we show in Corollary 1 of Theorem 5 that such sets S have relatively large gaps. Theorems 1-4 are based on Lemma 1, which provides a construction for sequences with small discrepancy at specific points. Theorems 5 and 6 are applications of Lemma 3 which is proved by a method of W.H. Schmidt.

KEY WORDS & PHRASES: *Discrepancy, irregularities of distribution, uniform distribution.*

^{*)} This report will be submitted for publication elsewhere.

1. INTRODUCTION

Let U be the unit interval consisting of numbers ξ with $0 \leq \xi < 1$, and let $\omega = \{\xi_1, \xi_2, \dots\}$ be a sequence of numbers in this interval. Given an α in U and a positive integer n , we write $Z_n(\omega, \alpha)$ for the number of integers i with $1 \leq i \leq n$ and $0 \leq \xi_i < \alpha$ and we put $D_n(\omega, \alpha) = Z_n(\omega, \alpha) - n\alpha$. For convenience we define $D_n(\omega, 1) = 0$ and $D_0(\omega, \alpha) = 0$ for all α , n and ω . Put $D(\omega, \alpha) = \sup_n |D_n(\omega, \alpha)|$.

In answering a question of J.G. van der CORPUT [2], Mrs. T. van AARDENNE-EHRENFEST [1] showed that there is no sequence ω in U for which $\sup_{\alpha \in U} D(\omega, \alpha)$ is bounded. P. ERDŐS [3] wondered whether for every sequence ω there exist numbers α such that $D(\omega, \alpha) = \infty$. This was answered by W.M. SCHMIDT [4] in the affirmative. Later SCHMIDT [7, p.40] proved that for every sequence ω even

$$\limsup_{n \rightarrow \infty} \frac{|D_n(\omega, \alpha)|}{\log \log n} > \frac{1}{2000}$$

for almost all α . SCHMIDT [5] also investigated sets at which D can remain bounded. He demonstrated that the set $S(\infty) := \{\alpha : D(\omega, \alpha) < \infty\}$ is countable for every sequence ω . Theorem 1 gives the opposite result that for every countable subset S of U there exists a sequence ω such that $D(\omega, \alpha) < \infty$ for every α in S . In the special case $S = \emptyset$ Theorem 3 gives a quantitative result which is in a sense the best possible. We remark that SCHMIDT [6] generalized his result on the countability of $S(\infty)$ in a very remarkable manner. See also L. SHAPIRO [8].

We call S a κ -discrepancy set if there exists a sequence ω such that $D(\omega, \alpha) < \kappa$ for every α in S . A bounded discrepancy set (BDS) is a set which is a κ -discrepancy set for some κ . Theorem 2 states that every finite set is a BDS. Recall that a number γ is a limit point of a set S if there is a sequence of distinct elements of S which converges to γ . The derivative $S^{(1)}$ of S consists of all the limit points of S . The higher derivatives are defined inductively by $S^{(d)} = (S^{(d-1)})^{(1)}$ ($d = 2, 3, \dots$). SCHMIDT [5] proved that $S^{(d)}$ is empty if S is a κ -discrepancy set and if $d > 4\kappa$. Furthermore he showed that $S^{(d)}$ need not be empty if S is a d -discrepancy set. This provides a necessary condition for being a BDS. The fact that $S = \{n^{-1}\}_{n=2}^{\infty}$ is not a BDS while $S^{(2)} = \emptyset$ shows that the condition is not sufficient. The

corollary of Theorem 5 gives a property of a BDS which this set does not fulfill: if S is a BDS then there is an $\varepsilon > 0$ such that every interval of length ℓ contains a subinterval J of length $\varepsilon\ell$ with $J \cap S = \emptyset$. It seems a difficult problem to characterize BDS' in a simple way, if possible at all. In Section 4 we argue that the essential problem already occurs for a monotonic decreasing sequence with limit 0. Theorem 4 gives a sufficient condition for being a BDS and in Theorem 6 we show that in a certain case the necessary and sufficient conditions coincide.

2. The basic tool for construction BDS' is the following lemma.

LEMMA 1. *Let α, β, γ be real numbers with $0 \leq \alpha < \beta < \gamma \leq 1$. Let $V \subseteq U$. Assume there is a sequence $\omega = \{\xi_n\}_{n=1}^{\infty}$ in V such that $D(\omega, \alpha) \leq A$ and $D(\omega, \gamma) \leq C$. Then there exists a sequence $\omega' = \{\xi'_n\}_{n=1}^{\infty}$ in $V \cup \{\alpha\} \cup \{\beta\}$ such that*

- (i) $\xi'_n = \xi_n$ if $\xi_n \in [0, \alpha) \cup [\gamma, 1)$,
- (ii) $\xi'_n \in \{\alpha, \beta\}$ if $\xi_n \in [\alpha, \gamma)$,
- (iii) $D(\omega', x) = D(\omega, x)$ for $x \in [0, \alpha] \cup [\gamma, 1)$,
- (iv) $D(\omega', \beta) \leq \frac{\gamma-\beta}{\gamma-\alpha} A + \frac{\beta-\alpha}{\gamma-\alpha} C + \frac{1}{2}$.

PROOF. We may assume without loss of generality that $\xi_n = \alpha$ if $\xi_n \in [\alpha, \gamma)$, since $D(\omega, x)$ for $x \in (\alpha, \gamma)$ is of no importance for the lemma. We shall prove by induction on m that we can define $\xi'_m \in \{\alpha, \beta\}$ in such a way that

$$(1) \quad -\frac{1}{2} \leq \Delta_m \leq \frac{1}{2}$$

where

$$(2) \quad \Delta_m = D_m(\omega', \beta) - \frac{\gamma-\beta}{\gamma-\alpha} D_m(\omega, \alpha) - \frac{\beta-\alpha}{\gamma-\alpha} D_m(\omega, \gamma).$$

It is obvious that $\Delta_0 = 0$ and that (1) holds for $m = 0$. Suppose that m is some non-negative integer for which the induction hypothesis holds. If $\xi_{m+1} \in [0, \alpha) \cup [\gamma, 1)$, then we put $\xi'_{m+1} = \xi_{m+1}$. It follows that

$$\Delta_{m+1} = \Delta_m + (1-\beta) - \frac{\gamma-\beta}{\gamma-\alpha} (1-\alpha) - \frac{\beta-\alpha}{\gamma-\alpha} (1-\gamma) = \Delta_m$$

if $\xi_{m+1} \in [0, \alpha)$ and that

$$\Delta_{m+1} = \Delta_m - \beta + \frac{\gamma - \beta}{\gamma - \alpha} \alpha + \frac{\beta - \alpha}{\gamma - \alpha} \gamma = \Delta_m$$

if $\xi_{m+1} \in [\gamma, 1)$. Hence (1) holds in this case. If $\xi_{m+1} = \alpha$ then put $\xi'_{m+1} = \alpha$ if $\Delta_m \leq (\beta - \alpha)/(\gamma - \alpha) - \frac{1}{2}$ and $\xi'_{m+1} = \beta$ otherwise. If $\xi'_{m+1} = \alpha$, then

$$\Delta_{m+1} = \Delta_m + (1 - \beta) + \frac{\gamma - \beta}{\gamma - \alpha} \alpha - \frac{\beta - \alpha}{\gamma - \alpha} (1 - \gamma) = \Delta_m + 1 - \frac{\beta - \alpha}{\gamma - \alpha}$$

and hence, by (1), $-\frac{1}{2} \leq \Delta_{m+1} \leq \frac{1}{2}$. If $\xi'_{m+1} = \beta$, then

$$\Delta_{m+1} = \Delta_m - \frac{\beta - \alpha}{\gamma - \alpha}$$

and hence, by (1), $-\frac{1}{2} \leq \Delta_{m+1} \leq \frac{1}{2}$. Thus (1) is valid with $m+1$ in place of m .

By the above construction a sequence $\omega' = \{\xi'_n\}_{n=1}^{\infty}$ is defined which satisfies (i) and (ii). Further (iii) is an immediate consequence of (i) and (ii). Finally it follows from (1) and (2) that

$$|D_m(\omega', \beta)| \leq \frac{\gamma - \beta}{\gamma - \alpha} |D_m(\omega, \alpha)| + \frac{\beta - \alpha}{\gamma - \alpha} |D_m(\omega, \gamma)| + \frac{1}{2}$$

for $m = 1, 2, \dots$. This implies (iv).

REMARK. Note that the discrepancy of ω' is bounded in both α and β and γ . Hence ω' assumes both values in $[\alpha, \beta)$ and in $[\beta, \gamma)$. By (i) and (ii) this implies that both α and β occur as terms of ω' .

3. SCHMIDT [5] proved that every $S(\infty)$ set is countable. The following theorem shows that every countable set is a $S(\infty)$ -set.

THEOREM 1. For every countable set $S = \{\alpha_1, \alpha_2, \dots\}$ in U there exists a sequence ω such that $D(\omega, \alpha_j) < \infty$ for $j = 1, 2, \dots$.

PROOF. Without loss of generality we may assume that $0, \alpha_1, \alpha_2, \dots$ are distinct numbers. We shall prove by induction on m that there exists a sequence $\omega_m = \{\xi_{m,1}, \xi_{m,2}, \dots\}$ in $\{0, \alpha_1, \dots, \alpha_m\}$ such that

- (i) $D(\omega_m, \alpha_j) = D(\omega_{m-1}, \alpha_j)$ for $j = 1, 2, \dots, m-1$,
- (ii) $D(\omega_m, \alpha_j) < \infty$ for $j = 1, 2, \dots, m$,
- (iii) If $1 \leq j < m$ and $\xi_{m-1,n}$ is the first element of ω_{m-1} with $\xi_{m-1,n} = \alpha_j$, then $\xi_{m,n} = \alpha_j$.

For $m = 1$ we apply Lemma 1 with $\alpha = 0$, $\beta = \alpha_1$, $\gamma = 1$, $A = C = 0$, $V = \{0\}$. Suppose that m is a non-negative integer for which the induction hypothesis holds. Let α be the largest element of the set $\{0, 1, \alpha_1, \dots, \alpha_m\}$ which is smaller than α_{m+1} and let γ be the smallest element of this set which is larger than α_{m+1} . Apply Lemma 1 with this α and γ and with $\beta = \alpha_{m+1}$. This gives a sequence ω'_{m+1} in $\{0, \alpha_1, \dots, \alpha_{m+1}\}$ satisfying (i) and (ii). Let n be the smallest integer with $\xi_{m,n} = \alpha$. If $\xi'_{m+1,n} = \alpha$, then put $\omega_{m+1} = \omega'_{m+1}$. If $\xi'_{m+1,n} = \beta$, then we form ω_{m+1} by interchanging the first α and the first β in ω'_{m+1} . This change does only affect the discrepancy in $(\alpha, \beta]$, in fact by at most 1 in absolute value. Since ω_{m+1} is derived from ω_m by merely replacing some α 's by β 's, the other α_j 's in ω_m remain unaltered. Thus ω_{m+1} satisfies (i) - (iii) and the induction step is complete.

By (iii) the sequence $\{\xi_{m,n}\}_{m=1}^{\infty}$ is constant from some $m_0 = m_0(n)$ on. Put $\xi_n = \xi_{m_0,n}$. This induces a sequence $\omega = \{\xi_1, \xi_2, \dots\}$. By the construction $\xi_n < \alpha_j$ if $\xi_{j,n} < \alpha_j$ and $\xi_n \geq \alpha_j$ if $\xi_{j,n} \geq \alpha_j$, for all j and n . Hence $D(\omega, \alpha_j) = D(\omega_j, \alpha_j) < \infty$ for $j = 1, 2, \dots$.

The following theorem shows that every finite set is a BDS and gives an upper bound which can only be improved by a constant factor in view of Corollary 2.

THEOREM 2. *For every finite set $S = \{\alpha_1, \alpha_2, \dots, \alpha_m\}$ in U there exists a sequence ω such that*

$$D(\omega, \alpha_j) \leq \frac{\log(2m)}{2 \log 2} \quad \text{for } j = 1, 2, \dots, m.$$

PROOF. We prove by induction on t that for every finite set $\{\alpha_1, \alpha_2, \dots, \alpha_{2^{t-1}}\}$ in U there exists a sequence ω_t such that $D(\omega_t, \alpha_j) \leq t/2$ for $j = 1, 2, \dots, 2^{t-1}$. For $t = 1$ we apply Lemma 1 with $\alpha = 0$, $\beta = \alpha_1$, $\gamma = 1$, $A = C = 0$. Suppose the induction hypothesis is true for t . Let $\{\alpha_1, \alpha_2, \dots, \alpha_{2^{t+1}-1}\} \subset U$. We may assume without loss of generality that $0 < \alpha_1 < \alpha_2 < \dots < \alpha_{2^{t+1}-1}$. Put

$\alpha_0 = 0$. There exists a sequence ω_t' in $\{\alpha_0, \alpha_2, \alpha_4, \dots, \alpha_{2^{t+1}-2}\}$ such that $D(\omega_t', \alpha_{2^i}) \leq t/2$ for $i = 0, 1, \dots, 2^t - 1$. On applying Lemma 1 with $\alpha = \alpha_{2^i}$, $\beta = \alpha_{2^{i+1}}$, $\gamma = \alpha_{2^{i+2}}$, $A = C = t/2$ for $i = 0, 1, \dots, 2^t - 1$ and combining the resulting sequences in an obvious way, we obtain a sequence ω_{t+1} such that $D(\omega_{t+1}, \alpha_i) \leq (t+1)/2$ for $i = 0, 1, \dots, 2^{t+1} - 1$. This proves the induction hypothesis for all values of t .

Let a set $S = \{\alpha_1, \alpha_2, \dots, \alpha_m\}$ be given. Let t be the integer with $2^{t-1} \leq m < 2^t$. We have shown that there exists a sequence $\omega = \omega_t$ with

$$D(\omega, \alpha_j) \leq \frac{1}{2} t < \frac{1}{2} (1 + \frac{\log m}{\log 2}) \quad \text{for } j = 1, 2, \dots, m.$$

The following result gives a quantitative form of Theorem 1 in the special case $S = \mathbb{Q}$ which is best possible in a similar way as Theorem 2 is.

THEOREM 3. *There exists a sequence ω such that*

$$D(\omega, \frac{p}{q}) \leq 1 + 4 \log q$$

for every p/q with $p, q \in \mathbb{Z}$ and $0 < p < q$.

PROOF. We prove by induction on t that there exists a sequence $\omega_t = \{\xi_{t,n}\}_{n=1}^{\infty}$ in a finite set V_t of at most 2^{3t} rational numbers with the following properties:

- (i) $V_{t-1} \subset V_t$ for $t \geq 2$,
- (ii) V_t contains all numbers $p2^{-2t}$ with $p \in \mathbb{Z}$ and $0 \leq p < 2^{2t}$,
- (iii) V_t contains all numbers pq^{-1} with $p, q \in \mathbb{Z}$ and $0 < p < q \leq 2^t$,
- (iv) if $\alpha \in V_{t-1}$ and $\xi_{t-1,n}$ is the first element of ω_{t-1} with $\xi_{t-1,n} = \alpha$, then $\xi_{t,n} = \alpha$,
- (v) $D(\omega_t, \alpha) \leq \frac{5}{2} t - \frac{3}{2}$ for every α in V_t .

For $t = 1$ we take $V_1 = \{0, \frac{1}{4}, \frac{1}{2}, \frac{3}{4}\}$ and by a double application of Lemma 1 there exists a sequence ω_1 in V_1 such that $D(\omega_1, \alpha) \leq 1$ for $\alpha \in V_1$. Suppose t is a positive integer for which the induction hypothesis is true. We construct V_{t+1} in three steps:

$$V'_t = V_t \cup \left\{ \frac{k}{2^{2t+1}} : k \in \mathbb{Z}, 0 < k < 2^{2t+1} \right\},$$

$$V''_t = V'_t \cup \left\{ \frac{k}{2^{2t+2}} : k \in \mathbb{Z}, 0 < k < 2^{2t+2} \right\},$$

$$V_{t+1} = V''_t \cup \left\{ \frac{p}{q} : p, q \in \mathbb{Z}, 0 < p < q \leq 2^{t+1} \right\}.$$

Observe that at each step any two "new" points are separated by an "old" point. Hence we can apply Lemma 1 as we did in the proof of Theorem 2 and we obtain sequences $\omega'_t, \omega''_t, \omega'''_t$ with discrepancy at V'_t, V''_t, V_{t+1} at most $\frac{5}{2}t - 1, \frac{5}{2}t - \frac{1}{2}, \frac{5}{2}t$ respectively. Clearly (i) - (iii) are fulfilled with $t+1$ in place of t . For every $\alpha \in V_t$ with the property that $\xi_{t+1,n} \neq \alpha$ where n is the smallest integer with $\xi_{t,n} = \alpha$ we make an interchange like in the proof of Theorem 1. In such a case $\xi_{t+1,n}$ is a number $\beta \in V_{t+1} \setminus V_t$ which is smaller than the smallest element of V_t which is larger than α . By interchanging the first α and the first β in ω'''_t the discrepancy function remains unchanged outside the interval $(\alpha, \beta]$ and changes by at most 1 in $(\alpha, \beta]$ in absolute value. Since these intervals $(\alpha, \beta]$ are disjoint, the sequence ω_{t+1} which results after all interchanges have been made, satisfies (iv) with $t+1$ in place of t and moreover $D(\omega_{t+1}, \alpha) \leq \frac{5}{2}t+1$ for every $\alpha \in V_{t+1}$. This completes the induction step.

By (iv) the sequence $\{\xi_{t,n}\}_{n=1}^{\infty}$ is constant from some $t_0 = t_0(n)$ on. Put $\xi_n = \xi_{t_0,n}$. This induces a sequence $\omega = \{\xi_1, \xi_2, \dots\}$. By the construction $\xi_n < \alpha$ if $\xi_{t,n} < \alpha$ and $\xi_n \geq \alpha$ if $\xi_{t,n} \geq \alpha$ for every α, n and t with $\alpha \in V_t$. Let $p/q \in \mathbb{Z}$ with $0 < p < q \leq 2^t$. Let t be the integer with $2^{t-1} < q \leq 2^t$. Then $p/q \in V_t$. Hence

$$D(\omega, \frac{p}{q}) = D(\omega_t, \frac{p}{q}) \leq \frac{5}{2}t - \frac{3}{2} < 1 + 5 \log q / 2 \log 2 < 1 + 4 \log q.$$

4. Suppose we want to decide whether a set S is a BDS. If it is, there exists a sequence ω and an integer d such that

$$(3) \quad D(\omega, \alpha) \leq d \quad \text{for every } \alpha \in S.$$

It follows from a result of SCHMIDT [5] that S has to be countable and $S^{(4d+1)} = \emptyset$. Note that $D_n(\omega, \alpha) = \lim_{\varepsilon \downarrow 0} D_n(\omega, \alpha + \varepsilon)$ for every α and n . Hence if α_0 is the limit of an increasing sequence in S and S satisfies (3) then $D(\omega, \alpha_0) \leq d$. If $\alpha_0 > 0$ is a limit point of S but not the limit of an increasing sequence in S , then we can replace every α_0 in ω by $\alpha_0 - \varepsilon$ for a sufficiently small $\varepsilon > 0$ without changing $D(\cdot, \alpha)$ for $\alpha \in S \cup S^{(1)} \setminus \{\alpha_0\}$. For this new sequence ω' we have $D(\omega', \alpha) = \lim_{\varepsilon \downarrow 0} D(\omega, \alpha + \varepsilon) \leq d$. Since we can do so for all such $\alpha_0 \in S^{(1)} \setminus S$ simultaneously, we conclude that S is a BDS if and only if $S \cup S^{(1)}$ is a BDS. We may therefore assume without loss of generality that S is closed. It further follows that $S^{(j)}$ ($j = 1, 2, \dots$) as a subsequence of S is also a BDS. So it is sufficient to be able to decide whether a set S is a BDS if it is known that $S^{(1)}$ is a BDS, for then one can apply the argument to make the transitions $S^{(4d+1)} \rightarrow S^{(4d)} \rightarrow \dots \rightarrow S^{(1)} \rightarrow S$.

Let S be a set such that $S^{(1)}$ is a BDS. For $\alpha \in S$ let $\phi(\alpha)$ denote an element in $S^{(1)}$ with $|\alpha - \phi(\alpha)|$ minimal. Let $\beta \in S^{(1)}$ and let $\alpha_1, \alpha_2, \dots$ be all elements of S with $\phi(\alpha_j) = \beta$ and $\alpha_j > \beta$ ordered in such a way that $\alpha_1 > \alpha_2 > \alpha_3 > \dots$. It is obvious that $\alpha_1, \alpha_2, \dots$ is a BDS if and only if $\alpha_1 - \beta, \alpha_2 - \beta, \dots$ is a BDS. For the points $\alpha \in S$ with $\phi(\alpha) = \beta$ and $\alpha < \beta$ a similar argument applies. So the essential difficulty is to decide whether a monotonic sequence $\alpha_1, \alpha_2, \dots$ in U with limit 0 is a κ -discrepancy set or not. If $S^{(1)}$ is a BDS and there exists a constant κ such that for every $\beta \in S^{(1)}$ both the points $\alpha \in S$ with $\phi(\alpha) = \beta$, $\alpha < \beta$ and the points $\alpha \in S$ with $\phi(\alpha) = \beta$, $\alpha > \beta$ are κ -discrepancy sets, then S is a BDS itself.

The following result gives a sufficient condition for a monotonic decreasing sequence with limit 0 to be a BDS. Necessary conditions for such sequences are given in Theorems 5 and 6.

THEOREM 4. Let $\{\alpha_n\}_{n=1}^{\infty}$ be a monotonic decreasing sequence in U with $\alpha_n \rightarrow 0$ as $n \rightarrow \infty$. If there exist a positive integer h and a constant c with $c < 1$ such that $\alpha_{n+h} < c\alpha_n$ for $n = 1, 2, \dots$, then there exists a sequence ω such that

$$D(\omega, \alpha_n) \leq \frac{1}{2-2c} + \frac{\log 2h}{2 \log 2} \quad \text{for } n = 1, 2, \dots$$

PROOF. We prove by induction on t that there exists a sequence

$\omega_t = \{\xi_{t,n}\}_{n=1}^{\infty}$ in $\{0, \alpha_{th}, \alpha_{th-1}, \dots, \alpha_1\}$ such that

$$(4) \quad D(\omega_t, \alpha_j) \leq \frac{1}{2-2c} \quad \text{for } j = 1, 2, \dots, t$$

and

$$(5) \quad D(\omega_t, \alpha_j) \leq \frac{1}{2-2c} + \frac{\log 2h}{2 \log 2} \quad \text{for } j = 1, 2, \dots, th.$$

For $t = 0$ the assertion is true. Suppose t is a non-negative integer for which the induction hypothesis holds. First apply Lemma 1 with $\alpha = 0$, $\beta = \alpha_{(t+1)h}$, $\gamma = \alpha_{th}$ ($\gamma = 1$ if $t = 0$), $A = C = (2-2c)^{-1}$. Hence, there exists a sequence ω'_t in $\{0, \alpha_{(t+1)h}, \alpha_{th}, \alpha_{th-1}, \alpha_{th-2}, \dots, \alpha_1\}$ such that

$$D(\omega'_t, \alpha_j) \leq \frac{c}{2-2c} + \frac{1}{2} = \frac{1}{2-2c} \quad \text{for } j = 1, 2, \dots, t+1$$

and

$$D(\omega'_t, \alpha_j) \leq \frac{1}{2-2c} + \frac{\log 2h}{2 \log 2} \quad \text{for } j = 1, 2, \dots, th.$$

Next we apply the argument used in the proof of Theorem 2 to the points $\alpha_{(t+1)h-1}, \dots, \alpha_{th+1}$. The only difference is that everywhere A and C have to be increased by $(2-2c)^{-1}$. So we obtain a sequence ω_{t+1} in $\{0, \alpha_{(t+1)h}, \alpha_{(t+1)h-1}, \dots, \alpha_1\}$ which satisfies (4) and (5) with $t+1$ instead of t .

Every sequence $\{\xi_{t,n}\}_{t=1}^{\infty}$ is constant from some $t_0 = t_0(n)$ on. Let $\xi_n = \lim_{t \rightarrow \infty} \xi_{t,n}$. This defines the sequence $\omega = \{\xi_n\}_{n=1}^{\infty}$. As before we have

$$D(\omega, \alpha_j) = D(\omega_j, \alpha_j) \leq \frac{1}{2-2c} + \frac{\log 2h}{2 \log 2} \quad \text{for } j = 1, 2, \dots$$

5. To derive further properties of a BDS we use a technique due to SCHMIDT [5]. Since we shall work from now on with one sequence ω only, we shall suppress the variable ω and write $D_n(\alpha)$, etc. Let I and J be real intervals. We shall use the following notations.

$$h_I(\alpha) = \max_{n \in I} D_n(\alpha) - \min_{n \in I} D_n(\alpha),$$

$$D_n(\alpha, \beta) = D_n(\beta) - D_n(\alpha) = Z(n, \beta) - Z(n, \alpha) - n(\beta - \alpha),$$

and

$$\begin{aligned} h_{I,J}(\alpha, \beta) &= \\ &= \max_{n \in I} (\min_{n \in I} D_n(\alpha, \beta) - \max_{n \in J} D_n(\alpha, \beta)) , \min_{n \in I} D_n(\alpha, \beta) - \max_{n \in I} D_n(\alpha, \beta) . \end{aligned}$$

The following lemma involves Schmidt's basic idea.

LEMMA 2. Suppose $\alpha, \beta \in U$ and suppose that J, K are subintervals of an interval I . Then

$$h_I(\alpha) + h_I(\beta) \geq h_{J,K}(\alpha, \beta) + \frac{1}{2}(h_J(\alpha) + h_J(\beta) + h_K(\alpha) + h_K(\beta)).$$

PROOF. [5, Lemma 5].

We use Lemma 2 to show that the average value of $h_I(\alpha)$ in a sequence of well-spaced points α cannot be very small.

LEMMA 3. Let λ be a real number with $0 < \lambda \leq \frac{1}{2}$. Let c and t be positive integers with $3\lambda c \leq 4$. Put $m = (4c)^t$. Let I be a real interval $[x, y)$ with $x \geq 0$ of length at least m/λ . Let $\alpha_0, \alpha_1, \dots, \alpha_{m-1}$ be real numbers satisfying $0 < \alpha_j - \alpha_{j-1} \leq \lambda c/m$ for $j = 1, 2, \dots, m-1$ and $\alpha_{j+m/2} - \alpha_j \geq \lambda$ for $j = 0, 1, \dots, \frac{1}{2}m-1$. Then, for any sequence ω in U ,

$$(6) \quad \frac{1}{m} \sum_{j=0}^{m-1} h_I(\alpha_j) > \frac{t}{64c}.$$

PROOF. Let $J = [v, w)$ be any interval of length $m/(4c\lambda)$ with $v \geq 0$. Take integers a and b such that $v \leq a < v+1$ and $w-1 \leq b < w$. Suppose

$$(7) \quad Z_b(\alpha_{m-1}) - Z_a(\alpha_{m-1}) - Z_b(\alpha_0) + Z_a(\alpha_0) \leq \frac{m}{8c}.$$

Then, for $j = 0, 1, \dots, \frac{1}{2}m-1$,

$$\begin{aligned}
& D_b(\alpha_{j+\frac{1}{2}m}) - D_a(\alpha_{j+\frac{1}{2}m}) - D_b(\alpha_j) + D_a(\alpha_j) \\
& \leq Z_b(\alpha_{m-1}) - Z_a(\alpha_{m-1}) - Z_b(\alpha_0) + Z_a(\alpha_0) - (b-a)(\alpha_{j+\frac{1}{2}m} - \alpha_j) \\
& \leq \frac{m}{8c} - \left(\frac{m}{4c\lambda} - 2\right)\lambda = -\frac{m}{8c} + 2\lambda.
\end{aligned}$$

Hence,

$$\begin{aligned}
& h_J(\alpha_{j+\frac{1}{2}m}) + h_J(\alpha_j) \\
& = \max_{n \in J} D_n(\alpha_{j+\frac{1}{2}m}) - \min_{n \in J} D_n(\alpha_{j+\frac{1}{2}m}) + \max_{n \in J} D_n(\alpha_j) - \min_{n \in J} D_n(\alpha_j) \\
& \geq \frac{m}{8c} - 2\lambda \geq \frac{m}{8c} - 1.
\end{aligned}$$

On summing over j we obtain that under the supposition (7)

$$(8) \quad \frac{1}{m} \sum_{j=0}^{m-1} h_J(\alpha_j) \geq \frac{m}{16c} - \frac{1}{2}$$

for any positive interval J of length $m/(4c\lambda)$.

We use induction on t . For $t = 1$ we have $D_n(\alpha) + n\alpha \in \mathbb{Z}$. Let $j \in \{0, 1, \dots, \frac{1}{2}m-1\}$. By the conditions of the lemma we have

$$\lambda \leq \alpha_{j+\frac{1}{2}m} - \alpha_j \leq \frac{1}{2} \lambda c \leq \frac{2}{3}.$$

Since $\min(\frac{1}{6}, \frac{1}{2}\lambda) \geq \frac{\lambda}{3}$, we have $\|\alpha_j\| \geq \lambda/3$ or $\|\alpha_{j+\frac{1}{2}m}\| \geq \lambda/3$, where $\|\alpha\|$ denotes the distance from α to the nearest integer. We can therefore choose integers $i \in \{j, j+\frac{1}{2}m\}$ and $r, s \in I$ such that $D_r(\alpha_i) - D_s(\alpha_i) \geq 1/4$. Hence $h_I(\alpha_i) \geq 1/4$ and therefore

$$\frac{1}{m} \sum_{j=0}^{m-1} h_I(\alpha_j) \geq \frac{1}{m} \cdot \frac{m}{2} \cdot \frac{1}{4} = \frac{1}{8}.$$

This proves the lemma in case $t = 1$.

We now assume that the assertion of the lemma holds for $t-1$ and we shall deduce it for t . Put

$$J_i = [x + \frac{(i-1)m}{4\lambda c}, x + \frac{im}{4\lambda c}] \quad \text{for } i = 1, 2, 3, 4.$$

Let z_j be the number of pairs (μ, ξ_μ) with $x+m/(4\lambda c) \leq \mu < x+2m/(4\lambda c)$ and $\xi_\mu - p \in [\alpha_{j-1}, \alpha_j]$ for some integer p . Hence z_j is a non-negative integer. We distinguish two cases.

- (a) Assume $\sum_{j=1}^{m-1} z_j \leq m/(8c)$. Then (7) is fulfilled for $v = x+m/(4\lambda c)$, $w = x+m/(2\lambda c)$. Hence, by (8),

$$\frac{1}{m} \sum_{j=0}^{m-1} h_{J_2}(\alpha_j) \geq \frac{m}{16c} - \frac{1}{2} \geq \frac{t}{16c}.$$

Since $J_2 \in I$, this implies inequality (6).

- (b) Assume $\sum_{j=1}^{m-1} z_j > m/(8c)$. For every $r \in J_1$ and $s \in J_3$ we have

$$\begin{aligned} D_s(\alpha_{j-1}, \alpha_j) - D_r(\alpha_{j-1}, \alpha_j) &\geq z_j - (s-r)(\alpha_j - \alpha_{j-1}) \\ &\geq z_j - \frac{3m}{4\lambda c} \cdot \frac{\lambda c}{m} = z_j - \frac{3}{4}. \end{aligned}$$

Hence, for $j = 0, 1, \dots, m-1$, in case $z_j \geq 1$,

$$h_{J_1, J_3}(\alpha_{j-1}, \alpha_j) \geq \frac{1}{4} z_j.$$

By Lemma 2, or obviously if $z_j = 0$,

$$h_I(\alpha_{j-1}) + h_I(\alpha_j) \geq \frac{1}{4} z_j + \frac{1}{2} (h_{J_1}(\alpha_{j-1}) + h_{J_1}(\alpha_j) + h_{J_3}(\alpha_{j-1}) + h_{J_3}(\alpha_j)).$$

Since $h_I(\alpha_j) \geq \max(h_{J_1}(\alpha_j), h_{J_3}(\alpha_j)) \geq \frac{1}{2} h_{J_1}(\alpha_j) + \frac{1}{2} h_{J_3}(\alpha_j)$, we have

$$\begin{aligned} 2 \sum_{j=0}^{m-1} h_I(\alpha_j) &\geq \sum_{j=0}^{m-1} (h_{J_1}(\alpha_j) + h_{J_3}(\alpha_j)) + \frac{1}{4} \sum_{j=1}^{m-1} z_j + h_I(\alpha_0) \\ &\quad + h_I(\alpha_{m-1}) - \frac{1}{2} h_{J_1}(\alpha_0) - \frac{1}{2} h_{J_1}(\alpha_{m-1}) - \frac{1}{2} h_{J_3}(\alpha_0) - \frac{1}{2} h_{J_3}(\alpha_{m-1}) \geq \\ &\geq \sum_{j=0}^{m-1} h_{J_1}(\alpha_j) + \sum_{j=0}^{m-1} h_{J_3}(\alpha_j) + \frac{m}{32c}. \end{aligned}$$

On applying the induction hypothesis to J_i and the point sets $\{\alpha_{4cl+k}\}_{l=0}^{m/(4c)-1}$, we obtain

$$\begin{aligned} \sum_{j=0}^{m-1} h_{J_i}(\alpha_j) &= \sum_{k=0}^{4c-1} \sum_{l=0}^{m/(4c)-1} h_{J_i}(\alpha_{4cl+k}) \\ &> \sum_{k=0}^{4c-1} \frac{m}{4c} \cdot \frac{t-1}{64c} = \frac{m}{64c} (t-1) \end{aligned}$$

for $j = 1$ and $j = 3$. Hence,

$$\frac{1}{m} \sum_{j=0}^{m-1} h_I(\alpha_j) > \frac{t-1}{64c} + \frac{1}{64c} = \frac{t}{64c}.$$

This proves Lemma 3.

6. As an application of Lemma 3 we derive the following theorem.

THEOREM 5. Let γ and δ be real numbers with $0 \leq \gamma < \delta \leq 1$. Let H be some positive integer. Let $\gamma = \alpha_1, \alpha_2, \dots, \alpha_N = \delta$ be real numbers satisfying $0 < \alpha_{i+1} - \alpha_i \leq (\delta - \gamma)/H$ for $i = 1, 2, \dots, N-1$. Then for every sequence ω

$$(8) \quad \max_{i=1,2,\dots,N} D(\omega, \alpha_i) \geq \frac{1}{2000} \log \frac{H}{48}.$$

PROOF. Put $\ell = \delta - \gamma$. Let $t = \lceil \log(H/3)/\log 16 \rceil$. So $H/48 < 16^t \leq H/3$. Split $[\gamma, \delta]$ into $3 \cdot 16^t$ parts of equal lengths and choose in every third part a point from $\{\alpha_1, \alpha_2, \dots, \alpha_N\}$. This is possible, since $\ell/3 \cdot 16^t \geq \ell/H$. This gives $m = 16^t$ points $\beta_1, \beta_2, \dots, \beta_m$ with $\beta_j - \beta_{j-1} \leq 4\ell/(3m)$. Further $\beta_{j+\frac{1}{2}m} - \beta_j \geq \ell/3$. We apply Lemma 3 with $\lambda = \ell/3$ and $c = 4$. Hence

$$\frac{1}{m} \sum_{j=0}^{m-1} h_I(\beta_j) > \frac{t}{256} > \frac{\log(H/48)}{256 \log 16} > \frac{1}{1000} \log \frac{H}{48}.$$

It follows that for any sequence ω

$$\max_{j=0,1,\dots,m-1} D(\omega, \beta_j) > \frac{1}{2000} \log \frac{H}{48}.$$

In particular (8) holds.

COROLLARY 1. Let S be a BDS. Then there exists an $\varepsilon > 0$ such that every subinterval of U of length ℓ contains a subinterval J of length at least $\varepsilon\ell$ with $J \cap S = \emptyset$.

PROOF. Let S be any BDS. Let ω be a sequence and κ a positive number such that

$$D(\omega, \alpha) \leq \kappa \quad \text{for every } \alpha \in S.$$

Let $[\gamma, \delta)$ be any subinterval of U . Choose H so large that

$$\frac{1}{2000} \log \frac{H}{48} > \kappa.$$

Put $\varepsilon = H^{-1}$. Then, by Theorem 5, $\max_{i=1, \dots, N} D(\omega, \alpha_i) > \kappa$ for any set $\{\alpha_1, \dots, \alpha_N\}$ in $[\gamma, \delta)$ with $0 < \alpha_{j+1} - \alpha_j \leq \varepsilon(\delta - \gamma)$ for $j = 1, 2, \dots, N-1$. Thus S does not contain such a subset. This proves the corollary.

The following result shows that Theorems 2 and 3 cannot be improved by more than a constant factor. (The constant $(4000)^{-1}$ can be improved considerably.)

COROLLARY 2. Let $n > 48^2$. Then for every sequence ω

$$\max_{j=0, 1, \dots, n-1} D(\omega, \frac{j}{n}) \geq \frac{1}{2000} \log \frac{n}{48} \geq \frac{1}{4000} \log n.$$

7. It follows from Corollary 1 that $S = \{\frac{1}{n}\}_{n=2}^{\infty}$ is not a BDS. This result is also a consequence of the following theorem which gives a necessary and sufficient condition for sequences satisfying a certain regularity condition.

THEOREM 6. Let $\alpha_1, \alpha_2, \dots$ be a strictly decreasing sequence with limit 0. Suppose there exists a constant c such that $\alpha_{n-1} - \alpha_n \leq c(\alpha_{m-1} - \alpha_m)$ for every n and m with $n \geq m$. Then $S = \{\alpha_1, \alpha_2, \dots\}$ is a BDS if and only if for some positive integer h

$$\limsup_{n \rightarrow \infty} \log \frac{\alpha_{n+h}}{\alpha_n} < 0.$$

PROOF. Suppose $\limsup_{n \rightarrow \infty} \log \frac{\alpha_{n+h}}{\alpha_n} \alpha_n^{-1} < 0$. Then there exists a constant $c < 1$ such that $\alpha_{n+h} < c\alpha_n$ for $n = 1, 2, \dots$. It follows from Theorem 4 that S is a BDS. (Here we did not use the regularity condition.)

Suppose S is a BDS. Then by Corollary 1 there exists a positive number ε such that every interval $[0, \alpha_n)$ contains an interval J of length $\varepsilon\alpha_n$ such that $S \cap J = \emptyset$. Let k be such that $J \subset (\alpha_{n+k}, \alpha_{n+k-1})$. Then

$$\max_{j=0, \dots, k} \alpha_{n+j-1} - \alpha_{n+j} > c^{-1}(\alpha_{n+k-1} - \alpha_{n+k}) \geq \varepsilon\alpha_n c^{-1}.$$

Hence,

$$\alpha_n \geq \alpha_n - \alpha_{n+k} \geq \varepsilon k \alpha_n c^{-1}.$$

Thus $k \leq c\varepsilon^{-1}$ is bounded, which implies that for $h = [c\varepsilon^{-1}]$

$$\limsup_{n \rightarrow \infty} \log \frac{\alpha_{n+h}}{\alpha_n} < \log(1-\varepsilon) < 0.$$

REFERENCES

- [1] AARDENNE-EHRENFEST, T. van, *Proof of the impossibility of a just distribution*, Indag. Math. 7 (1945), 71-76.
- [2] CORPUT, J.G. van der, *Verteilungsfunktionen, I*. Proc. Kon. Nederl. Akad. Wetensch. 38 (1935), 813-821.
- [3] ERDÖS, P., *Problems and results on diophantine approximation*, Compositio Math. 16 (1964), 52-66.
- [4] SCHMIDT, W.M., *Irregularities of distribution*, Quart. J. Math. (Oxford), 19 (1968), 181-191.
- [5] _____, *Irregularities of distribution VI*, Compositio Math. 24 (1972), 63-74.
- [6] _____, *Irregularities of distribution VIII*, Trans. Amer. Math. Soc. 198 (1974), 1-22.
- [7] _____, *Lectures on irregularities of distribution*, Tata Institute of Fundamental Research, Bombay, 1977.

- [8] SHAPIRO, L., *Regularities of distribution*, Studies in probability and ergodic theory, *Advances in mathematics supplementary studies*, vol. 2, Academic Press, New York etc., 1978, pp. 135-154.

ONTVANGEN 1 2 DEC. 1979