

**stichting
mathematisch
centrum**



AFDELING ZUIVERE WISKUNDE

ZW43/75

APRIL

K. HARDY AND I. JUHÁSZ

NORMALITY AND THE WEAK cb PROPERTY

Prepublication

2e boerhaavestraat 49 amsterdam

BIBLIOTHEEK MATHEMATISCH CENTRUM
AMSTERDAM

Printed at the Mathematical Centre, 49, 2e Boerhaavestraat, Amsterdam.

The Mathematical Centre, founded the 11-th of February 1946, is a non-profit institution aiming at the promotion of pure mathematics and its applications. It is sponsored by the Netherlands Government through the Netherlands Organization for the Advancement of Pure Research (Z.W.O), by the Municipality of Amsterdam, by the University of Amsterdam, by the Free University at Amsterdam, and by industries.

Normality and the weak cb property

by

K. Hardy and I. Juhász

ABSTRACT

It is demonstrated that the Alexandroff duplicate of a Dowker space is again a Dowker space which is not weak cb, while the existence of weak cb Dowker spaces is made manifest.

KEY WORDS AND PHRASES: *Dowker space, cb-space, weak cb-space, Alexandroff duplicate.*

A non-metrizable, first countable compact space was created by ALEXANDROFF in [1] and the construction has been subsequently generalized and employed (ENGELKING [2,3], JUHÁSZ [7,8]). The present note concentrates on some properties of the Alexandroff duplicate $A(X)$ which, in particular, show that a normal space need not have the weak cb property, thus resolving the open question in MACK [11, p.240].

1. PRELIMINARIES

No separation axioms are implicitly assumed for the topological space X . The Hewitt-Nachbin realcompactification of a Tychonoff space X is denoted by νX . We will write $A_n \searrow \emptyset$ to indicate that (A_n) is a decreasing sequence of subsets of X such that $\bigcap_n A_n = \emptyset$. \mathbb{N} denotes the natural numbers. A set A is regular closed if $A = \text{cl}_X \text{int}_X A$, and ∂A denotes the boundary of A .

PROPOSITION 1.1. (MACK [10,11]) *A space X is cb (weak cb) if and only if for each sequence $A_n \searrow \emptyset$ of closed (regular closed) subsets of X , there exists a sequence of zero sets (Z_n) with $A_n \subseteq Z_n$ and $\bigcap_n Z_n = \emptyset$.*

cb-spaces originated in HORNE [5] and were studied by MACK in [10]. Every normal, countably paracompact space is cb and every cb-space is countably paracompact. Weak cb-spaces were defined in [11]. They form a natural generalization of cb-spaces and include the Tychonoff pseudocompact spaces and all extremally disconnected spaces. Interest in weak cb-spaces is centered in the theorem ([11]) that for a Tychonoff space X , the Dedekind completion of $C(X)$ is isomorphic to $C(Y)$, for some space Y , if and only if νX is weak cb. It should be noted that if X is Tychonoff and weak cb, then any space T with $X \subseteq T \subseteq \nu X$ is weak cb. The converse fails in general (see HARDY & WOODS [4]) but the following result is evident and will be needed below.

PROPOSITION 1.2. *Let X be Tychonoff and consider the statements*

(a) *For any sequence $A_n \searrow \emptyset$ of regular closed sets in X we have*

$$\bigcap_n \text{cl}_{\nu X} A_n = \emptyset$$

(b) For any decreasing sequence (A_n) of regular closed sets in X we have

$$\bigcap_n \text{cl}_{\cup X} A_n = \text{cl}_{\cup X} \bigcap_n A_n$$

(c) If $\cup X$ is weak cb, then any space T , with $X \subseteq T \subseteq \cup X$ is weak cb.

Then (a) if and only if (b); and (a) or (b) implies (c).

PROOF. We merely recall that if X is dense in T and A is a regular closed subset of X then $\text{cl}_T A = B$ is the unique regular closed subset of T with $A = B \cap X$. \square

According to a result of ISHIKAWA [6], a space X is countably paracompact if and only if for each sequence $A_n \searrow \emptyset$ of closed subsets of X , there exists a sequence (G_n) of open sets such that $A_n \subseteq G_n$ and $\bigcap_n \text{cl}_X G_n = \emptyset$. The following observation will be useful below and may have independent interest.

PROPOSITION 1.3. *The following statements are equivalent*

- (a) X is countable paracompact.
- (b) For each sequence $F_n \searrow \emptyset$ of closed nowhere dense subsets of X , there exists a sequence (G_n) of open sets such that $F_n \subseteq G_n$ and $\bigcap_n \text{cl}_X G_n = \emptyset$.
- (c) Each countable increasing cover ([10]) by dense open sets has a countable closed refinement whose interiors cover X .

PROOF. It is enough to show (b) implies (a). Let $A_n \searrow \emptyset$ be an arbitrary sequence of closed sets and define a sequence of open sets (G_n) with $A_n \subseteq G_n$ and $\bigcap_n \text{cl}_X G_n = \emptyset$ in the following manner:

- (i) If $\text{int}_X A_m = \emptyset$ for some $m \geq 1$, there exist open sets G_k with $A_k \subseteq G_k$, $k \geq m$ and $\bigcap_k \text{cl}_X G_k = \emptyset$; put $G_n = X$ for $1 \leq n < m$. Now assume that $\text{int}_X A_n \neq \emptyset$ for all n .
- (ii) If a subsequence (A_{n_k}) exists with $A_{n_{k+1}} \subseteq \text{int}_X A_{n_k}$, let $G_{n_{k+1}} = \text{int}_X A_{n_k}$ and $G_n = X$ otherwise.
- (iii) If there exists $m \geq 1$ such that $F_k = \partial A_k \cap \partial A_{k+1} \neq \emptyset$ for $k \geq m$ then $F_k \searrow \emptyset$ is a sequence of closed nowhere dense sets and there exists a sequence of open sets (U_k) with $F_k \subseteq U_k$ and $\bigcap_k \text{cl}_X U_k = \emptyset$. Define $G_{k+1} = \text{int}_X A_k \cup U_k$ for $k \geq m$ and $G_n = X$ for $1 \leq n \leq m$. \square

In order to exploit the use of nowhere dense closed subsets, we venture to make the following

DEFINITION 1.4. *X is an nd-space if for each sequence $F_n \searrow \emptyset$ of closed nowhere dense sets, there exists a sequence of zero sets (Z_n) with $F_n \subseteq Z_n$ and $\bigcap_n Z_n = \emptyset$.*

Every cb-space is an nd-space. Since every zero set Z is a regular G_δ -set (a countable intersection of closed sets whose interiors contain Z), we may adapt the proof of Proposition 1.3 to conclude that every nd-space is countable paracompact. A space is cb if and only if it is both a weak cb and an nd-space. The example on p.240 of [11] is countably paracompact but not an nd-space. It is conjectured that an nd-space need not be cb, although an example at the present time is not forthcoming.

2. PROPERTIES OF $A(X)$

Recall the construction in [7]. Given an arbitrary topological space X , consider the set $A(X) = X \cup X'$, where X' is a disjoint copy of X . For any $x \in X$, let x' denote the corresponding point of X' and if $S \subseteq X$ define $S' = \{x' \mid x \in S\}$. A topology is introduced to $A(X)$ by defining a base $\{B(z) \mid z \in A(X)\}$ as follows:

$$B(x') = \{\{x'\}\} \text{ and } B(x) = \{V \cup (V' \setminus \{x'\}) \mid V \in \mathcal{V}(x)\},$$

where $\mathcal{V}(x)$ is a neighbourhood base of x in X . The resulting space, also denoted by $A(X)$, generalizes the original construction in ALEXANDROFF & URYSOHN [1] and is called the Alexandroff duplicate of X . It is clear that X is a closed, C -embedded subspace of $A(X)$.

Many properties of X are shared with $A(X)$. It has been noticed that $A(X)$ is compact ([2]), α -compact (for any infinite cardinal α), realcompact and Tychonoff ([7]), if X has the corresponding property. We will now expand this list of properties.

Observe that a space is normal if and only if each pair of disjoint closed nowhere dense sets can be separated by disjoint open neighbourhoods.

PROPOSITION 2.1. *X is normal if and only if $A(X)$ is normal.*

PROOF. Let A and B be disjoint closed nowhere dense subsets of $A(X)$. Then A and B are closed and disjoint in X and can be separated by disjoint open sets U and V in X. The sets $U \cup U'$ and $V \cup V'$ are open disjoint neighbourhoods of A and B in $A(X)$. \square

PROPOSITION 2.2. *X is countably paracompact if and only if $A(X)$ is countably paracompact.*

PROOF. For the necessity, let $F_n \searrow \emptyset$ be a sequence of closed nowhere dense subsets of $A(X)$. Then $F_n \subseteq X$ and there exists a sequence (V_n) of open subsets of X with $F_n \subseteq U_n$ and $\bigcap_n \text{cl}_X U_n = \emptyset$. Define $G_n = U_n \cup U'_n$ and note that $\text{cl}_{A(X)} G_n = \text{cl}_X U_n \cup U'_n$, so that $F_n \subseteq G_n$ and $\bigcap_n \text{cl}_{A(X)} G_n = \emptyset$. \square

PROPOSITION 2.3. *If $A(X)$ is weak cb then both X and $A(X)$ are cb.*

PROOF. To show that X is cb, take a sequence $A_n \searrow \emptyset$ of closed sets in X. Then $B_n = A_n \cup A'_n$ is regular closed in $A(X)$ and $B_n \searrow \emptyset$. There exist zero sets W_n in $A(X)$ with $B_n \subseteq W_n$ and $\bigcap_n W_n = \emptyset$. Then $Z_n = W_n \cap X$ is a zero set in X and $A_n \subseteq Z_n$ with $\bigcap_n Z_n = \emptyset$. If X is cb then both X and $A(X)$ are countably paracompact, hence $A(X)$ is cb. \square

One may show that $A(X)$ is countably compact if and only if X is. Furthermore, if X contains a C-embedded copy of \mathbb{N} , so does $A(X)$ so that $A(X)$ is pseudocompact implies that X is also. However, if X is pseudocompact (Tychonoff) but not countable compact then $A(X)$ is not weak cb, in particular, not pseudocompact.

3. DOWKER SPACES

A Dowker space is a normal Hausdorff space which is not countably paracompact. Such spaces exist within Zermelo-Fraenkel set theory; the axiom of choice implies the existence of a zero-dimensional P-space which is Dowker (RUDIN [12]) and more recently a certain combinatorial principle called \diamond implies existence of a locally compact, first countable, hereditarily sep-

arable Dowker space (JUHÁSZ et al. [9]).

The open question in [11, p.240] may be phrased as follows: *Must every Dowker space have the weak cb property?* It follows from Propositions 2.1 and 2.2 that $A(X)$ is a Dowker space if and only if X is such. Since no Dowker space can be even an nd -space, 2.3 implies that for any Dowker space X , the space $A(X)$ answers the above question negatively. It may be of interest however that the Dowker space of M.E. RUDIN [12] is weak cb, as is now shown.

The reader is referred to [12] for details. With the same notation as in [12], define

$$F = \{f: \mathbb{N} \rightarrow \omega_\omega \mid f(n) \leq \omega_n \text{ for all } n \in \mathbb{N}\}.$$

$$X = \{f \in F \mid \omega_1 \leq cf(f(n)) \leq \omega_k \text{ for all } n \in \mathbb{N} \text{ and some } k \in \mathbb{N}\},$$

$$X' = \{f \in F \mid \omega_1 \leq cf(f(n)) \text{ for all } n \in \mathbb{N}\}.$$

F carries a topology generated by the basic open-and-closed sets

$$(f, g] = \{h \in F \mid f(n) < h(n) \leq g(n) \text{ for all } n \in \mathbb{N}\}.$$

Then $X \subseteq X' \subseteq F$ are subspaces and $\cup X = X'$ is paracompact, and hence a weak cb-space.

To show that X is weak cb, let $A_n \searrow \emptyset$ be a sequence of regular closed subsets of X and suppose $g \in \bigcap_{n \in \mathbb{N}} cl_{\cup X} A_n$. We will define an increasing sequence $\{f_\alpha \in X \mid \alpha < \omega_1\}$ as follows:

1) Choose any $f_0 \in \text{int}_{X^1} A_1$ with $f_0 \leq g$.

2) Assume $f_\beta \in X$ is defined for all $\beta < \alpha$, and

(a) if $\alpha = \beta + 1$, let $i \in \mathbb{N}$ be the smallest integer with $f_\beta \notin \text{int}_{X^1} A_i$ and choose $f_\alpha \in (\text{int}_{X^1} A_i) \cap (f_\beta, g]$.

(b) if α is a limit ordinal, let $h_\alpha(n) = \sup\{f_\beta(n) \mid \beta < \alpha\}$ and choose $f_\alpha \in (\text{int}_{X^1} A_1) \cap (h_\alpha, g]$.

Now define $f(n) = \sup\{f_\alpha(n) \mid \alpha < \omega_1\}$. Then $f \leq g$ and $cf(f(n)) = \omega_1$ for all $n \in \mathbb{N}$ implies that $f \in X$. However, $f \in A_k$ for all $k \in \mathbb{N}$: let $h < f$ and for each $n \in \mathbb{N}$ there is $f_{\alpha_n} \in \{f_\alpha \mid \alpha < \omega_1\}$ with $h(n) < f_{\alpha_n}(n)$. Let $\beta = \sup\{\alpha_n \mid n \in \mathbb{N}\}$ and then $f_{\beta+k} \in (\text{int}_{X^1} A_k) \cap (h, f]$, that is

$$f \in \text{cl}_{X^k}^{\text{int}} A_k = A_k.$$

We have a contradiction and so $A_n \setminus \emptyset$ implies $\bigcap_n \text{cl}_{\cup X} A_n = \emptyset$. Finally, apply Proposition 1.2 to infer that X is weak cb.

4. REMARKS

Since the Dowker space X in [12] is weak cb, it follows from [4] that $E(\cup X) = \cup E(X)$, where $E(X)$ denotes the absolute of X (see for example [4, p.652]). Thus, $\cup E(X)$ is paracompact. However, it has been shown by E.K. VAN DOUWEN that $E(X)$ is not normal. It would seem natural therefore to pose the following questions. 1) Is there a normal (non-paracompact) space X with normal absolute $E(X)$; 2) Is there an extremally disconnected Dowker space; and ultimately 3) Is there a Dowker space X with Dowker absolute $E(X)$.

REFERENCES

- [1] ALEXANDROFF, P. & P. URYSOHN, *Mémoire sur les espaces topologiques compacts*, Verh. Akad. Wetensch. Amsterdam, 14 (1929), 1-96.
- [2] ENGELKING, R., *On the double circumference of Alexandroff*, Bull. Acad. Pol. Sci., 16 (1968), 629-634.
- [3] ENGELKING, R., *On functions defined on Cartesian products*, Fund. Math., 59 (1966), 221-231.
- [4] HARDY, K. & R.G. WOODS, *On c -realcompact spaces and locally bounded normal functions*, Pacific J. Math., 43 (1972), 646-656.
- [5] HORNE, J.G., *Countable paracompactness and cb-spaces*, Notices Amer. Math. Soc. 6 (1959), 629-630.
- [6] ISHIKAWA, F., *On countably paracompact spaces*, Proc. Japan Acad., 31 (1955), 686-687.
- [7] JUHÁSZ, I., *On closed discrete subspaces of product spaces*, Bull. Acad. Pol. Sci., 17 (1969), 219-223.

- [8] JUHÁSZ, I. & S. MRÓWKA, *E-compactness and the Alexandroff duplicate*, Indagationes Math., 73 (1970), 26-29.
- [9] JUHÁSZ, I., K. KUNEN, & M.E. RUDIN, *Examples of Dowker spaces*, to appear.
- [10] MACK, J., *On a class of countably paracompact spaces*, Proc. Amer. Math. Soc., 16 (1965), 467-472.
- [11] MACK, J. & D. JOHNSON, *The Dedekind completion of $C(X)$* , Pacific J. Math., 20 (1967), 231-243.
- [12] RUDIN, M.E., *A normal space X for which $X \times I$ is not normal*, Fund. Math., 73 (1971), 179-186.