

STICHTING
MATHEMATISCH CENTRUM
2e BOERHAAVESTRAAT 49
AMSTERDAM

ZW 1956-026^{*u*}

On the Minkowski-Hlawka theorem

C.G. Lekkerkerker



MATHEMATICS

ON THE MINKOWSKI-HLAWKA THEOREM

BY

C. G. LEKKERKERKER

(Communicated by Prof. J. F. KOKSMA at the meeting of March 24, 1956)

Let $R_n (n \geq 2)$ be the real Euclidean space of points $x = (x_1, \dots, x_n)$. Let O be the origin. The theorem of MINKOWSKI-HLAWKA asserts that, if K is a bounded star domain in R_n , of volume V and symmetric about O , there exists a lattice, which is admissible for K and whose determinant does not exceed $V/2\zeta(n)$, where $\zeta(n) = 1 + 2^{-n} + 3^{-n} + \dots$. Various proofs of this theorem have been given [1, 2, 3, 4, 5]. In all these proofs the lattice found may have a rather contorted form: if S is any sphere about O and the volume of K has a fixed value, then, for suitably chosen K , these proofs lead to a lattice which does not have a basis contained in S . It is also clear that the proofs of SIEGEL [6] and WEIL [7] do not give us any information concerning the form of the lattices which have the required properties.

By applying the BRUNN-MINKOWSKI theorem, MAHLER [8] and DAVENPORT-ROGERS [9] obtained improvements of the theorem of MINKOWSKI-HLAWKA in the case of convex bodies. The results of these authors are as follows. For $n \geq 2$, let c_n denote the lower bound of V/Δ for all centrally symmetric convex bodies in R_n , V being the volume and Δ denoting the critical determinant of K ; it is the largest positive number such that, for all K , there exists a K -admissible lattice with determinant not exceeding V/c_n . By the theorem of MINKOWSKI-HLAWKA, $c_n \geq 2\zeta(n)$. Now MAHLER found that

$$(1) \quad c_2 \geq 2\sqrt{3}, \quad c_n > 2\zeta(n) + 1/6 \text{ for all } n,$$

whereas DAVENPORT-ROGERS proved that

$$\liminf_{n \rightarrow \infty} c_n \geq c,$$

where $c = 4.921 \dots$ is the solution of

$$(2) \quad c \log c = 2(c-1) \quad (c > 1).$$

In this note I shall prove two theorems. Firstly, I shall show

Theorem 1. *Let c be defined by (2). Then $c_n > c$ for $n \geq 5$.*

Next, I shall show that, for an arbitrary convex body K , by using the method of proof employed by MAHLER and DAVENPORT-ROGERS, one can find a lattice which has the properties discussed and, in addition,

has a basis contained in some relatively small sphere about O . More precisely, I shall prove the following

Theorem 2. *Let the numbers $\bar{c}_2, \bar{c}_3, \dots$ be given by*

$$(3) \quad \bar{c}_2=3, \bar{c}_3=3.82, \bar{c}_4=4.41, \bar{c}_5=4.80, \bar{c}_6=c \text{ for } n \geq 6.$$

Let κ_n be the volume of the n -dimensional unit sphere. Let K be a convex body in R_n , of volume V and symmetric about O . Then there exists a K -admissible lattice, whose determinant does not exceed V/\bar{c}_n and which, after a suitable rotation about O , has a basis contained in the cube defined by

$$(4) \quad |x_i| < b(V/\kappa_n)^{1/n} \quad (i = 1, \dots, n),$$

where, for all n , one may take $b=2.13$.

We note that, since $\kappa_n = \pi^{n/2} / \Gamma(\frac{n+2}{2})$, $\kappa_n^{-1/n}$ is asymptotically equal to $\sqrt{n/(2\pi e)}$. Further, the cube defined by (4) is contained, also after an arbitrary rotation, in the sphere with centre at O and radius $\sqrt{n} \cdot b(V/\kappa_n)^{1/n}$. So the assertion of theorem 2 can also be stated in the following somewhat weaker form: *there exists a K -admissible lattice, whose determinant does not exceed V/\bar{c}_n and which has a basis contained in the sphere*

$$x_1^2 + \dots + x_n^2 < (b_1 n V^{1/n})^2,$$

where b_1 is some positive constant not depending on K and n .

The proofs of the above theorems will be preceded by a number of lemmas. The first two of these lemmas are the main steps in the proof of DAVENPORT-ROGERS and are also fundamental for our purpose. For the proofs I refer to the paper mentioned above. Further, in particular, lemmas 4 and 5 deal with the volumes of the sections of a convex body or a star body by planes through O . Lemma 5 is perhaps of interest by itself.

The following notations will be used. For given K and real a , denote by $V(a)$ the $(n-1)$ -dimensional volume of the section of K by $x_n = a$. For each $(n-1)$ -dimensional hyperplane Π through O , let $v(\Pi)$ denote the $(n-1)$ -dimensional volume of $K \cap \Pi$. Further, for each point x , write $|x| = (x_1^2 + \dots + x_n^2)^{1/2}$. If $x \neq O$, then let Π_x denote the $(n-1)$ -dimensional hyperplane, which passes through O and is orthogonal to the vector x . Finally, let ω_n denote the area of the unit sphere in R_n , so that

$$(5) \quad \omega_n = n\kappa_n \quad (n \geq 2).$$

Lemma 1.¹⁾ *Let \mathcal{L} be an $(n-1)$ -dimensional lattice in the space*

¹⁾ Lemma 1 says a little more than lemma 2 in the paper of DAVENPORT-ROGERS, but follows immediately from the proof of that lemma. Lemma 2 follows from their lemma 3, if one puts $\beta = (1 + \delta/n)^{-n+1}$ and notes that

$$1 < \frac{n(1-\beta^{1/(n-1)})}{1-\beta^{n/(n-1)}} < n, \text{ if } 0 < \beta < 1.$$

$x_n = 0$, with determinant $d(\mathcal{L})$, which has no point (except O) in K . Suppose that $\alpha > 0$, and

$$(6) \quad \sum_{t=1}^{\infty} V(\alpha t) < d(\mathcal{L}).$$

Then there exists a point g of the form $g = (g_1, \dots, g_{n-1}, \alpha)$, such that the lattice generated by \mathcal{L} and g is admissible for K .

Lemma 2. Let β be a real number with $0 < \beta < 1$ and let α be defined by

$$(7) \quad \alpha = \frac{V}{2V(0)} \cdot \frac{n(1-\beta)^{1/(n-1)}}{1-\beta^{n/(n-1)}}.$$

Then

$$(8) \quad \sum_{t=0}^{\infty} V(\alpha t) \leq \beta V(0).$$

It is convenient to deduce from lemma 1 the following

Lemma 3. The assertion of lemma 1 remains true, if \mathcal{L} is allowed to have points on the boundary of K and if the condition (6) is replaced by

$$(6') \quad \sum_{t=1}^{\infty} V(\alpha t) \leq d(\mathcal{L}).$$

Proof. Let \mathcal{L} and α be such that (6') holds and that \mathcal{L} has no point (except O) in the interior of K . For each positive integer r , let \mathcal{L}_r be the lattice $(1+r^{-1})\mathcal{L}$. Then \mathcal{L}_r has no point (except O) in K and (6) holds, with \mathcal{L}_r instead of \mathcal{L} . So, by lemma 1, there exists a point $g^{(r)}$ in the plane $x_n = \alpha$, such that the lattice \mathcal{A}_r generated by \mathcal{L}_r and $g^{(r)}$ is admissible for K . Clearly there exists an increasing sequence of positive integers r_1, r_2, \dots , such that \mathcal{A}_{r_k} , for $k \rightarrow \infty$, converges to a lattice \mathcal{A} generated by \mathcal{L} and some point g in the plane $x_n = \alpha$. By a familiar argument¹⁾, \mathcal{A} is admissible for K . This proves the lemma.

Lemma 4. Let K be a convex body in R_n , of volume $V = \kappa_n$, symmetric about O . Then there exists an $(n-1)$ -dimensional hyperplane Π , which passes through O , such that

$$(9) \quad \frac{1}{2} \kappa_n \leq v(\Pi) \leq \frac{1}{2} n \kappa_n.$$

Proof. Let x' be a point on the boundary of K , for which $|x'|$ is maximal. Since $V = \kappa_n$, K is not properly contained in the unit sphere, and so $|x'| \geq 1$. Write $\Pi' = \Pi_{x'}$, and consider the cone C , with vertex at x' and with the intersection $K \cap \Pi'$ as a basis.

Since K is convex, C is contained in K . Hence, since K is also symmetric about O ,

$$V = \kappa_n \geq 2V(C) = \frac{2}{n} |x'| \cdot v(\Pi') \geq \frac{2}{n} v(\Pi'),$$

hence

$$v(\Pi') \leq \frac{1}{2} n \kappa_n.$$

¹⁾ See e.g. MAHLER [10], proof of theorem 8.

Next, let x'' be a point on the boundary of K , such that $|x''|$ is minimal. Since the unit sphere is not properly contained in K , we now have $|x''| \leq 1$. Write $\Pi'' = \Pi_{x''}$ and, for real a , denote by $W(a)$ the $(n-1)$ -dimensional volume of the intersection of K and the plane which passes through ax'' and is parallel to Π'' . In particular, $W(0) = v(\Pi'')$. In virtue of the BRUNN-MINKOWSKI theorem¹⁾, the expression $\sqrt[n-1]{W(a)}$ is a concave function of a . Further, by the symmetry of K , this function is even. Hence $W(a) \leq W(0)$, for all a , and so

$$V = \kappa_n \leq 2|x''| \cdot W(0) \leq 2W(0) = 2v(\Pi'').$$

Hence

$$v(\Pi'') \geq \frac{1}{2}\kappa_n.$$

Since the point of intersection of the boundary of K and a straight line through O varies continuously, as the direction of this line varies, the volume $v(\Pi)$ varies continuously with Π . Then it follows from the above estimates for $v(\Pi')$ and $v(\Pi'')$ that Π may be chosen such that (9) holds.

When K is the n -dimensional unit sphere, then, for all Π , $v(\Pi)$ is equal to κ_{n-1} . Here, since $\kappa_n = \tau^{n/2} / \Gamma(\frac{n+2}{2})$, the number κ_{n-1} is asymptotically equal to $\sqrt{n/(2\pi)} \cdot \kappa_n$. One might conjecture that in lemma 4 the plane Π can always be chosen in such a way that $v(\Pi) = \kappa_{n-1}$. But it seems difficult to decide whether this is true. In the following lemma I shall prove (for a much wider class of bodies) that a certain mean value of $v(\Pi)$ is at most equal to κ_{n-1} . As a consequence, in lemma 4 the plane Π may be chosen such that instead of (9) we have

$$(9') \quad \frac{1}{2}\kappa_n \leq v(\Pi) \leq \kappa_{n-1}.$$

Lemma 5. *Let K be a bounded star domain (not necessarily symmetric about O), of volume κ_n . Let S_{n-1} be the sphere $x_1^2 + \dots + x_n^2 = 1$. For $x \in S_{n-1}$, let $v(\Pi_x)$ be the $(n-1)$ -dimensional volume of $K \cap \Pi_x$. Then we have*

$$(10) \quad \left[\frac{1}{\omega_n} \int_{S_{n-1}} \{v(\Pi_x)\}^{n/(n-1)} dx \right]^{(n-1)/n} \leq \kappa_{n-1}.$$

Proof. For $x \in S_{n-1}$, denote by $f(x)$ the uniquely determined positive number λ , for which λx belongs to the boundary of K . Clearly $f(x)$ is a positive, continuous function. Expressing the volume of K in terms of $f(x)$ we get

$$\kappa_n = \frac{1}{n} \int_{S_{n-1}} \{f(x)\}^n dx.$$

For given $x \in S_{n-1}$, let us denote by $S_{n-2}(x)$ the set of points y with

$$y_1^2 + \dots + y_n^2 = 1, \quad y_1 x_1 + \dots + y_n x_n = 0.$$

¹⁾ See BONNESEN-FENCHEL [11], pp. 71 and 88.

Then for $v(\Pi_x)$ we have the expression

$$v(\Pi_x) = \frac{1}{n-1} \int_{S_{n-1}(x)} \{f(y)\}^{n-1} dy.$$

To the last integral we apply HÖLDER'S inequality. This gives

$$\begin{aligned} v(\Pi_x) &\leq \frac{1}{n-1} \left[\int_{S_{n-2}(x)} \{f(y)\}^n dy \right]^{(n-1)/n} \cdot \left[\int_{S_{n-2}(x)} dy \right]^{1/n} \\ &= \frac{1}{n-1} \omega_{n-1}^{1/n} \left[\int_{S_{n-2}(x)} \{f(y)\}^n dy \right]^{(n-1)/n}, \end{aligned}$$

hence, on account of (5),

$$\{v(\Pi_x)\}^{n/(n-1)} \leq \frac{1}{n-1} \kappa_{n-1}^{1/(n-1)} \int_{S_{n-2}(x)} \{f(y)\}^n dy.$$

The inequality (10) will follow if we can prove that

$$(11) \quad \int_{S_{n-1}} dx \int_{S_{n-2}(x)} \{f(y)\}^n dy = \omega_{n-1} \int_{S_{n-1}} \{f(y)\}^n dy;$$

for then we have

$$\int_{S_{n-1}} \{v(\Pi_x)\}^{n/(n-1)} dx \leq \frac{1}{n-1} \kappa_{n-1}^{1/(n-1)} \omega_{n-1} \cdot n \kappa_n = \kappa_{n-1}^{n/(n-1)} \omega_n.$$

Let δ be a small positive number. We define a function $\phi(x, y)$ of two independent variables x, y , as follows:

$$\phi(x, y) = \begin{cases} \{f(y)\}^n & \text{if } |xy| \leq \delta \\ 0 & \text{if } |xy| > \delta \end{cases} \quad (x, y \in S_{n-1}),$$

where $xy = x_1 y_1 + \dots + x_n y_n$. Since $f(y)$ is a continuous function of y , we certainly have

$$\int_{S_{n-1}} \int_{S_{n-1}} \phi(x, y) dx dy = \int_{S_{n-1}} \int_{S_{n-1}} \phi(x, y) dy dx.$$

We further have

$$\begin{aligned} \int_{S_{n-1}} \phi(x, y) dy &\sim 2\delta \int_{S_{n-2}(x)} \{f(y)\}^n dy \text{ as } \delta \rightarrow 0, \\ \int_{S_{n-1}} \phi(x, y) dx &= \{f(y)\}^n \int_{|xy| \leq \delta} dx \sim 2\delta \omega_{n-1} \{f(x)\}^n \text{ as } \delta \rightarrow 0. \end{aligned}$$

From these relations (11) follows. This proves the lemma.

Lemma 6. *Let c be the solution of (2). Then, if $a > c$ and $n \geq 5$,*

$$(12) \quad \frac{2}{n} \frac{a^{n/(n-1)} - 1}{a^{1/(n-1)} - 1} > c.$$

Proof. Since the left-hand member of (12) can be written as a polynomial in $a^{1/(n-1)}$, with positive coefficients, it is an increasing function of a . Hence it is sufficient to prove that

$$(12') \quad \frac{2}{n} \frac{c^{n/(n-1)} - 1}{c^{1/(n-1)} - 1} \geq c \quad \text{for } n \geq 5.$$

Put $y_n = c^{1/(n-1)} - 1$. Then $n = 1 + \log c / \log (y_n + 1)$ and so the inequality (12') takes the form

$$(12'') \quad \frac{2 \log (y_n + 1)}{\log c + \log (y_n + 1)} \left(c + \frac{c-1}{y_n} \right) \geq c \quad \text{for } n \geq 5.$$

The quantity y_n is positive and tends to zero for $n \rightarrow \infty$. We now consider the function

$$f(y) = \frac{2 \log (y+1)}{\log c + \log (y+1)} \left(c + \frac{c-1}{y} \right) \quad (y > 0).$$

Clearly, by (2),

$$\lim_{y \rightarrow +0} f(y) = \frac{2}{\log c} (c-1) = c.$$

We shall prove that $f(y)$ is a steadily increasing function of y for $0 < y < 0.5$.

Differentiating $f(y)$ and using (2) we get

$$\begin{aligned} f'(y) &= -\frac{c-1}{y^2} \cdot \frac{2 \log (y+1)}{\log c + \log (y+1)} + \left(c + \frac{c-1}{y} \right) \cdot \frac{2 \log c}{(y+1)[\log c + \log (y+1)]^2} \\ &= \frac{2(c-1)}{y^2(y+1)[\log c + \log (y+1)]^2} \psi(y), \end{aligned}$$

with

$$\psi(y) = -(y+1) \log (y+1) [\log c + \log (y+1)] + 2y^2 + \left(2 - \frac{2}{c} \right) y.$$

We have $\psi(0) = 0$. Further,

$$\begin{aligned} \psi'(y) &= -[\log^2 (y+1) + (2 + \log c) \log (y+1) + \log c] + 4y + 2 - \frac{2}{c}, \\ \psi''(y) &= -\frac{1}{y+1} [2 \log (y+1) + 2 + \log c] + 4 \quad (y \geq 0). \end{aligned}$$

Clearly $\psi'(0) = -\log c + 2 - \frac{2}{c} = 0$. Next,

$$\begin{aligned} 2 \log (y+1) + 2 + \log c &< 0.4 + 2 + 1.6 = 4 \quad \text{for } 0 < y < 0.2, \\ 2 \log (y+1) + 2 + \log c &< 1 + 2 + 1.6 < 4.8 \quad \text{for } 0.2 \leq y < 0.5. \end{aligned}$$

Hence $\psi''(y) > 0$, and so $\psi'(y) > 0$, for $0 < y < 0.5$. Then also $\psi(y) > 0$, and so $f'(y) > 0$ for $0 < y < 0.5$.

It follows that $f(y) > c$ for $0 < y < 0.5$. Since $0 < y_n < 0.5$ for $n \geq 5$, this proves (12'') and so proves the lemma.

Lemma 7. *If a is a number with $c < a < 6$, and $n \geq 6$, then*

$$(13) \quad \frac{94}{100} a < \frac{2}{n} \frac{a^{n/(n-1)} - 1}{a^{1/(n-1)} - 1} < 6.$$

Proof. Put $n-1 = m$, so that $m \geq 5$. Write $a_m = \frac{2}{m+1} \frac{a^{(m+1)/m} - 1}{a^{1/m} - 1}$. Since a_m is a steadily increasing function of a , we have

$$\begin{aligned} a_m &< \frac{2}{m+1} \frac{6^{(m+1)/m} - 1}{6^{1/m} - 1} = \frac{2}{m+1} \left(6 + \frac{5}{6^{1/m} - 1} \right) \\ &< \frac{2}{m+1} \left(6 + \frac{5}{m^{-1} \log 6 + \frac{1}{2} m^{-2} \log^2 6} \right) = \frac{6}{m+1} \left(2 + \frac{10}{6 \log 6} \cdot \frac{m}{1 + \frac{1}{2} m^{-1} \log 6} \right). \end{aligned}$$

Hence $a_m < 6$, since

$$\frac{10}{6 \log 6} = 0.9416 \dots < (1 - m^{-1}) (1 + \frac{1}{2} m^{-1} \log 6) \quad \text{for } m \geq 5.$$

Further, since $\frac{a-1}{a \log a}$ is steadily decreasing for $c < a$,

$$\begin{aligned} \frac{a_m}{a} &= \frac{2}{m+1} \left(1 + \frac{a-1}{a(a^{1/m}-1)} \right) > \frac{2}{m+1} \left(1 + m \cdot \frac{a-1}{a \log a} \left(1 - \frac{1}{2m} \log a \right) \right) \\ &> \frac{2}{m+1} \left(1 + \frac{5m}{6 \log 6} - \frac{a-1}{2a} \right) > \frac{10}{6 \log 6} > \frac{94}{100}. \end{aligned}$$

This proves (13).

Proof of theorem 1. By (2), $c_2 \geq 2\sqrt{3} > 1$. Suppose $n \geq 3$, and that $c_{n-1} > 1$. Let K be a convex body in R_n , of volume V , symmetric about O . Consider the section of K by $x_n = 0$. It is an $(n-1)$ -dimensional convex body, symmetric about O , of volume $V(0)$. By the definition of c_{n-1} , the critical determinant of this body is at most equal to $V(0)/c_{n-1}$, and so there exists an $(n-1)$ -dimensional lattice \mathcal{L} in the plane $x_n = 0$, of determinant $d(\mathcal{L}) = V(0)/c_{n-1}$, which has no point (except O) in the interior of K .

Put $\beta = c_{n-1}^{-1}$. Since we assumed that $c_{n-1} > 1$, we have $0 < \beta < 1$. For this β let α be defined by (7). Then, from lemma 2,

$$\sum_{t=1}^{\infty} V(\alpha t) \leq \beta V(0) = V(0)/c_{n-1} = d(\mathcal{L}).$$

In virtue of the lemmas 1 and 3 there now exists a point g of the form $g = (g_1, \dots, g_{n-1}, \alpha)$, such that the lattice Δ generated by \mathcal{L} and g is admissible for K . It follows that the critical determinant of K , Δ say, is at most equal to $\alpha d(\mathcal{L})$. Hence

$$V/\Delta \geq \frac{c_{n-1}V}{\alpha V(0)} = \frac{2c_{n-1}}{n} \frac{1 - c_{n-1}^{-n/(n-1)}}{1 - c_{n-1}^{-1/(n-1)}} = \frac{2}{n} \cdot \frac{c_{n-1}^{n/(n-1)} - 1}{c_{n-1}^{1/(n-1)} - 1}.$$

From the arbitrariness of K it then follows that

$$(14) \quad c_n \geq \frac{2}{n} \frac{c_{n-1}^{n/(n-1)} - 1}{c_{n-1}^{1/(n-1)} - 1}.$$

From (14) it follows that $c_n > 1$, since we assumed that $c_{n-1} > 1$. Then it follows, by induction on n , that (14) holds for all $n \geq 3$. Using the relation $c_2 \geq 2\sqrt{3}$ and applying (14) with $n = 3, 4, 5$ we find that $c_5 > c = 4.921\dots$ Hence, by (14) and lemma 6, we have $c_n > c$ for $n \geq 5$. This proves the theorem.

Proof of theorem 2. We define numbers d_2, d_3, \dots as follows:

$$(15) \quad d_2 = 3, \quad d_n = \frac{2}{n} \frac{d_{n-1}^{n/(n-1)} - 1}{d_{n-1}^{1/(n-1)} - 1} \quad \text{for } n \geq 3.$$

Clearly $d_n > 1$ for all n . By induction on n , we shall prove the following assertion:

Let b be any number $\geq \frac{200}{94}$. Then there exists a lattice Δ with the following properties:

1. the lattice A is admissible for K
2. the determinant of A is equal to V/d_n
3. A has a basis contained in the cube

$$W: |x_i| < b(V/\kappa_n)^{1/n} \quad (i = 1, 2, \dots, n)$$

4. for each point x of the space there exists a point $y \in A$, such that $x - y \in W$.

In proving this assertion it is no loss of generality to suppose that $V = \kappa_n$.

First consider the case $n=2$ and suppose that $V = \kappa_2 = \pi$. Then there exists a point x' on the boundary of K at distance 1 from O . It is no loss of generality to suppose that x' is the point $(1, 0)$. Then the line $x_2 = \pi/3$ intersects K in a segment of length ≤ 1 , since otherwise the volume of K would be greater than π . For a similar reason the line $x_2 = 2\pi/3$ does not intersect K . Consequently, there exists a K -admissible lattice A , generated by the point $(1, 0)$ and a point of the form $(a, \pi/3)$. Here we may take a such that $|a| \leq \frac{1}{2}$. Hence A has a basis contained in the square

$$|x_i| \leq \pi/3 < b \quad (i = 1, 2).$$

Next, for each point x of the plane there exist integers u and v , such that $x - u \cdot (a, \pi/3) - v \cdot (1, 0)$ is contained in the square $|y_i| < b \quad (i = 1, 2)$. Finally, A has determinant $d(A) = \pi/3$. This proves the assertion in the case $n=2$.

Now let $n \geq 3$ and suppose that the assertion is true, with n replaced by $n-1$. Let K be a centrally symmetric convex in n dimensions, of volume κ_n . As we already remarked above (see the relation (9')), it follows from the lemmas 4 and 5 that there exists an $(n-1)$ -dimensional plane Π through O , such that the $(n-1)$ -dimensional volume of $K \cap \Pi$ is comprised between $\frac{1}{2}\kappa_n$ and κ_{n-1} . It is no loss of generality to suppose that Π is the plane $x_n = 0$, so that

$$(16) \quad \frac{1}{2}\kappa_n \leq V(0) \leq \kappa_{n-1}.$$

Hence $V(0)/\kappa_{n-1} \leq 1$. So, according to the induction hypothesis, there exists an $(n-1)$ -dimensional lattice \mathcal{L} in the plane $x_n = 0$ with the following properties:

- 1) \mathcal{L} has no point (except O) in the interior of K
- 2) \mathcal{L} has determinant $d(\mathcal{L}) = V(0)/d_{n-1}$
- 3) \mathcal{L} has a basis contained in the cube

$$|x_i| < b \quad (i = 1, 2, \dots, n-1), \quad x_n = 0$$

- 4) for each point x in the space $x_n = 0$ there exists a point $y \in \mathcal{L}$, such that $|x_i - y_i| < b$ for $i = 1, 2, \dots, n-1$.

We now apply the lemmas 2 and 3. Take $\beta = 1/d_{n-1}$. Then $0 < \beta < 1$. With this value of β , let α be defined by (6). Then, by 2) and lemma 2,

$$\sum_{t=1}^{\infty} V(\alpha t) \leq d(\mathcal{L}).$$

Hence, in virtue of 1) and lemma 3, there exists a point g of the form $g = (g_1, \dots, g_{n-1}, \alpha)$, such that the lattice Λ generated by \mathcal{L} and g is admissible for K . We shall prove that this lattice possesses also the properties 2, 3, 4.

Using 2) and the definitions of α and d_n we find that the determinant of Λ is given by

$$d(\Lambda) = \alpha d(\mathcal{L}) = \frac{V}{2d_{n-1}} \frac{n(1-d_{n-1}^{-1/(n-1)})}{1-d_{n-1}^{-n/(n-1)}} = \frac{V}{2} \frac{n(d_{n-1}^{1/(n-1)} - 1)}{d_{n-1}^{n/(n-1)} - 1} = V/d_n.$$

Next, in virtue of 4), there exists a point $y \in \mathcal{L}$, such that $|g_i - y_i| < b$ for $i = 1, 2, \dots, n-1$. Put $g - y = x$. Then Λ is generated by \mathcal{L} and x . Further $|x_i| < b$ for $i = 1, 2, \dots, n-1$ and $x_n = \alpha$. We prove that $\alpha < b$. We have $\alpha = \frac{V}{V(0)} \frac{d_{n-1}}{d_n}$. On account of (16), $\frac{V}{V(0)} \leq 2$. From (15) one easily finds that $d_2 < d_3 < d_4 < d_5 < d_6 < 6$ and $d_6 > c$. Then, by lemma 6, $d_n > c$ for $n > 6$. Hence, by lemma 7, $\frac{94}{100} d_{n-1} < d_n < 6$ for $n \geq 7$. Hence $d_{n-1}/d_n < \frac{100}{94}$ for $n = 3, 4, \dots$. This shows that $\alpha < b$. It follows that Λ has a basis contained in the cube $|x_i| < b$ ($i = 1, 2, \dots, n$). Finally, let x be an arbitrary point. There exists an integer u_n , such that $|x_n - u_n \alpha| < b$. In virtue of 4), there further exists a point $y \in \mathcal{L}$, such that $|x_i - u_n g_i - y_i| < b$ for $i = 1, 2, \dots, n-1$. Hence the point $x - u_n g - y$ is contained in the cube $|x_i| < b$ ($i = 1, 2, \dots, n$).

This proves that Λ possesses the properties 1, 2, 3, 4. By induction on n , the truth of the assertion follows. Calculation gives $d_3 > 3.82$, $d_4 > 4.41$, $d_5 > 4.80$, whereas $d_n > c$ for $n \geq 6$. From this the theorem follows.

Mathematisch Centrum, Amsterdam

REFERENCES

1. Hlawka, E., Zur Geometrie der Zahlen, Math. Z. **49**, 285-312 (1943).
2. Rogers, C. A., Existence theorems in the geometry of numbers, Ann. of Math. (2) **48**, 994-1002 (1947).
3. ———, The number of lattice points in a star body, J. London Math. Soc. **26**, 307-310 (1951).
4. Malýšev, A. V., On the Minkowski-Hlawka theorem concerning a star body, Uspehi Matem. Nauk. (N.S.) **7**, no 2 (48), 168-171 (1952). (Russian).
5. Cassels, J. W. S., A short proof of the Minkowski-Hlawka theorem, Proc. Cambridge Philos. Soc. **49**, 165-166 (1953).
6. Siegel, C. L., A mean value theorem in geometry of numbers, Ann. of Math. (2) **46**, 340-347 (1945).
7. Weil, A., Sur quelques résultats de Siegel, Summa Brasiliensis Math. **1**, fasc. 4, 21-39 (1946).
8. Mahler, K., The theorem of Minkowski-Hlawka, Duke Math. J. **13**, 611-621 (1946).
9. Davenport, H. and C. A. Rogers, Hlawka's theorem in the geometry of numbers, Duke Math. J. **14**, 367-375 (1947).
10. Mahler, K., On lattice points in n -dimensional star bodies, I. Existence theorems, Proc. Royal Soc. A **187**, 151-187 (1947).
11. Bonnesen, T. and W. Fenchel, Theorie der konvexen Körper (Chelsea, 1947).