

STICHTING
MATHEMATISCH CENTRUM
2e BOERHAAVESTRAAT 49
AMSTERDAM


AFDELING ZUIVERE WISKUNDE

ZW 1963-005

Topological semigroups II

by

Aida Paalman - de Miranda



May 1963

BIBLIOTHEEK MATHEMATISCH CENTRUM
AMSTERDAM

CHAPTER II

Semigroups with zero and identity.

§ 1. Semigroups with zero.

Let S be a mob with 0 , and a an element of S . If $a^n \rightarrow 0$, i.e. if for every neighbourhood U of 0 there exists an integer n_0 , such that $a^n \in U$ if $n \geq n_0$, then a is termed a nilpotent element.

We denote by N the set of all nilpotent elements of S . An ideal (right, left) A of S with the property $A^n \rightarrow 0$ is called a nilpotent ideal.

A nil-ideal A is an ideal consisting entirely of nilpotent elements.

Then it is clear that every nilpotent ideal is a nil-ideal, and that the join of a family of (right, left) nilideals is again a (right, left) nilideal of S .

Example: Let S be the unit interval with the usual multiplication. Then $I = [0, 1]$ is an ideal consisting entirely of nilpotent elements.

I is not a nilpotent ideal, since $I^n = I$ for all n .

Lemma 1: Every right (left) nilideal of S is contained in some nilideal of S .

Proof: Let A be a right nilideal of S . Then SA is an ideal of S . Suppose $x = sa \in SA$, and let U be any neighbourhood of 0 . Then there exists a neighbourhood V of 0 such that $s \vee a \in U$. As A is a right nilideal of S , $as \in A$, and $(as)^n \in V$ for $n \geq n_0$. Hence if $m \geq n_0 + 1$ we have $(sa)^m = s(as)^{m-1} a \in s \vee a \in U$. Therefore SA is a nilideal of S , and hence $A \cup SA$ is a nilideal of S containing A .

Definition 1: The join R of all nil-ideals of a mob S with zero is called the radical of S .

Lemma 1 implies that R is a nil-ideal, which contains every right and every left nilideal of S .

Hence R is the maximal right and the maximal left nil-ideal.

If $S = R$ i.e if S consists only of nilpotent elements, then S is called a nil - semigroup.

Let $a \in S$, then we shall denote by $\Gamma(a)$, the closure of the set $\{a^n\}_{n=1}^{\infty}$. $\Gamma(a) = \{a^n\}_{n=1}^{\infty}$.

Lemma 2: Let S be a mob and let A be a compact part of S such that $Ax \subset A$, with $\Gamma(x)$ compact.

Then $\bigcap_{n=1}^{\infty} Ax^n = Ae$, with $e = e^2 \in \Gamma(x)$.

Proof: Let $p \in \bigcap_{n=1}^{\infty} Ax^n$

Then $p = a_1x = a_2x^2 = \dots$

Hence from §1 lemma 2 it follows that there is an element $a \in \{a_1\}^-$ such that $p = ae$, where $e = e^2 \in \Gamma(x)$ (see §1 th. 4).

This implies $\bigcap_{n=1}^{\infty} Ax^n \subset Ae$.

Now let $a_1 \notin Ax^k$. Then we can find a neighbourhood V of e such that $a_1V \cap Ax^k = \emptyset$.

But since $p \in \Gamma(x)$, there is a $k_0 \gg k$ such that $x^{k_0} \in V$ and hence $a_1x^{k_0} \notin Ax^k$.

This is a contradiction, since $Ax \subset A$ implies $Ax^{k_0} \subset Ax^k$.

Hence $Ae \subset Ax^k \Rightarrow \bigcap_{n=1}^{\infty} Ax^n = Ae$.

Theorem 1: Let S be an element - wise compact mob with zero (i.e for every a , $\Gamma(a)$ is compact).

Then every (right, left) ideal of S is either a nil-ideal or contains non - zero idempotents.

Proof: Let a be a non-nilpotent element of the ideal A .

Then the identity e of the group $D = \bigcap_{n=1}^{\infty} \{a^n\}_{1 \leq n}^-$ is not equal to zero.

Furthermore $a \in D$, and $a \in A \cap D \subseteq A$, since A is an ideal. Hence $D \cap A \neq \emptyset$, so that $D \subseteq A$, since no group can properly contain an ideal. Thus $e \in A$.

Theorem 2: Let e be a non-zero idempotent of the compact monoid S with zero.

Then these are equivalent.

- 1) $eSe \setminus \{0\}$ is a group
- 2) e is primitive
- 3) eSe is a minimal non-nil left ideal
- 4) eSe is a minimal non-nil ideal
- 5) each idempotent of eSe is primitive.

Proof: $(1 \rightarrow 2)$: If $eSe \setminus \{0\}$ is a group, then e is the only idempotent in $eSe \setminus \{0\}$, since no idempotent $\neq 0$, can be nilpotent.

Hence e is primitive

$(2 \rightarrow 3)$: Let L be a non-nil left ideal $L \subseteq eSe$.

Then by theorem 1 there is an idempotent $f \in L$, $f \neq 0$.

Since $f \in eSe$, we have $fe = f$, and $(ef)(ef) = ef$.

Thus ef is an idempotent $\neq 0$ and $ef \in eSe$.

Hence since e is primitive $ef = e$.

This implies that $ef = e \in eL \subseteq L \Rightarrow L = eSe$.

$(3 \rightarrow 4)$ Let I be a non-nil ideal $I \subseteq eSe$.

Then there exists an idempotent $f \in I$, $f \neq 0$, and elements $a, b \in S$, such that $aeb = f$.

We can choose b such that $bf = b$.

Let $g = bae$. Then $g^2 = baebae = bfae = g$.

Furthermore $g \neq 0$, since otherwise $0 = gb = baeb = bf = b$.

Now $g \in eSe$ and $g \in SfS$.

Hence by (3) $eSe = Sg \subseteq SfS$, and we conclude $eSe = SfS = I$.

$(4 \rightarrow 5)$ Let f be a non-zero idempotent of eSe , and let $g = g^2 \neq 0 \in fSf$.

Since $f, g \in eSe$, we have $SgS = SfS = eSe$ and $f \in SgS$. Hence $f = agb$,

and we may assume $ag = a$, $gb = b$.

Since $gf = fg = g$, this implies $afb = agfb = agb = f$.

Hence $f = a^n g b^n$.

It follows from §1 lemma 2 that there is an idempotent

$g^* \in P(a) \in Sg$ and $b' \in P(b)$ such that $f = g^* g b'$.

We note that $g^* g = g^*$, hence $g^* f = f = g^* g f = g^*$, and

$f = g^* = g^* g = fg = g$.

(5 \Rightarrow 1) Since every idempotent in SeS is primitive, e is primitive and hence $Se = L$ is a minimal non-nil left ideal.

Now let $a \in eSe \setminus N$, then $a \in \{Se \cap eS\} \setminus N$.

Since L is minimal. $a = ea \in La = L$.

Hence there is $\bar{a} \in L$ such that $\bar{a}a = e$.

Let $e\bar{a} = a'$, then $a' \in eSe$ and $a'a = e$.

$(aa')(aa') = aea' = aa'$. Hence aa' is an idempotent and

$aa' \in eSe \setminus N$. Since e is primitive $aa' = e$.

So we can find for every $a \in eSe \setminus N$ an element $a' \in eSe$ such that $aa' = e = a'a$.

This implies that $eSe \setminus N$ is a group, since $a' \notin N$.

For if $a' \in N$, then $\bigcap_{n=1}^{\infty} S(a')^n = S \cdot 0 = 0$ by lemma 2.

This is in contradiction with $aa' = a^2(a')^2 = a^n(a')^n = e$.

Definition: A mob S with zero is said to be an N -semigroup if its nilpotent elements form an open set.

Lemma 3: Let S be a mob with zero, and let $a \in S$.

If a^n is nilpotent for some $n \geq 0$, then a itself is a nilpotent element.

Proof: Let U be an arbitrary neighbourhood of 0 , then since

$a^j 0 = 0$, there is a neighbourhood V of 0 , such that

$a^j V \subset U$ ($j = 1, 2, \dots, n$).

Since a^n is nilpotent there exists an integer $k_0 \geq 0$ such that $(a^n)^k \in V$ for $k \geq k_0$.

Thus $a^j a^{nk} = a^{nk+j} \in U$. $j = 1, 2, \dots, n$ $k \geq k_0$.

This implies that for $N \geq nk_0$ $a^N \in U$.

Hence a is nilpotent.

Theorem 3: If a mob S with O has a neighbourhood U of O , which consists entirely of nilpotent elements, then S is an N -semigroup.

Proof: Let $p \in N$, then there is an n such that $p^n \in U$.

Therefore there is a neighbourhood V of p , such that $V^n \subset U$.

Hence every point of V^n is nilpotent.

Lemma 3 then implies that $V \subset N$.

Theorem 4: A locally compact mob S with O having a neighbourhood U of O which contains no non-zero idempotents is an N -semigroup.

Proof: Since S is locally compact and Hausdorff. S is regular, and we can find a neighbourhood W of O , such that $\bar{W} \subset U$, and \bar{W} is compact.

The continuity of multiplication and the compactness of \bar{W} imply, that there is a neighbourhood V of O , with $V \bar{W} \subset W$ $V \subset W$.

Hence $V^2 \subset V$. $\bar{W} \subset W$, and $V^n \subset W$.

The set $A = \bigcup_{i=0}^{\infty} V^i$ is a mob contained in W .

Therefore \bar{A} is a compact mob contained in U .

Since \bar{A} contains no non-zero idempotents \bar{A} is a nil-semigroup (theorem 1).

Hence V consists entirely of nilpotent elements, and by theorem 3 S is an N -semigroup.

Corollary: A locally compact semigroup with O , which is not an N -semigroup contains a set of non-zero idempotents with clusterpoint O .

Theorem 5: The radical of a compact N-semigroup is open.

Proof: Let $a \in R$, then for every $s \in S$ $sa \in R \subset N$.

Since N is open and S compact, there exists a neighbourhood V of a such that $SV \subset N$, $V \subset N$.

Since $V \cup SV$ is a left nil-ideal, $V \cup SV \subset R$.

Hence $V \subset R$ and R is open.

Theorem 6: Let S be a compact N-semigroup which is not a nil-semigroup.

Then any non-nilideal I of S contains a minimal non-nil ideal I^* , such that I^*/R^* is completely simple, where $R^* = I^* \cap R$ is the radical of I^* .

Proof: Since I is a non-nilideal of S , I contains non-zero idempotents.

Let $E^* = E - \{0\}$. Then E^* is closed, since N is open and E is closed.

Let $E_\lambda = E^* \cap Se_\lambda S$, $e_\lambda \in E^* \cap I$.

Then E_λ is closed and non-empty.

Suppose E_ν is a minimal member of $\{E_\lambda\}$. E_ν exists since S is compact.

We shall now prove that e_ν is a primitive idempotent.

Suppose $0 \neq f = f^2 \in e_\nu Se_\nu \Rightarrow f \in I$. Then $SfS \subset Se_\nu S$.

Since E_ν is minimal;

$E^* \cap SfS = E^* \cap Se_\nu S$. Hence $e_\nu = s_1 f s_2$, with $e_\nu s_1 = s_1, s_1 f = s_1$.

$$\begin{aligned} s_1^n f s_2^n &= s_1^{n-1} s_1 f s_2 s_2^{n-1} = s_1^{n-1} f s_1 f s_2 s_2^{n-1} = s_1^{n-1} f e_\nu s_2^{n-1} = \\ &= s_1^{n-1} f s_2^{n-1} \end{aligned}$$

Hence $s_1^n e_\nu s_2^n = e_\nu$.

Thus there is an idempotent $g \in \Gamma(s_1)$ and an element $s \in \Gamma(s_2)$ so that $ge_\nu s = e_\nu$.

We note that since $\Gamma(s_1) \in SfS$ $gf = g$.

Hence $e_\nu = ge_\nu = gfe_\nu = gf = g \Rightarrow f = e_\nu f = gf = g = e_\nu$.

Thus e_ν is a non-zero primitive idempotent.

Theorem 2 then implies that $Se_\nu S = I^* \subset I$ is a minimal non nil-ideal.

Now we shall prove that $R^* = I^* \cap R$.

Since $I^* \cap R$ is a nil-ideal of I^* we have $I^* \cap R \subset R^*$.

Furthermore $SR^*S \subset SI^*S \subset I^*$.

If $SR^*S = I^*$, then $I^*SR^*SI^* = I^{*3} = I^*$, and so $I^* = I^*SR^*SI^* \subset I^*R^*I^* \subset R^*$. This contradicts the fact that I^* is a non nil-ideal.

Hence SR^*S is an ideal of S properly contained in I^* .

This implies that SR^*S must be a nil-ideal i.e. $SR^*S \subset R^* \Rightarrow R^*$ is a nil-ideal of $S \Rightarrow R^* \subset I^* \cap R$.

Since R^* is a maximal proper ideal of I^* , §3 th.3 implies that I^* / R^* is completely simple.

Corollary: Let S be a compact mob with zero; then S contains a non-zero primitive idempotent if and only if there is a non-zero idempotent e with $(eSe) \setminus N$ closed.

Proof: If $e = e^2 \neq 0$, e primitive $eSe \setminus N$ is a maximal subgroup. (th.2). On the other hand if $(eSe) \setminus N$ is closed and $e \neq 0$, then $eSe \setminus N$ is the set of nilpotent elements of eSe , and $eSe \cap N$ is open in eSe .

We conclude from theorem 6 that eSe contains a non-zero primitive idempotent. Hence so does S .

Theorem 7: Let e be a non-zero primitive idempotent of the compact mob S with zero. Then $Se \setminus N$ and $(Se) \cap N$ are submobs and $Se \setminus N$ is the disjoint union of the maximal groups $e_\alpha Se_\alpha \setminus N$ where e_α runs over the non-zero idempotents of Se .

Proof: Suppose $a, b \in Se \setminus N$, then $a^n, b^n \in Se \setminus N$. Let $ab \in N$. Then since Se is a minimal non-nil left ideal, we know that $Sa = Sb = Se \Rightarrow Sa^n = Sb^n = Se$. Hence $Sab = Sb^2 = Se \Rightarrow S(ab)^n = Se$.

Thus $Se = \bigcap_n S(ab)^n = S0 = 0$ (lemma 2).

This is a contradiction with $e \neq 0$.

Suppose now $a_1 b \in Se \cap N$ and $ab \notin N$.

Then $(ab)^2 \notin N$ and hence $Sab = Se$, since Se is a minimal non-nil left ideal.

Since $a \in Se$, we have $Sa \subset Se = Sab$.

Hence $Sa \subset Sab \subset Sab^2 \subset Sab^3 \subset \dots$

But since $ab^n \in Se$, $Sab^n = Se$.

This implies that $Se = \bigcap_n Sab^n = Sa \cdot 0 = 0$, a contradiction.

Finally let $a \in Se \setminus N$. Then $Sa = Se$.

Choose an idempotent f in $\Gamma(a)$; then $Sf = Se = Sa$, and f is a right unit for Se .

Let D be the subgroup of S contained in $\Gamma(a)$. Then D is an ideal of $\Gamma(a)$ (§ 1 th.4). Hence $\Gamma(a) f \subset D \Rightarrow \Gamma(a) = D$ and $\Gamma(a)$ is a group. Thus $Se \setminus N$ is the union of groups.

For any $e_\alpha = e_\alpha^2 \neq 0$, $e_\alpha \in Se$, $Se_\alpha = Se$, so that e_α is primitive and $e_\alpha Se_\alpha \setminus N$ is a group.

Now the maximal group containing e_α is contained in $e_\alpha Se_\alpha$, moreover since any group which meets N must be zero, we conclude that $e_\alpha Se_\alpha \setminus N$ is a maximal group.

§ 2. 0 - simple semigroups.

As in Ch. 1 § 3 we call a semigroup S simple if it does not contain a proper non-zero ideal.

By a 0-simple semigroup we mean a simple semigroup containing a zero element.

A completely 0-simple semigroup is a completely simple semigroup with a zero element.

If S is completely 0-simple then S contains a non-zero idempotent and this implies that S cannot be a nil-semigroup.

On the other hand if S is not a nil-semigroup and S is 0-simple, then every right or left nilideal of S is the zero ideal $\{0\}$, since (§ 1 lemma 1) every right (left) nilideal of S is contained in some nilideal of S .

We shall call a (left, right) ideal I of a mob S with zero 0-minimal if $I \neq \{0\}$ and $\{0\}$ is the only (left, right) ideal of S properly contained in I .

Hence every minimal non-nil left ideal of a 0-simple mob is a 0-minimal left ideal.

Lemma 1: Let L be a 0-minimal left ideal of a 0-simple mob S and let $a \in L \setminus 0$. Then $Sa = L$.

Proof: Since Sa is a left ideal of S contained in L , it follows that $Sa = 0$ or $Sa = L$.

If $Sa = 0$, then $SaS = 0$, in contradiction with $SaS = S$.

If S is an element-wise compact mob with zero, then every non-nil (left, right) ideal of S , contains a non-zero idempotent.

So in this case if L is a minimal non-nil left ideal of S , then there is an idempotent $e \in L$, with $Se = L$.

Lemma 2: Let L be a 0-minimal left ideal of a 0-simple mob S , and let $s \in S$. Then Ls is either $\{0\}$ or a 0-minimal left ideal of S .

Proof: Assume $Ls \neq 0$. Evidently Ls is a left ideal of S .

Now let L_0 be a left ideal of S contained in Ls . $L_0 \subset Ls$.

Let A be the set of all $a \in L$ with $as \in L_0$.

Then $As = L_0$, and $A \subset L$.

Furthermore $SA \subset SL_0 \subset L_0$ and $SA \subset SL \subset L$.

Hence $SA \subset A$ and A is a left ideal of S .

From the minimality of L , either $A = 0$ or $A = L$, and we have correspondingly $L_0 = 0$ or $L_0 = Ls$.

Theorem 1: Let S be a compact 0-simple mob. Then S is the union of all minimal (i.e minimal non-nil) left ideals of S .

Proof: Since S is compact, S is completely 0-simple and hence contains a non-zero primitive idempotent e .

From §1 th.2 it then follows that Se is a minimal non-nil left ideal, and hence a 0-minimal left ideal.

Now let A be the union of all the 0-minimal left ideals of S .

Clearly A is a left ideal of S and $A \neq \{0\}$.

Now we show that A is, also a right ideal.

Let $a \in A$ and $s \in S$. Then $a \in L$ for some 0-minimal left ideal L of S .

By lemma 2 $Ls = 0$ or Ls is a 0-minimal left ideal.

Hence $Ls \subset A$ and $as \in A$.

Thus A is a non-zero ideal of S , whence $A = S$.

An analogous result holds for 0-minimal right ideals.

Lemma 3: Let L and R be 0-minimal left and right ideals of a 0-simple mob, such that $LR \neq 0$.

Then $RL = R \cap L$ is a group with zero and the identity e of $RL \setminus \{0\}$ is a primitive idempotent of S .

Proof: Since LR is a non-zero ideal of S , we must have

$LR = S$. Furthermore $RL \neq 0$, since $S = S^2 = LRLR$.

Now let $a \in RL \setminus 0$, then $a \in L \setminus 0$ and $a \in R \setminus 0$, and hence $Sa \equiv L$ (lemma 1), and $aR = 0$ or $aR = R$.

Since $S = LR = SaR$, it follows that $aR \neq 0$.

Consequently $aRL = RL$.

In the same way we can prove that $RLa = RL$.

From this we conclude that RL is a group with zero.

Now let e be the identity of RL .

Then since $R = eS$ and $L = Se$, we have $R \cap L = eS \cap Se = eSe$ and $RL = eSSe = eSe$.

Since eSe is a group with zero, e is primitive.

Theorem 2: Let S be a compact 0-simple mob and let e and f be non-zero primitive idempotents of S .

Then the maximal subgroups $H(e)$ and $H(f)$ containing e and f respectively are topological isomorphic compact groups.

Proof: Since Se and Sf are 0-minimal left ideals and eS and fS 0-minimal right ideals (§ 1 th.2) it follows from lemma 3 that $eSe \setminus \{0\}$ and $fSf \setminus \{0\}$ are groups.

Since $H(e) \subset eSe \setminus \{0\}$ we have $H(e) = eSe \setminus \{0\}$, $H(f) = fSf \setminus \{0\}$. Now $eSSf \neq 0$, since $eSSfS = eS^2 = eS$.

Hence $eS \cap Sf \neq 0$.

Let $a \neq 0 \in eS \cap Sf$. Then $ea = a = af$.

Since $eS = aS$ and $Sf = Sa$ (lemma 1), there exists a_1 and $a_2 \in S$ such that $e = aa_1$ $f = a_2a$.

Now let $b = fa_1e$, then $b \neq 0$ and

$ab = afa_1e = aa_1e = ee = e$; $ba = fba = a_2aba = a_2ea = f$.

Furthermore $bS = fS$, $Sb = Se$.

We now proof that the mappings $\varphi : x \rightarrow bxa$ and $\psi : y \rightarrow ayb$ are mutually inverse one-to-one mappings of $H(e)$ and $H(f)$ upon each other.

For let $x \in H(e)$ then $bxa \in bS \cap Sa = fS \cap Sf = H(f) \cup \{0\}$.

Similarly $y \in H(f)$ implies $ayb \in aS \cap Sb = eS \cap Se = H(e) \cup \{0\}$.

And if $x \in H(e)$ $a(bxa)b = exe = x$.

φ is an isomorphism since $(bx_1a)(bx_2a) = bx_1ex_2a = bx_1x_2a$.

Since φ is continuous and one-to-one φ is topological.

Corollary: Let S be a compact 0-simple mob.

Then S is the disjoint union of isomorphic compact groups $H(e)$ and of sets A_α with the property $A_\alpha^2 = 0$.

Corollary: Let S be a commutative compact 0-simple mob.

Then S is a group with zero.

Proof: By lemma 3 we have $S^2 = S \cap S = S$ is a group with zero since S is both a 0-minimal left and right ideal.

Theorem 3: Let J be a maximal proper ideal of the compact mob S .

Then the following are equivalent.

- 1°) $S-J$ is the disjoint union of groups.
- 2°) for each element of $S-J$, there exists a unit element
- 3°) $a \in S-J$ implies $a^2 \in S-J$
- 4°) J is a completely prime ideal
- 5°) $S-J$ contains an idempotent, and the product of two idempotents of $S-J$ lies in $S-J$.

Proof:

(1) clearly implies (2).

(2) \rightarrow (3). Let $a \in S-J$ and $ax = xa = a$.

Then $ae = ea = a$, with $e = e^2 \in \Gamma(x)$ and $e \in S-J$.

Hence since $S-J \cup \{0\} = \bigcup H(e_\alpha) \cup \bigcup A_\alpha$, we have $a \in H(e)$ which implies $a^2 \in H(e) \Rightarrow a^2 \in S-J$.

(3) \rightarrow (4) Let $a, b \in S-J$ and suppose $ab \in J$.

Then $I = \{x \mid x \in S \text{ } xb \in J\}$ is a left ideal with $I \supset J$.

Now let $x \in I$, $xs \notin I$, then $xs b \notin J$, and hence $xs b x s b \notin J \Rightarrow bx \notin J \Rightarrow b x b x \notin J \Rightarrow x b \notin J$ a contradiction.

Since I is an ideal containing J , we have $I = S \Rightarrow b^2 \in J$ a contradiction.

(4) \rightarrow (5) This follows from the fact that $J = J_0(S-e)$.

(5) \rightarrow (1) Since $e \in S-J$, we have S/J completely simple and $S/J = \bigcup H(e_\alpha) \cup \bigcup A_\beta$.

Now let $a \neq 0 \in A_\beta$, then $a \in Se$ and $a \in fS$. with $SefS = 0$, or else it would follow from lemma 3 that $a \in SenfS = H(e_\alpha) \cup \{0\}$. Since $ef \notin J$, we have however $SefS \neq 0$, a contradiction. Hence $A_\beta = \emptyset$ and $S-J = \bigcup H(e_\alpha)$.

From theorem 3 it follows that $S-J$ is a group if and only $S-J$ contains a unique idempotent.

§ 3. Connected semigroups

Lemma 1: If S is connected, then each minimal (left, right) ideal of S is connected.

Proof:

Let L be a minimal left ideal of S , then for any $a \in L$, $Sa = L$ and hence L is connected.

If K is the minimal ideal of S , then $K = SaS$ for each $a \in K$. Hence $K = \bigcup_{s_\alpha \in S} Sas_\alpha$: Since each Sas_α is connected and meets the connected set aaS it follows that K is connected.

Lemma 2: If S is connected, then each ideal of S is connected, provided S has a left or right unit.

Proof:

Let I be an ideal of S . Then $I = \bigcup_{x \in I} Sx$ if e is a left unit of S . Since each Sx meets aS with $a \in I$ we have that I is connected.

Example: Let $S = \{(x, y) \mid 0 \leq x \leq 1 \quad 0 \leq y \leq 1\}$.

For (x_1, y_1) and $(x_2, y_2) \in S$ define the product $(x_1, y_1) \cdot (x_2, y_2)$ to be $(0, y_1 y_2)$.

Then S is a compact connected commutative mob.

Let $I = \{(x, y) \mid x = 0, 1 \quad 0 \leq y \leq 1\}$.

And $I^* = \{(x, y) \mid 0 \leq x < \frac{1}{4} \quad \frac{3}{4} < x \leq 1 \quad 0 \leq y < 1\}$.

Then I is a disconnected closed ideal, and I^* is a disconnected open ideal.

Theorem 1: If S is connected and I an ideal of S , then one and only one component of I is an ideal of S .

Proof:

Let $I^* = SI \cup IS$. Then I^* is connected and the component of I which contains I^* is an ideal of S .

Furthermore it is readily seen that this is the only component of S which is an ideal. This ideal will be called the component ideal of I .

Lemma 3: Let S be a compact connected mob and U a proper open subset of S with $J_0(U) \neq \emptyset$.

Let C_0 be the component ideal of $J_0(U)$, then C_0 intersects $\bar{U} \setminus U$.

Proof:

If $\bar{C}_0 \cap \bar{U} \setminus U = \emptyset$, then $\bar{C}_0 \subset U$, and since \bar{C}_0 is an ideal, we have $\bar{C}_0 \subset J_0(U)$ and $C_0 = \bar{C}_0$.

Furthermore $J_0(U)$ is open and hence we can find an open set V , with $C_0 \subset V \subset \bar{V} \subset J_0(U)$.

Since C_0 is a component of the compact set \bar{V} of the connected set S , we have $C_0 \cap \bar{V} \setminus V \neq \emptyset$ a contradiction.

Corollary 1:

Let S be a compact connected mob and F a closed subset of $S \setminus K$, with the property that if $F \cap I \neq \emptyset$, then $F \subset I$ for any ideal I of S .

Then if C is the component of $S \setminus F$ which contains K then $F = \bar{C} \setminus C$.

Proof:

Since C is closed in $S \setminus F$ we have $\bar{C} \cap S \setminus F = C \Rightarrow F \supset \bar{C} \setminus C$.

Furthermore it follows from lemma 3 that if C_0 is the component ideal of $J_0(S \setminus F)$, then $K \subset C_0$ and \bar{C}_0 intersects $\overline{S \setminus F} \setminus S \setminus F \subset F$.

Hence $F \subset \bar{C}_0 \subset \bar{C}$.

Since $F \cap C = \emptyset$ we have $F \subset \bar{C} \setminus C$.

If we take in corollary 1 $F = H(e)$ with $e \in E \setminus K$ and if C is the component of $S \setminus H(e)$ which contains K , then $H(e) = \bar{C} \setminus C$.

This follows immediately from corollary 1, since if $H(e) \cap I \neq \emptyset$, then $H(e) \subset I$ for any ideal I of S . Furthermore it follows that $H(e)$ with $e \in E \setminus K$ can contain no innerpoints.

Theorem 2: Let S be a compact connected mob. If K is not the cartesian product of two non-degenerate connected sets, then

either K is a group or the multiplication in K is of type (a) or (b).

$$(a) \quad xy = x \quad \text{all } x, y \in S$$

$$(b) \quad xy = y \quad \text{all } x, y \in S.$$

Proof:

From Ch I. §2 lemma 4 we know that

$$K = \{ Se \cap E \} \cdot eSe \cdot \{ eS \cap E \} \quad e \in E \cap K.$$

Now let $K^* = (Se \cap E) \times (eSe) \times (eS \cap E)$ and $\phi : K^* \rightarrow K$

$$\phi(x, y, z) = xyz.$$

Then ϕ is clearly a continuous mapping of K^* onto K .

Now let $x_1 y_1 z_1 = x_2 y_2 z_2$ with $x_1, x_2 \in Se \cap E$, $z_1, z_2 \in eS \cap E$

$$y_1, y_2 \in eSe.$$

Then since $x_1 S$ and $x_2 S$ are minimal ideals with $x_1 S \cap x_2 S \neq \emptyset$ we have $x_1 x_2 = x_2$.

Furthermore since

$$x_1, x_2 \in Se, \quad Se = Sx_1 = Sx_2 \Rightarrow x_1 e = x_1, \quad x_2 e = x_2, \quad ex_1 = ex_2 = e.$$

$$\text{Hence } x_2 = x_1 x_2 = x_1 (ex_2) = x_1 e = x_1.$$

In the same way we can prove $z_1 = z_2$.

$$\text{Since } x_1 y_1 z_1 = x_2 y_2 z_2 \text{ we have } ex_1 y_1 z_1 e = ex_2 y_2 z_2 e \Rightarrow ey_1 e = ey_2 e \Rightarrow y_1 = y_2.$$

Hence ϕ is one to one and K is homeomorphic to K^* .

Since K is connected, each of eSe , $Se \cap E$ and $eS \cap E$ must be connected.

Hence at least two of the factors must consist of single elements.

If $eS \cap E = Se \cap E = e$, then $K = eSe$ and hence a group.

If $eS \cap E = eSe = e$, then $K = Se$, and if $x, y \in K$ we have $xy = (xe)(ye) = x(eye) = xe = x$.

If $Se \cap E$ and eSe are both e , then the multiplication is of type (b).

Corollary: Let S be a compact connected mob. If K contains a cutpoint, then the multiplication in K is of type (a) or (b).

Proof:

If K contains a cutpoint, then K is not the cartesian product of two non-degenerate connected sets.

Hence from theorem 2 it follows that K is a group or the multiplication is of type (a) or (b).

Since a compact connected group, contains no cutpoints, the corollary follows.

Definition 1: A clan is a compact connected mob with a unit element.

Lemma 4: Let B be the solid unit ball in Euclidean n -space and let f be a map of B into itself, such that $|x - f(x)| < \frac{1}{2}$ for all $x \in B$. Then $0 \in f(B)$.

Proof:

Let $x = (x_1, \dots, x_n)$ $f(x) = (f_1(x), \dots, f_n(x))$.

We now consider the mapping $h(x) = (x_1, \dots, x_n) - (f_1(x), \dots, f_n(x))$.

This mapping transforms the ball $|x| \leq \frac{1}{2}$ into itself and hence by Brouwer's fixed point theorem there is a point x^* for which $h(x^*) = 0$.

i.e. $(x_1^*, \dots, x_n^*) = (x_1^*, \dots, x_n^*) - (f_1(x^*), \dots, f_n(x^*))$ $f(x^*) = 0$.

Theorem 3: Let S be a mob with unit element u having an Euclidean neighbourhood U of u .

Then $H(u)$ is an open subset of S and is a Lie group.

Proof:

We identify U with E^n and let $F_\epsilon = \{x \mid |u-x| \leq \epsilon\}$.

Since the multiplication on F is uniformly continuous there is a δ such that $|x-xy| < \epsilon/2$, $|x-yx| < \epsilon/2$ whenever $|u-y| < \delta$.

By lemma 4 $u \in F_\epsilon y$ and $u \in y F_\epsilon$, hence y has an inverse y^{-1} in F_ϵ and the mapping $y \rightarrow y^{-1}$ is continuous.

Therefore $H(u)$ is a topological group, and since it contains an open set it must be open in S .

Furthermore $H(u)$ is locally Euclidean and hence a Lie group.

Corollary 1: If S is a clan having a Euclidean neighbourhood of the identity then S is a Lie group.

Proof:

By theorem 3 $H(u)$ is open. $H(u)$ is closed since S is compact, and hence $H(u)$, must be all of S . Thus if S is a clan and S is an n -sphere, then S is a topological group, and hence $n = 0, 1$ or 3 .

In general a compact manifold which admits a continuous associative multiplication with identity must be a group.

Corollary 2: Let S be a clan and F a closed subset of S such that $S-F$ is locally Euclidean.

Then either S is a group or $H(u) \subset F$.

Proof:

Let $h \in H(u)$ and $h \notin F$. Then h has a Euclidean neighbourhood V . Since $h^{-1}V$ is a Euclidean neighbourhood of u , it follows from corollary 1 that S is a group.

In case S is a subset of Euclidean space, then it follows from corollary 2 that $H(u) \subset \text{boundary of } S$ or S a top. group.

If S contains interior points, then it cannot be a group and we have $H(u) \subset \text{Bd}(S)$.

Definition 2: A subset C of a space X is a C -set provided that $C \neq X$ and if M is a continuum with $C \cap M \neq \emptyset$ then $M \subset C$ or $C \subset M$. It can easily be shown that if C is a C -set of a compact connected Hausd. space, then the interior of C is empty and C is connected.

For let x be an interior point of C , then there is an open set V with $x \in V \subset \bar{V} \subset C$.

Now let $y \in X-C$. Then the component M of y in $X-V$ has a non-empty intersection with the boundary of $X-V \subset \bar{V}$.

Hence M is a continuum with $M \cap C \neq \emptyset$ and $C \not\subset M$, $M \not\subset C$.

Theorem 4: (Gleason).

Let G be a compact Lie group which acts on a completely regular space X . Let $p \in X$ such that $g(p) \neq p$ unless g is the identity; $g \in G$.

Then there exists a closed neighbourhood N of p and a closed subset C of N , such that the orbit of every point of N has exactly one point in common with C .

Proof: See Gleason Pr A.M.S. 1 1950.

Lemma 5: Let G be a compact group and let U be an open neighbourhood of the identity.

Then U contains an invariant subgroup H of G such that G/H is a Lie group.

Proof: See Montgomery Zippin: Topological transformation groups.

Theorem 5: Let S be a clan, S no group G a compact invariant subgroup of $H(u) = H$, such that H/G is a Lie group.

Then S contains a continuum M such that M meets H and the complement of H , and such that $u \in M \cap H \subseteq G$.

Proof:

We can consider H as transformation group acting on S . Let $H' = H/G$ and S' the space of orbits of G . Then H' is a compact Lie group acting on S' .

By theorem 4 there exists a closed neighbourhood N of $u' = u.G$ and a closed set $C \subset N$ such that $n H' \cap C$ is a single point for each $n \in N$.

Now let S'' be the space of orbits under H .

Then we have the following canonical mappings $\alpha: S \rightarrow S'$,

$\beta: S' \rightarrow S''$ $\gamma: S \rightarrow S''$, with $\gamma = \alpha \cdot \beta$.

Since α and γ are open maps, β is also open.

Let N^0 be the interior of N then βN^0 is open and $\beta(u') \in \beta(N^0)$.

Let P be the component of $\beta(N)$ which contains $\beta(u')$.

Then P meets the boundary of $\beta(N)$ and hence P is non-degenerate.

Now let $\beta^* = \beta|C$.

Then since $n H' \cap C$ is a single point for each $n \in N$ it follows that β^* is a homeomorphism between C and $N\beta$.

$\beta^{*-1}(P)$ is a continuum which meets H' only at $C \cap H'$, and hence $\beta^{*-1}(P)$ also meets the complement of H' .

Now let K be a component of $\alpha^{-1} \beta^{*-1}(P)$.

Since α is an open mapping we have $\alpha(K) = \beta^{*-1}(P)$.

Hence K is a continuum which meets H and the complement of H and $K \cap H \subset \alpha^{-1}(c)$, where $c = C \cap M$.

Let $h \in K \cap H$, then $K \cap H \subset hG$.

Suppose now $M = h^{-1}K$, then $u \in M \cap H$ and $M \cap H \subset G$ and if $k \in K$, $k \notin H$, then $h^{-1}k \in M$; $h^{-1}k \notin H$, since $S-H$ is an ideal of S . q.e.d.

Theorem 6: Let S be a clan which is no group.

Then the identity u of S belongs to no non-trivial C -set.

Proof:

Let $u \in C$, with C a C -set. We first prove that $C \subset H(u)$.

If $x \in C$, then since xS is a continuum which meets C , we have $C \subset xS$ or $xS \subset C$.

If $u \in xS$, then x has an inverse and is thus included in $H(u)$.

Now let $u \notin xS$, then $xS \subset C$; $xS \neq C$, and there is an open set V with $xS \subset V$; $C \setminus V \neq \emptyset$. Since $xK \subset K$ we have $K \cap C \neq \emptyset$.

If $u \in K$ then S is a group, hence $u \notin K \Rightarrow K \subset C$.

We can find now an open set W with $x \in W$ $WS \subset V$.

Since C contains no interior points there exists a $y \in W \setminus C$ with $yS \subset V$.

Clearly yS is a continuum which meets both C and $S \setminus C$ and $C \not\subset yS$ a contradiction.

Hence $u \in xS$ and thus $x \in H(u) \Rightarrow C \subset H(u)$.

Now let U be a neighbourhood of u such that $C \not\subset U$.

Then by lemma 5 there is a subgroup $G \subset U$ such that H/G is a Lie group and $C \not\subset G$.

Theorem 5 implies that we can find a continuum M such that $u \in M \cap H \subset G$ and such that M meets the complement of H .

Hence $M \cap C \neq \emptyset$ and since $C \subset H$ M meets the complement of C .
 $\Rightarrow C \subset M$.

Since $M \cap H \subset G$ and $C \not\subset G \Rightarrow C \not\subset M$ a contradiction. q.e.d.

Example: Let $A = \{(x, y) \mid y = \sin \frac{1}{x} \ 0 < x \leq 1\}$
 $B = \{(2-x, y) \mid (x, y) \in A\}$.
 $C = \{(0, y) \cup (2, y) \mid -1 \leq y \leq 1\}$.

and let $S = A \cup B \cup C$.

We will show that S does not admit the structure of a clan.

For suppose that S is a clan.

Since S is not homogeneous, S cannot be a topological group and hence $S \neq H(u)$.

Then $S \setminus H(u) = J \neq \emptyset$ is the maximal proper ideal of S . Since J is open, dense and connected we have $A \cup B \subset J$ and hence $u \in C$. But since C is the union of two C -sets, u cannot be in C .

Lemma 6: Let S be a clan and C a C -set of S . If g is an idempotent with $g \notin K$, then $g \notin C$.

Proof:

Suppose $g \in C$. Since gSg is a continuum we have $C \subset gSg$ or $gSg \subset C$.

g is the identity of the clan gSg and gSg is not a group since $g \notin K$ (Ch I. § 3 th.6). Hence theorem 6 implies that $C \not\subset gSg$.

Now suppose $gSg \subset C \Rightarrow K \cap C \neq \emptyset$ and since $g \in C$ $C \setminus K \neq \emptyset$.

Let U and V be neighbourhoods of K with $SK=K \subset U \subset \bar{U} \subset V$.

while $g \notin V$.

Since S is compact there is a neighbourhood W of K such that $SW \subset U$

\overline{SW} is a continuum and hence $\overline{SW} \subset C$.

Furthermore $W \subset \overline{SW}$ and this would imply that C contains inner points; a contradiction.

Theorem 7: Let S be a clan and C a C -set of S , then $C \subseteq K$.

Proof:

From the proof of lemma 6 it follows that if $K \cap C \neq \emptyset$, then $C \subseteq K$.

Suppose now $C \cap K = \emptyset$ and let $x \in C$ and U a neighbourhood of x with $C \setminus U \neq \emptyset$.

Let e be a minimal member of the partial ordered set E with $xe = x$.

e exists since $E_x = \{e \mid e^2 = e \text{ } xe = x\} \neq \emptyset$ and compact.

Furthermore $e \notin K$ since $x \notin K$.

Hence $H(e) \neq eSe$ and we can find a neighbourhood V of e such that $xV \subset U$ and a continuum $M \subset eSe$ such that $e \in M \subset V$ and $M \cap \{eSe \setminus H(e)\} \neq \emptyset$.

Since $x \in xM$ we have $xM \subset C$.

Let $m \in M \cap \{eSe \setminus H(e)\}$, then $C \subset xSm$.

This implies that $x = xsm = xesem = xp$ with $p \in \{eSe \setminus H(e)\}$. since $\{eSe \setminus H(e)\}$ is an ideal of eSe .

Hence $x = xf$ with $f = f^2 \in \Gamma(p) \subset eSe$, and thus $ef = fe = f \Rightarrow f \leq e$.

But since e is minimal we have $f = e$.

Furthermore $pe = p = ep \Rightarrow pf = p = fp \Rightarrow p \in H(f) = H(e)$; a contradiction.

Theorem 8: If S is a clan and if K is a C -set, then K is a maximal subgroup of S .

Proof:

If $S = K$, then S is a group and the result follows.

If $S \neq K$, then K has no interior points since K is a C -set.

Let $\{a_\lambda \mid \lambda \in \Lambda\}$ be a directed set of points of $S \setminus K$ with $a_\lambda \rightarrow e$ where $e = e^2 \in K$.

Since $K \cap a_\lambda S \neq \emptyset$ $K \cap Sa_\lambda \neq \emptyset$ and $a_\lambda \in a_\lambda S \cap Sa_\lambda$ we have

$K \subset a_\lambda S \cap Sa_\lambda \Rightarrow K \subset eS \cap Se = eSe$.

Now $e \in K$ gives $H(e) = eSe$ and thus $K = H(e)$.

Theorem 9: If a clan is an indecomposable continuum it is a group.

Proof:

If $S = K$, then S is a group.

Suppose now $K \neq S$. Then there exists an open set V with $K \subset V \subset \bar{V} \neq S$. Let $J_0(V)$ be the union of all ideals of S contained in V , then $J_0(V)$ is open and connected and $K \subset J_0(V) \subset J_0(\bar{V}) \neq S$.

Since $S = \overline{J_0(V)} \cup S - \overline{J_0(V)}$ and S is indecomposable we have $S - \overline{J_0(V)}$ not connected.

Let $S - \overline{J_0(V)} = A \cup B$ $A \cap B = \emptyset$ A, B open.

Then we have $\overline{J_0(V)} \cup A$ connected and $\overline{J_0(V)} \cup B$ connected and hence S not indecomposable; a contradiction.

§ 4. I-semigroups

Definition 1:

Let $J = [a, b]$ denote a closed interval on the real line.
If J is a mob such that a acts as a zero-element and b as an identity then J will be called an I-semigroup.

We will identify J usually with $[0, 1]$, so that $0x = x0 = 0$ and $1x = x1 = x$ for all $x \in I$.

Example: $J_1 = [0, 1]$ under the usual multiplication

$J_2 = [\frac{1}{2}, 1]$ with multiplication defined by $xoy = \max(\frac{1}{2}, xy)$ where xy denotes the usual multiplication of real numbers.

$J_3 = [0, 1]$ with multiplication defined by $xoy = \min(x, y)$.

J_1 and J_2 have just the two idempotents zero and identity, but in J_3 every element is an idempotent.

Furthermore every non-idempotent element in J_2 is algebraically nil-potent i.e. for every $x \in J_2$ there exists an n such that x^n is equal to zero.

Lemma 1: If J is an I-semigroup, then $xJ = Jx = [0, x]$ for all $x \in J$.

Proof:

Since xJ is connected and $0, x \in xJ$ we have $[0, x] \subset xJ$ and by the same argument $Jx \supset [0, x]$.

$J_0([0, x)) = J_0$ is open and connected and hence $x \in \bar{J}_0$ and \bar{J}_0 an ideal of J .

Hence $Jx \subset J\bar{J}_0 \subset \bar{J}_0 \subset [0, x]$ and $xJ \subset [0, x]$.

Thus $xJ = Jx = [0, x]$.

Corollary: If J is an I-semigroup, then $x \leq y$ and $w \leq v \Rightarrow xw \leq yv$.

Proof: Since $x \leq y$ there is a z such that $x = zy$.

Hence $xw = z(yw) \leq yw$.

In the same way we can prove $yw \leq yv \Rightarrow xw \leq yv$.

Theorem 1: If J is an I-semigroup with just the two idempotents 0 and 1 and with no nilpotent elements, then J is isomorphic to J_1 .

Proof: We first show that if $xy = xz \neq 0$ then $y = z$.

Assume $y < z$. Then by lemma 1 there is a w such that $y = zw$.

Hence $xy = x(zw) = xyw \Rightarrow xy = (xy)w^n$ for every $n > 0$.

Thus $xy = (xy)e$, with $e = e^2 \in \Gamma(w)$.

Since $1 \notin \Gamma(w)$, we have $e = 0 \Rightarrow xy = 0$ a contradiction.

We now prove that if $x \neq 0$, then x has a unique square root.

The function $f: J \rightarrow J$ defined by $f(x) = x^2$ is continuous and leaves 0 and 1 fixed. Hence f is a map of J onto J so that square roots exist for every element.

Assume $a^2 = b^2 \neq 0$ and let $a \leq b$.

Then by lemma 1 $a^2 \leq ab \leq b^2$. Hence $ab = a^2 \Rightarrow b = a$.

This establishes that for $x \neq 0$, x has a unique square root and by induction that x has unique 2^n -th roots.

Let x_n be the 2^n -th root of $x \neq 0$ and for $r = p/2^n$ define $x^r = x_n^p$.

Then it is easy to prove that $x^r \cdot x^s = x^{r+s}$, where r, s are positive dyadic rationals.

Furthermore if $r < s$, then $x^r > x^s$. For by lemma 1 $x^r \geq x^s$, and if $x^r = x^s$, then $x^{r-s} = 1$, a contradiction.

This implies that $\lim x_n = 1$.

Since $x_n < x_{n+1}$ $\lim x_n$ exist. Assume $\lim x_n = y \neq 1$.

Then since $y \rightarrow 0$ there is an n_0 such that $y^{n_0} < x$.

Hence $y < x_{n_0}$ a contradiction.

Now let $D = \{x^r \mid r \text{ a positive dyadic rational}\}$.

Then D is a commutative submob of J and $\bar{D} = J$.

Assume $\bar{D} \neq J$. Then there is an open interval $P \subset J \setminus \bar{D}$.

$P = (a, b)$ and $b \in \bar{D}$.

Now since $x_n \rightarrow 1$, $x_n b \rightarrow b$, and $x_n b \leq b$ by lemma 1.

If $x_n b = b$, then $x_n = 1$ a contradiction.

Hence $x_n b < b$ and $x_n b \in P$ for n sufficiently large.

Since $b \in \bar{D}$ and $x_n \in \bar{D}$, we have $x_n b \in \bar{D}$ a contradiction

And thus $\bar{D} = J$.

Now let $g: D \rightarrow J_1$ be defined by $g(x^r) = \frac{1}{2}^r$.

$g(D)$ is dense in J_1 and g is one- to one continuous and order preserving.

Hence g can be extended to an isomorphism of J onto J_1 .

Theorem 2: If J is an I-semigroup with just the two idempotents 0 and 1 and with at least one nilpotent, then J is isomorphic to J_1 .

Proof:

Let $d = \sup \{ x \mid x^2 = 0 \}$. Then $d \neq 0$.

For let $y \neq 0$ be nilpotent, then $y^n = 0$, $y^{n-1} \neq 0$ for some $n > 1$.

Clearly $(y^{n-1})^2 = 0$. Hence $d \geq y^{n-1}$.

As shown in theorem 1, d has a unique 2^n th root and if r and s are positive dyadic rationals, then $d^r < d^s$ if $r > s$ and $d^s \neq 0$, and $d^r d^s = d^{r+s}$.

Now let $D = \{ d^r \mid r \text{ a positive dyadic rational} \}$. Then by the same type of argument used in the proof of theorem 1, $\bar{D} = J$.

We define $g: D \rightarrow J_2$ by $g(d^r) = (\sqrt{\frac{1}{2}})^r$. Then g is one to one and continuous and $g(D)$ is dense in J_2 .

Since g is order preserving it can be extended to an isomorphism of J onto J_2 .

Theorem 3: Let J be an I-semigroup. Then E is closed and if $e, f \in E$, then $e.f = \min(e, f)$.

The complement of E is the union of disjoint intervals.

Let P be the closure of one of these. Then P is isomorphic to either J_1 or J_2 . Furthermore if $x \in P$, $y \notin P$ then $xy = \min(x, y)$.

Proof:

Let $e, f \in E$ $e < f$. Then by lemma 1 $e.e \leq ef \Rightarrow e \leq ef$.

Since $ef \leq e$, we have $e = ef$.

Now let $Q = [e, f]$.

Then for any $x, y \in [e, f]$ we have $e.e \leq x.y \leq f.f$.

Hence Q is a submob of J .

Furthermore if $e \leq x$, then $e \geq ex \geq e.e = e \Rightarrow ex = e$.

In other words e acts as a zero for $[e, 1]$.

If $x \leq f$, then by lemma 1 $x = fy$ and thus $fx = x$.

f acts as an identity for $[0, f]$.

So we have in particular P an I-semigroup with only two idempotents and hence P is isomorphic either to J_1 or J_2 .

If $x \in P$, $y \notin P$, $x \leq y$ then there is an $e \in E$, with $x \leq e \leq y$.

Hence $xy = (xe)y = x(ey) = xe = x$.

It follows from theorem 3, that every I-semigroup is commutative.

Theorem 4: Let S be the closed interval $[a, b]$. If S is a mob such that a and b are idempotents and S contains no other idempotents, then S is abelian.

Proof:

Let $e \in E \cap K$. Then $H(e) = eSe$.

Since S has the fixed point property and $H(e)$ is a retract of S , $H(e)$ has the fixed point property and hence $H(e) = e$.

Consequently every element of K is idempotent.

Since K is connected, $K = a$ or $K = b$.

If $K = a$, then a is a zero for S and g an identity since $gS = Sg = S$.

Hence S is an I-semigroup and abelian.

Theorem 5: Let S be the closed interval $[a, b]$. If S is a clan such that both a and b are idempotents, then S is abelian if and only if S has a zero.

Proof:

Let S be commutative, then K is a group and by the same argument used in the proof of th. 4, the maximal subgroups in K are single elements, hence K consists of only one element, a zero.

Now let S have a zero. If either a or b is the zero element, then the other is obviously a unit and the result follows by theorem 3.

Now let $a < 0 < b$. Then $S' = [a, 0]$ is a submob of S .

For suppose there exists $x, y \in S'$ with $xy \in (0, b]$.

Then since a acts as a unit on S' , we have $x, xy \in x[a, y]$.

Hence there is an $s^* \in [a, y]$ with $xs^* = 0$.

Since $0, s^* \in S'$, we have $y = s^*q$.

Hence $xy = xs^*q = 0q = 0$ a contradiction.

In the same way we can prove that $S'' = [0, b]$ is a submob of S and both S' and S'' are commutative since they are I-semigroups. It also follows that the unit of S is either a or b . Suppose b is the unit element. Then in the same way as above we can prove that $aS'' = S''a = [0, a]$.

Hence if $x'' \in S''$ then $ax'' = y''a = (y''a)a = a(x''a) = a(az'') = az'' = x''a$.

Furthermore if $x' \in S'$ and $x'' \in S''$, then $x'x'' = (x'a)x'' = x'(ax'') = (ax'')x' = (x''a)x' = x''x'$.

Theorem 6: Let S be the closed interval $[a, b]$. If S is a mob such that a and b are idempotents, then S is abelian if and only if S has a zero and $ab = ba$.

Proof:

If S is commutative, S has a zero by the same argument as in theorem 5, and obviously $ab = ba$.

Now let S have a zero and let $ab = ba$.

Then again the result follows if either a or b is a zero.

If $a < 0 < b$, then $S' = [a, 0]$ and $S'' = [0, b]$ are abelian submobs of S .

Suppose now $ab \in S'$, then $bS' = baS' = abS' = [ab, 0]$ by lemma 1.

Hence $bS = Sb = [ab, b]$, and $[ab, b]$ is an abelian submob by theorem 5.

To prove the theorem it suffices to show that if $x \in [0, ab]$ and $y \in [ab, b]$ then $xy = yx$.

Now $xy = (xa)(by) = (xab)y$, and $xab \in [ab, 0]$.

Hence $(xab)y = y(xab) = y(xb) = (yb)xb = y(bxb) = ybbx = yx$.