

STICHTING
MATHEMATISCH CENTRUM
2e BOERHAAVESTRAAT 49
AMSTERDAM

SP 90

(S 345)

. J. Fabius

A probabilistic Example of a nowhere analytic $C -$
function

(Zeitschrift fuer wahescheinlichkeitstheorie
und verwandte Gebiete, 5 (1966), p 173-174)



A Probabilistic Example of a Nowhere Analytic C^∞ -Function*

J. FABIUS

Received May 24, 1965

It is well known that there exist functions which are infinitely differentiable but nowhere analytic. In fact, MORGENSTERN [1] has shown that the nowhere analytic functions are by far in the majority among the infinitely differentiable functions on $[0, 1]$. A number of examples, as well as an extensive bibliography, are given by SALZMANN and ZELLER [2].

It is the purpose of this note to show that another example is provided by the restriction on $[0, 1]$ of the distribution function F of $X = \sum_{n=1}^{\infty} 2^{-n} \xi_n$, where ξ_1, ξ_2, \dots are independent random variables, uniformly distributed on $[0, 1]$.

One easily verifies that $F(0) = 0$, $F(1) = 1$ and $0 < F(x) < 1$ for all $x \in (0, 1)$. Moreover, since $2 \sum_{n=2}^{\infty} 2^{-n} \xi_n$ is independent of ξ_1 , and has the same distribution as X ,

$$(1) \quad F(x) = \int_0^1 F(2x - y) dy = \begin{cases} \int_0^{2x} F(t) dt & \text{for } x \in [0, \frac{1}{2}], \\ \int_{2x-1}^1 F(t) dt + 2x - 1 & \text{for } x \in (\frac{1}{2}, 1], \end{cases}$$

which implies that F is a continuous function. For notational convenience we introduce another function f , which we take to be the unique function on $[0, \infty)$ which coincides with F on $[0, 1]$ and satisfies the relations

$$(2) \quad f(x) = \begin{cases} 1 - f(x - 1) & \text{for } x \in (1, 2] \\ -f(x - 2^n) & \text{for } x \in (2^n, 2^{n+1}], \quad n = 1, 2, \dots \end{cases}$$

Clearly f is continuous and vanishes only at the nonnegative even integers. Moreover,

$$(3) \quad f(x) = \int_0^{2x} f(t) dt$$

for all $x \in [0, \infty)$. For $x \in [0, \frac{1}{2}]$ this is just a restatement of (1), for $x \in (\frac{1}{2}, 1]$ and for $x \in (1, 2]$ it follows from (1) and (2) by easy computations, and for $x \in (2^n, 2^{n+1}]$, $n = 1, 2, \dots$ we prove (3) by induction on n : If (3) holds for all $x \in [0, 2^n]$ and some $n \geq 1$, then we have for $x \in (2^n, 2^{n+1}]$

$$f(x) = -f(x - 2^n) = -\int_0^{2x-2^{n+1}} f(t) dt = \int_{2^{n+1}}^{2x} f(t) dt = \int_0^{2x} f(t) dt,$$

* Report S 345, Stat. Dept., Mathematisch Centrum, Amsterdam.

since (2) implies that

$$\int_0^{2^{n+1}} f(t) dt = 0$$

for all $n \geq 1$.

It follows from (3) that f , being continuous, is differentiable with

$$f'(x) = 2f(2x)$$

for all $x \in [0, \infty)$ and hence, that f is infinitely differentiable with

$$(4) \quad f^{(n)}(x) = 2^{n(n+1)/2} f(2^n x)$$

for all $x \in [0, \infty)$, $n = 1, 2, \dots$

In view of the fact that f vanishes only at the nonnegative even integers, (4) implies that at any binary rational of the form $x = (2k - 1)2^{-n}$ with k and n positive integers all derivatives except the first n vanish. Consequently the Taylor series expansion of f around such a point is a polynomial of degree n , which cannot possibly coincide with f on any neighborhood, since at every binary irrational point all derivatives of f are nonzero. Thus f is nowhere analytic, being singular at all binary rationals.

References

- [1] MORGENSTERN, D.: Unendlich oft differenzierbare nichtanalytische Funktionen. Math. Nachr. **12**, 74 (1954).
- [2] SALZMANN, H., und K. ZELLER: Singularitäten unendlich oft differenzierbarer Funktionen. Math. Z. **62**, 354–367 (1955).

Stat. Dept., Math. Centrum
Amsterdam, Netherlands