

**stichting
mathematisch
centrum**



AFDELING NUMERIEKE WISKUNDE
(DEPARTMENT OF NUMERICAL MATHEMATICS)

NW 73/79

NOVEMBER

M. BAKKER

A NOTE ON C^0 GALERKIN METHODS FOR TWO-POINT
BOUNDARY PROBLEMS

Preprint

2e boerhaavestraat 49 amsterdam

Printed at the Mathematical Centre, 49, 2e Boerhaavestraat, Amsterdam.

The Mathematical Centre, founded the 11-th of February 1946, is a non-profit institution aiming at the promotion of pure mathematics and its applications. It is sponsored by the Netherlands Government through the Netherlands Organization for the Advancement of Pure Research (Z.W.O).

A note on C^0 Galerkin methods for two-point boundary problems^{*)}

by

M. Bakker

ABSTRACT

As is known [4], the C^0 Galerkin solution of a two-point boundary problem using piecewise polynomial functions, has $O(h^{2k})$ convergence at the knots, where k is the degree of the finite element space. In this note, it is proved that on any segment there are $k-1$ interior points where the Galerkin solution is of $O(h^{k+2})$, one point better than the global order of convergence. These points are the Lobatto points.

KEY WORDS & PHRASES: *two-point boundary problems, finite element method, superconvergence, Lobatto points.*

^{*)} This paper will be submitted for publication elsewhere.

1. INTRODUCTION

We consider the two-point boundary problem

$$\begin{aligned} Lu &\equiv -(p(x)u')' + q(x)u = f(x), & x \in [0,1] = I; \\ (1) \quad u(0) &= u(1) = 0. \end{aligned}$$

We suppose that p , q and f are such that (1) has a unique and sufficiently smooth solution.

Let, for a constant integer N , $\Delta: 0 = x_0 < x_1 < \dots < x_N = 1$ be a partition of I with

$$h = N^{-1}; \quad x_j = jh; \quad I_j = [x_{j-1}, x_j]$$

and let for a constant integer $k \geq 2$ and for any interval $E \subset I$, $P_k(E)$ be the class of polynomials of degree at most k restricted to E .

We define for $m \geq 0$ and $s \geq 1$

$$\begin{aligned} W^{m,s}(E) &= \{v \mid D^j v \in L^s(E), j = 0, \dots, m\}; \\ H^m(E) &= W^{m,2}(E); \\ H_0^1(I) &= \{v \mid v \in H^1(I); v(0) = v(1) = 0\}; \\ (2) \quad M_0^k(\Delta) &= \{v \mid v \in H_0^1(I); v \in P_k(I_j), j = 1, \dots, N\}; \\ \|v\|_{W^{m,s}(E)} &= \left[\sum_{j=0}^m \|D^j v\|_{L^s(E)}^2 \right]^{1/2}; \\ \|v\|_{H^m(E)} &= \left[\sum_{j=0}^m (D^j v, D^j v)_{L^2(E)} \right]^{1/2}, \end{aligned}$$

where D^j denotes d^j/dx^j . If $E = I$, we write (α, β) instead of $(\alpha, \beta)_{L^2(I)}$ and $\|\alpha\|_m$ instead of $\|\alpha\|_{H^m(I)}$.

Let $U \in M_0^k(\Delta)$ be the unique solution of

$$(3) \quad B(U, V) = (f, V), \quad V \in M_0^k(\Delta),$$

where $B: H_0^1(I) \times H_0^1(I) \rightarrow \mathbb{R}$ is defined by

$$(4) \quad B(u, v) = (pu', v') + (qu, v); \quad u, v \in H_0^1(I).$$

We assume that B is strongly coercive, i.e. there exists a $C > 0$ such that

$$(5) \quad B(v, v) \geq C \|v\|_1^2, \quad v \in H_0^1(I).$$

In the sequel, we assume that C, C_1, C_2 , etc., are generic positive constants not necessarily the same.

LEMMA 1. *Let $u \in H_0^1(I) \cap H^{k+1}(I)$ be the solution of (1) and let $U \in M_0^k(\Delta)$ be the solution of (3). Then the error function $e(x) = u(x) - U(x)$ has the bounds*

$$\|e\|_\ell \leq Ch^{k+1-\ell} \|u\|_{k+1}, \quad \ell = 0, 1;$$

$$(6) \quad |e(x_j)| \leq Ch^{2k} \|u\|_{k+1}, \quad j = 1, \dots, N-1;$$

$$\|e\|_{L^\infty(I)} \leq Ch^{k+1} \|u\|_{k+1}.$$

PROOF. See [5], [4] and [6]. \square

In the next §, we prove that the local order of convergence improves slightly at specific points interior on I_j , if u satisfies stricter smoothness requirements on the interior of I_j .

2. ORDER OF CONVERGENCE AT LOBATTO POINTS

On the segment $[-1, +1]$, we define the Lobatto points $\sigma_0, \dots, \sigma_k$ by

$$(7) \quad (1 - \sigma_\ell^2) \frac{d}{d\sigma} P_k(\sigma_\ell) = 0, \quad \ell = 0, \dots, k,$$

where $P_k(\sigma)$ is the k -th degree Legendre polynomial. Associated to this polynomial is the quadrature formula (see [1, formula 2.5.4.32])

$$\int_{-1}^{+1} f(\sigma) d\sigma = \sum_{\ell=0}^k w_{\ell} f(\sigma_{\ell}) + R_k f^{(2k)}(s \in (-1, +1));$$

(8)

$$w_{\ell} = \frac{2}{k(k+1) [P_k(\sigma_{\ell})]^2}, \quad \ell = 0, \dots, k.$$

From (7) and (8), we define

$$\begin{aligned} \xi_{j\ell} &= x_{j-1} + \frac{h}{2}(1+\sigma_{\ell}); \quad \ell = 0, \dots, k; \quad j = 1, \dots, N; \\ (\alpha, \beta)_j^* &= \frac{h}{2} \sum_{\ell=0}^k w_{\ell} \alpha(\xi_{j\ell}) \beta(\xi_{j\ell}); \quad \alpha, \beta \in W^{2k, \infty}(I_j); \quad j = 1, \dots, N; \\ (\alpha, \beta)_h &= \sum_{j=1}^N (\alpha, \beta)_j^*. \end{aligned}$$

(9)

We return to problems (1) and (3). It is known that

$$(10) \quad B(e, V) = 0, \quad V \in M_0^K(\Delta).$$

For any I_j , we define

$$(11) \quad M_0^k(I_j) = \{V \mid V \in M_0^K(\Delta), \text{ supp}(V) = I_j\}.$$

We temporarily drop the subscript j from the numbers $\xi_{\ell j}$. We define a natural basis $\{\phi_i\}_{i=1}^{k-1}$ for $M_0^k(I_j)$ by

$$(12) \quad \phi_i(\xi_{\ell}) = \delta_{i\ell}, \quad 1 \leq i, \quad \ell \leq k-1,$$

where $\delta_{i\ell}$ is the Kronecker symbol. If we elaborate (10) for $V = \phi_i$, $i = 1, \dots, k-1$, we get

$$(13) \quad (e, L\phi_i) = [p(x)e(x)\phi_i^1(x)]_{\xi_0}^{\xi_k}, \quad i = 1, \dots, k-1.$$

Approximation of $(e, L\phi_i)$ by Lobatto quadrature yields

$$\begin{aligned}
\sum_{\ell=1}^{k-1} w_{\ell} L\phi_i(\xi_{\ell}) e(\xi_{\ell}) &= 2h^{-1} [p(x) e(x) \phi_i^1(x)]_{\xi_0}^{\xi_k} - w_0 e(\xi_0) L\phi_i(\xi_0) - \\
(14) \quad &- w_k e(\xi_k) L\phi_i(\xi_k) + Ch^{2k} D^{2k}(eL\phi_i)(\xi \in I_j), \\
i &= 1, \dots, k-1.
\end{aligned}$$

This is a linear system for $e(\xi_1), \dots, e(\xi_{k-1})$. We have to prove the non-singularity of $(w_{\ell} L\phi_i(\xi_{\ell}))$ and to compute the order of the solution.

Take a $V \in M_0^k(I_j)$ represented by

$$V(x) = \sum_{i=1}^{k-1} v_i \phi_i(x).$$

Suppose

$$\sum_{h=1}^{k-1} w_{\ell} L\phi_i(\xi_{\ell}) v_{\ell} = 0, \quad i = 1, \dots, k-1.$$

Then

$$\begin{aligned}
0 &= \frac{h}{2} \sum_{\ell=1}^{k-1} w_{\ell} v_{\ell} \sum_{i=1}^{k-1} v_i L\phi_i(\xi_{\ell}) = \frac{h}{2} \sum_{\ell=1}^{k-1} w_{\ell}^V(\xi_{\ell}) LV(\xi_{\ell}) \\
&= (V, LV) + Ch^{2k+1} D^{2k}(VLV)(\xi \in I_j),
\end{aligned}$$

Hence

$$\begin{aligned}
\|V\|_{H^1(I_j)}^2 &\leq CB(V, V) \leq Ch^{2k+1} \|D^{2k}(VLV)\|_{L^{\infty}(I_j)} \leq Ch^{2k+1} \|V\|_{W^{k, \infty}(I_j)}^2 \\
&\leq Ch^{2k+1} \|V\|_{H^k(I_j)}^2 \leq Ch^3 \|V\|_{H^1(I_j)}^2,
\end{aligned}$$

which is only true if $V \equiv 0$, which proves the nonsingularity of $(w_{\ell} L\phi_i(\xi_{\ell}))$, if h is small enough. Since $h^2 w_{\ell} L\phi_i(\xi_{\ell}) \sim -h^2 w_{\ell} p(\xi_{\ell}) \phi_i''(\xi_{\ell}) = c_{i\ell}$, where $c_{i\ell}$ is of $O(1)$, as $h \rightarrow 0$, all the entries of $(w_{\ell} L\phi_i(\xi_{\ell}))$ are of $O(h^{-2})$, hence the entries of its inverse are of $O(h^2)$.

We turn to the second part of our problem. The first three terms of the

right hand side of (14) are of $O(h^{2k-2}\|u\|_{k+1})$. For the last term, we prove that

$$(16) \quad \|D^{2k}(eL\phi_i)\|_{L^\infty(I_j)} \leq C\|e\|_{W^{2k,\infty}(I_j)} \|L\phi_i\|_{W^{2k,\infty}(I_j)}.$$

From [3], it can be proved that

$$(17) \quad \|D^\ell e\|_{L^\infty(I_j)} \leq \begin{cases} Ch^{k+1-\ell}\|u\|_{k+1}, & \ell \leq k; \\ \|D^\ell u\|_{L^\infty(I_j)}, & \ell > k. \end{cases}$$

Furthermore,

$$(18) \quad \|L\phi_i\|_{W^{2k,\infty}} \leq Ch^{-k},$$

hence we summarily have

$$(19) \quad \left| \sum_{\ell=1}^{k-1} w_\ell L\phi_i(\xi_\ell) e(\xi_\ell) \right| \leq Ch^k [\|u\|_{k+1} h^{k-2} + \|u\|_{W^{2k,\infty}(I_j)}],$$

$$i = 1, \dots, k-1.$$

This was the last step in the proof of

THEOREM 1. Let $u \in H_0^1(I) \cap H^{k+1}(I) \cap \bigcap_{j=1}^N W^{2k,\infty}(I_j)$ be the solution of (1) and let $U \in M_0^k(\Delta)$ be the solution of (3). Then the error function has the local error bounds.

$$(20) \quad |e(\xi_{j\ell})| \leq Ch^{k+2} [\|u\|_{k+1} h^{k-2} + \|u\|_{W^{2k,\infty}(I_j)}],$$

$$j = 1, \dots, N; \quad \ell = 1, \dots, k-1. \quad \square$$

3. LOBATTO QUADRATURE

Usually, $B(\cdot)$ and (\cdot) are to be evaluated by numerical quadrature. We will show that Lobatto quadrature leaves the order of convergence at the

Lobatto points invariant.

We define

$$(21) \quad B_h(\alpha, \beta) = (p\alpha', \beta')_h + (q, \alpha, \beta)_h; \quad \alpha, \beta \in \bigcap_{j=1}^N W^{2k, \infty}(I_j),$$

where $(\cdot)_h$ is defined by (9).

LEMMA 2. Let $Y \in M_0^k(\Delta)$ be the solution of

$$(22) \quad B_h(Y, V) = (f, V)_h, \quad V \in M_0^k(\Delta)$$

and let $u \in H_0^1(I) \cap H^{k+1}(I) \cap \bigcap_{j=1}^N W^{2k, \infty}(I_j)$ be the solution of (1). Then the error function $\eta = u - Y$ has the bounds

$$|\eta(x_j)| \leq Ch^{2k} \|f\|_{2k, \Delta}; \quad j = 1, \dots, N-1,$$

if h is small enough, with

$$(23) \quad \|f\|_{\ell, \Delta} = \left[\sum_{j=1}^N \|f\|_{H^{\ell}(I_j)}^2 \right]^{\frac{1}{2}}.$$

PROOF. See [4]. \square

We now consider $\varepsilon(x) = U(x) - Y(x)$, where U is the solution of (3). From (3) and (22), we obtain for every I_j

$$\begin{aligned} |B(\varepsilon, V)| &\leq |(f, V) - (f, V)_h| + |B_h(Y, V) - B(Y, V)| \\ &\leq Ch^{2k+1} \|V\|_{H^k(I_j)} [\|f\|_{H^{2k}(I_j)} + \|Y\|_{H^k(I_j)}], \quad V \in M_0^k(I_j). \end{aligned}$$

If we take for V any of the basis functions ϕ_i of $M_0^k(I_j)$, as defined by (12), we have

$$(25) \quad |B(\varepsilon, \phi_i)| \leq Ch^{k+1} [\|f\|_{H^{2k}(I_j)} + \|Y\|_{H^k(I_j)}], \quad i = 1, \dots, k-1.$$

Since

$$\begin{aligned}
& \sum_{\ell=1}^{k-1} w_{\ell} \varepsilon(\xi_{\ell}) L\phi_i(\xi_{\ell}) = 2h^{-1} B(\varepsilon, \phi_i) \\
(26) \quad & - w_0 \varepsilon(\xi_0) L\phi_i(\xi_0) - w_k \varepsilon(\xi_k) L\phi_i(\xi_k) \\
& - \frac{2}{h} [p(x) \varepsilon(x) \phi_i'(x)]_{\xi_0}^{\xi_k} + Ch^{2k} D^{2k}(\varepsilon L\phi_i)(\xi \in I_j)
\end{aligned}$$

and

$$\begin{aligned}
(27) \quad & \|D^{2k}(\varepsilon L\phi_i)\|_{L^{\infty}(I_j)} \leq C \|\varepsilon\|_{W^{k,\infty}(I_j)} \|\phi_i\|_{W^{k,\infty}(I_j)} \\
& \leq Ch^{-2k} \|\varepsilon\|_{L^{\infty}(I_j)} \leq Ch^{-k+1} \|f\|_{2k,\Delta},
\end{aligned}$$

we have

$$\begin{aligned}
(28) \quad & \left| \sum_{\ell=1}^{k-1} w_{\ell} \varepsilon(\xi_{\ell}) L\phi_i(\xi_{\ell}) \right| \leq C_1 h^k [\|f\|_{H^{2k}(I_j)} + \|y\|_{H^k(I_j)}] \\
& + C_2 h^{2k-2} \|f\|_{2k,\Delta} + C_3 h^{k+1} \|f\|_{2k,\Delta}.
\end{aligned}$$

The nonsingularity of $(w_{\ell} L\phi_i(\xi_{\ell}))$ has already been proved, its inverse is of $O(h^2)$, hence we have

$$(29) \quad |\varepsilon(\xi_{\ell})| \leq C_1 h^{k+2} [\|f\|_{H^{2k}(I_j)} + \|y\|_{H^k(I_j)}] + C_2 h^{k+3} \|f\|_{2k,\Delta}.$$

Since (see [3]).

$$\begin{aligned}
(30) \quad & \|y\|_{H^k(I_j)} \leq \|\eta\|_{H^k(I_j)} + \|u\|_{H^k(I_j)} \leq Ch \|u\|_{k+1} + \|u\|_{H^k(I_j)} \\
& \leq C \|u\|_{k+1},
\end{aligned}$$

we can prove by combination of (20), (29) and (30)

THEOREM 2. Let $u \in H_0^1(I) \cap H^{k+1}(I) \cap \bigcap_{j=1}^N W^{2k,\infty}(I_j)$ be the solution of (1) and let $Y \in M_0^k(\Delta)$ be the solution of (22). Then the error function η has the bounds

$$|\eta(\xi_{\ell j})| \leq C_1 h^{k+2} [\|f\|_{H^{2k}(I_j)} + \|u\|_{k+1}] + C_2 h^{k+3} \|f\|_{2k,\Delta};$$

$$j = 1, \dots, N; \quad \ell = 1, \dots, k-1. \quad \square$$

4. CONCLUSIONS

We have found a lighter form of superconvergence at other points than the knots. The findings of this paper stress the important part that Lobatto points play in the C^0 Galerkin method for two-point boundary problems. This is especially true for $k = 2$, since in that case the error is of $O(h^4)$ at all Lobatto points.

The results of this paper can be easily applied to the case of two-point initial boundary problems, (see [2]) and probably to other cases, as nonlinear boundary problems.

REFERENCES

- [1] ABRAMOWITZ, A. & I. STEGUN, *Handbook of mathematical functions*, Dover Publications, 1968.
- [2] BAKKER, M., *On the numerical solution of parabolic equations in a single space variable by the continuous time Galerkin method*, SIAM J. Num. Anal. 17, no. 1 (1980).
- [3] CIARLET, P.G. & P.A. RAVIART, *General Lagrange and Hermite interpolation in R^N with applications to finite element methods*, Arch. Rat. Mech. An. 46 (1972), pp. 177-199.
- [4] DOUGLAS, J. Jr. & T. DUPONT, *Galerkin approximations for the two-point boundary problem using continuous, piecewise polynomial spaces*, Num. Mat. 22 (1974), pp. 99-109.
- [5] STRANG, G. & G.J. FIX, *An analysis of the finite element method*, Prentice-Hall, Englewood Cliffs, New Jersey, 1973.

- [6] WHEELER, M.F., *An optional L_∞ error estimate for Galerkin approximations to solutions of two-point boundary problems*, SIAM J. Num. Anal. 10, pp. 914-917 (1973).