

# NAG C Library Function Document

## nag\_1d\_pade (e02rac)

### 1 Purpose

nag\_1d\_pade (e02rac) calculates the coefficients in a Padé approximant to a function from its user-supplied Maclaurin expansion.

### 2 Specification

```
void nag_1d_pade (Integer ia, Integer ib, const double c[], double a[], double b[],
                 NagError *fail)
```

### 3 Description

Given a power series

$$c_0 + c_1x + c_2x^2 + \dots + c_{l+m}x^{l+m} + \dots$$

nag\_1d\_pade (e02rac) uses the coefficients  $c_i$ , for  $i = 0, 1, \dots, l + m$ , to form the  $[l/m]$  Padé approximant of the form

$$\frac{a_0 + a_1x + a_2x^2 + \dots + a_lx^l}{b_0 + b_1x + b_2x^2 + \dots + b_mx^m}$$

with  $b_0$  defined to be unity. The two sets of coefficients  $a_j$ , for  $j = 0, 1, \dots, l$  and  $b_k$ , for  $k = 0, 1, \dots, m$  in the numerator and denominator are calculated by direct solution of the Padé equations (see Graves–Morris (1979)); these values are returned through the argument list unless the approximant is degenerate.

Padé approximation is a useful technique when values of a function are to be obtained from its Maclaurin expansion but convergence of the series is unacceptably slow or even non-existent. It is based on the hypothesis of the existence of a sequence of convergent rational approximations, as described in Baker and Graves–Morris (1981) and Graves–Morris (1979).

Unless there are reasons to the contrary (as discussed in Baker and Graves–Morris (1981) Chapter 4, Section 2, Chapters 5 and 6), one normally uses the diagonal sequence of Padé approximants, namely

$$\{[m/m], m = 0, 1, 2, \dots\}.$$

Subsequent evaluation of the approximant at a given value of  $x$  may be carried out using nag\_1d\_pade\_eval (e02rbc).

### 4 References

Baker G A Jr and Graves–Morris P R (1981) Padé approximants, Part 1: Basic theory *encyclopaedia of Mathematics and its Applications* Addison–Wesley

Graves–Morris P R (1979) The numerical calculation of Padé approximants *Padé Approximation and its Applications. Lecture Notes in Mathematics* (ed L Wuytack) 765 231–245 Addison–Wesley

### 5 Parameters

1: **ia** – Integer Input  
 2: **ib** – Integer Input

*On entry:* **ia** must specify  $l + 1$  and **ib** must specify  $m + 1$ , where  $l$  and  $m$  are the degrees of the numerator and denominator of the approximant, respectively.

*Constraint:* **ia**  $\geq 1$  and **ib**  $\geq 1$ .

- 3: **c[ia + ib - 1]** – const double *Input*  
*On entry:* **c[i - 1]** must specify, for  $i = 1, 2, \dots, l + m + 1$ , the coefficient of  $x^{i-1}$  in the given power series.
- 4: **a[ia]** – double *Output*  
*On exit:* **a[j]**, for  $j = 1, 2, \dots, l + 1$ , contains the coefficient  $a_j$  in the numerator of the approximant.
- 5: **b[ib]** – double *Output*  
*On exit:* **b[k]**, for  $k = 1, 2, \dots, m + 1$ , contains the coefficient  $b_k$  in the denominator of the approximant.
- 6: **fail** – NagError \* *Input/Output*  
 The NAG error parameter (see the Essential Introduction).

## 6 Error Indicators and Warnings

### NE\_INT\_2

On entry, **ib** = *<value>*, **ia** = *<value>*.  
 Constraint: **ia**  $\geq 1$  and **ib**  $\geq 1$ .

### NE\_DEGENERATE

The Pade approximant is degenerate.

### NE\_ALLOC\_FAIL

Memory allocation failed.

### NE\_BAD\_PARAM

On entry, parameter *<value>* had an illegal value.

### NE\_INTERNAL\_ERROR

An internal error has occurred in this function. Check the function call and any array sizes. If the call is correct then please consult NAG for assistance.

## 7 Accuracy

The solution should be the best possible to the extent to which the solution is determined by the input coefficients. It is recommended that the user determines the locations of the zeros of the numerator and denominator polynomials, both to examine compatibility with the analytic structure of the given function and to detect defects. (Defects are nearby pole-zero pairs; defects close to  $x = 0.0$  characterise ill-conditioning in the construction of the approximant.) Defects occur in regions where the approximation is necessarily inaccurate. The example program calls `nag_zeros_real_poly (c02age)` to determine the above zeros.

It is easy to test the stability of the computed numerator and denominator coefficients by making small perturbations of the original Maclaurin series coefficients (e.g.,  $c_l$  or  $c_{l+m}$ ). These questions of intrinsic error of the approximants and computational error in their calculation are discussed in Chapter 2 of Baker and Graves–Morris (1981).

## 8 Further Comments

The time taken is approximately proportional to  $m^3$ .

## 9 Example

The example program calculates the  $[4/4]$  Padé approximant of  $e^x$  (whose power-series coefficients are first stored in the array **c**). The poles and zeros are then calculated to check the character of the  $[4/4]$  Padé approximant.

### 9.1 Program Text

```

/* nag_ld_pade (e02rac) Example Program.
 *
 * Copyright 2001 Numerical Algorithms Group.
 *
 * Mark 7, 2001.
 */

#include <stdio.h>
#include <nag.h>
#include <nag_stdlib.h>
#include <nagc02.h>
#include <nage02.h>

int main(void)
{
    /* Scalars */
    Integer exit_status, i, ifail, l, m, ia, ib, ic;
    NagError fail;

    /* Arrays */
    double *aa = 0, *bb = 0, *cc = 0, *dd = 0;
    Complex *z = 0;

    INIT_FAIL(fail);
    exit_status = 0;
    Vprintf("e02rac Example Program Results\n");

    l = 4;
    m = 4;
    ia = l + 1;
    ib = m + 1;
    ic = ia + ib - 1;

    /* Allocate memory */
    if ( !(aa = NAG_ALLOC(ia, double)) ||
        !(bb = NAG_ALLOC(ib, double)) ||
        !(cc = NAG_ALLOC(ic, double)) ||
        !(dd = NAG_ALLOC(ia + ib, double)) ||
        !(z = NAG_ALLOC(l+m, Complex)) )
    {
        Vprintf("Allocation failure\n");
        exit_status = -1;
        goto END;
    }

    /* Power series coefficients in cc */
    cc[0] = 1.0;
    for (i = 1; i <= 8; ++i)
        cc[i] = cc[i-1] / (double) i;

    Vprintf("\n");

    Vprintf("The given series coefficients are\n");

    for (i = 1; i <= ic; ++i)
    {
        Vprintf("%13.4e", cc[i-1]);
        Vprintf(i%5 == 0 || i == ic ? "\n" : " ");
    }

    ifail = 0;

```

```

e02rac(ia, ib, cc, aa, bb, &fail);
if (fail.code != NE_NOERROR)
{
    Vprintf("Error from e02rac.\n%s\n", fail.message);
    exit_status = 1;
    goto END;
}

Vprintf("\n");
Vprintf("Numerator coefficients\n");

for (i = 1; i <= ia; ++i)
{
    Vprintf("%13.4e", aa[i-1]);
    Vprintf(i%5 == 0 || i == ia ? "\n": " ");
}

Vprintf("\n");
Vprintf("Denominator coefficients\n");

for (i = 1; i <= ib; ++i)
{
    Vprintf("%13.4e", bb[i-1]);
    Vprintf(i%5 == 0 || i == ib ? "\n" : " ");
}

/* Calculate zeros of the approximant using c02agc */
/* First need to reverse order of coefficients */
for (i = 1; i <= ia; ++i)
    dd[ia-i] = aa[i-1];

ifail = 0;
c02agc(1, dd, TRUE, z, &fail);
if (fail.code != NE_NOERROR)
{
    Vprintf("Error from c02agc, 1st call.\n%s\n", fail.message);
    exit_status = 1;
    goto END;
}

Vprintf("\n");
Vprintf("Zeros of approximant are at\n");
Vprintf("    Real part    Imag part\n");
for (i = 1; i <= 1; ++i)
    Vprintf("%13.4e%13.4e\n", z[i-1].re, z[i-1].im);

/* Calculate poles of the approximant using c02agc */
/* Reverse order of coefficients */
for (i = 1; i <= ib; ++i)
    dd[ib-i] = bb[i-1];

ifail = 0;
c02agc(m, dd, TRUE, z, &fail);
if (fail.code != NE_NOERROR)
{
    Vprintf("Error from c02agc, 2nd call.\n%s\n", fail.message);
    exit_status = 1;
    goto END;
}

Vprintf("\n");
Vprintf("Poles of approximant are at\n");
Vprintf("    Real part    Imag part\n");
for (i = 1; i <= m; ++i)
    Vprintf("%13.4e%13.4e\n", z[i-1].re, z[i-1].im);

END:
if (aa) NAG_FREE(aa);
if (bb) NAG_FREE(bb);
if (cc) NAG_FREE(cc);
if (dd) NAG_FREE(dd);

```

```

    if (z) NAG_FREE(z);
    return exit_status;
}

```

## 9.2 Program Data

None.

## 9.3 Program Results

e02rac Example Program Results

The given series coefficients are

1.0000e+00	1.0000e+00	5.0000e-01	1.6667e-01	4.1667e-02
8.3333e-03	1.3889e-03	1.9841e-04	2.4802e-05	

Numerator coefficients

1.0000e+00	5.0000e-01	1.0714e-01	1.1905e-02	5.9524e-04
------------	------------	------------	------------	------------

Denominator coefficients

1.0000e+00	-5.0000e-01	1.0714e-01	-1.1905e-02	5.9524e-04
------------	-------------	------------	-------------	------------

Zeros of approximant are at

Real part	Imag part
-5.7924e+00	1.7345e+00
-5.7924e+00	-1.7345e+00
-4.2076e+00	5.3148e+00
-4.2076e+00	-5.3148e+00

Poles of approximant are at

Real part	Imag part
5.7924e+00	1.7345e+00
5.7924e+00	-1.7345e+00
4.2076e+00	5.3148e+00
4.2076e+00	-5.3148e+00

---