

NAG C Library Chapter Introduction**f01 – Matrix Factorizations****Contents**

1	Scope of the Chapter	2
2	Background to the Problems	2
2.1	Matrix Inversion	2
2.2	Matrix Factorizations	3
3	Recommendations on Choice and Use of Available Functions	3
3.1	Matrix Inversion	3
3.2	Matrix Factorizations	4
4	Decision Trees	4
5	Index	6
6	Functions Withdrawn or Scheduled for Withdrawal	6
7	References	6

1 Scope of the Chapter

This chapter together with Chapters f07, f08 and f11 provides facilities for two types of problem:

- (i) Matrix Inversion (see Chapter f07)
- (ii) Matrix Factorizations (see Chapters f01, f07, f08 and f11)

These problems are discussed separately in Section 2.1 and Section 2.2 .

2 Background to the Problems

2.1 Matrix Inversion

- (i) Non-singular square matrices of order n .

If A , a square matrix of order n , is non-singular (has rank n), then its inverse X exists and satisfies the equations $AX = XA = I$ (the identity or unit matrix).

It is worth noting that if $AX - I = R$, so that R is the ‘residual’ matrix, then a bound on the relative error is given by $\|R\|$, i.e.,

$$\frac{\|X - A^{-1}\|}{\|A^{-1}\|} \leq \|R\|.$$

- (ii) General real rectangular matrices.

A real matrix A has no inverse if it is square (n by n) and singular (has rank $< n$), or if it is of shape (m by n) with $m \neq n$, but there is a **Generalized or Pseudo Inverse** Z which satisfies the equations

$$AZA = A, \quad ZAZ = Z, \quad (AZ)^T = AZ, \quad (ZA)^T = ZA$$

(which of course are also satisfied by the inverse X of A if A is square and non-singular).

- (a) if $m \geq n$ and $\text{rank}(A) = n$ then A can be factorized using a **QR factorization**, given by

$$A = Q \begin{pmatrix} R \\ 0 \end{pmatrix},$$

where Q is an m by m orthogonal matrix and R is an n by n , non-singular, upper triangular matrix. The pseudo-inverse of A is then given by

$$Z = R^{-1} \tilde{Q}^T,$$

where \tilde{Q} consists of the first n columns of Q .

- (b) if $m \leq n$ and $\text{rank}(A) = m$ then A can be factorized using an **RQ factorization**, given by

$$A = (R \ 0)P^T$$

where P is an n by n orthogonal matrix and R is an m by m , non-singular, upper triangular matrix. The pseudo-inverse of A is then given by

$$Z = \tilde{P}R^{-1},$$

where \tilde{P} consists of the first m columns of P .

- (c) if $m \geq n$ and $\text{rank}(A) = r \leq n$ then A can be factorized using a **QR factorization**, with column interchanges, as

$$A = Q \begin{pmatrix} R \\ 0 \end{pmatrix} P^T,$$

where Q is an m by m orthogonal matrix, R is an r by n upper trapezoidal matrix and P is an n by n permutation matrix. The pseudo inverse of A is then given by

$$Z = PR^T (RR^T)^{-1} \tilde{Q}^T,$$

where \tilde{Q} consists of the first r columns of Q .

(d) if $\text{rank}(A) = r \leq k = \min(m, n)$, then A can be factorized as the **singular value decomposition**

$$A = QDP^T,$$

where Q is an m by m orthogonal matrix, P is an n by n orthogonal matrix and D is an m by n diagonal matrix with non-negative diagonal elements. The first k columns of Q and P are the **left- and right-hand singular vectors** of A respectively and the k diagonal elements of D are the **singular values** of A . D may be chosen so that

$$d_1 \geq d_2 \geq \cdots \geq d_k \geq 0$$

and in this case if $\text{rank}(A) = r$ then

$$d_1 \geq d_2 \geq \cdots \geq d_r > 0, \quad d_{r+1} = \cdots = d_k = 0.$$

If \tilde{Q} and \tilde{P} consist of the first r columns of Q and P respectively and Σ is an r by r diagonal matrix with diagonal elements d_1, d_2, \dots, d_r then A is given by

$$A = \tilde{Q}\Sigma\tilde{P}^T$$

and the pseudo inverse of A is given by

$$Z = \tilde{P}\Sigma^{-1}\tilde{Q}^T.$$

Notice that

$$A^T A = P(D^T D)P^T$$

which is the classical eigenvalue (spectral) factorization of $A^T A$.

(e) if A is complex then the above relationships are still true if we use ‘unitary’ in place of ‘orthogonal’ and conjugate transpose in place of transpose. For example, the singular value decomposition of A is

$$A = QDP^H,$$

where Q and P are unitary, P^H the conjugate transpose of P and D is as in (d) above.

2.2 Matrix Factorizations

The functions in this section perform matrix factorizations which are required for the solution of systems of linear equations with various special structures. A few functions which perform associated computations are also included.

Other functions for matrix factorizations are to be found in Chapters f03, f07, f08 and f11.

3 Recommendations on Choice and Use of Available Functions

3.1 Matrix Inversion

Note: before using any function for matrix inversion, consider carefully whether it is really needed.

Although the solution of a set of linear equations $Ax = b$ can be written as $x = A^{-1}b$, the solution should **never** be computed by first inverting A and then computing $A^{-1}b$; the functions in Chapters f04 or f07 should **always** be used to solve such sets of equations directly; they are faster in execution, and numerically more stable and accurate. Similar remarks apply to the solution of least-squares problems which again should be solved by using the functions in Chapters f02 or f08 rather than by computing a pseudo inverse.

(a) Non-singular square matrices of order n

This chapter describes techniques for inverting a general real matrix A and matrices which are positive-definite (have all eigenvalues positive) and are either real and symmetric or complex and Hermitian. It is wasteful and uneconomical not to use the appropriate function when a matrix is known to have one of these special forms. A general function must be used when the matrix is not known to be positive-definite. In most functions the inverse is computed by solving the linear equations $Ax_i = e_i$, for $i = 1, 2, \dots, n$, where e_i is the i th column of the identity matrix.

The residual matrix $R = AX - I$, where X is a computed inverse of A , conveys useful information in that $\|R\|$ is a bound on the relative error in X .

The decision trees for inversion show which functions in Chapter f07 should be used for the inversion of other special types of matrices not treated in the chapter.

(b) General real rectangular matrices

For real matrices `nag_dgeqrf` (f08aec) returns a QR factorization of A and `nag_dgeqpf` (f08bec) returns the QR factorization with column interchanges. The corresponding complex functions are `nag_zgeqrf` (f08asc) and `nag_zgeqpf` (f08bsc) respectively. Functions are also provided to form the orthogonal matrices and transform by the orthogonal matrices following the use of the above functions.

Combinations of functions in Chapter f08 (e.g., `nag_dgebrd` (f08kec), `nag_dorgbr` (f08kfc) and `nag_dbdsqr` (f08mec)) compute the singular value decomposition as described in Section 2 for real and complex matrices respectively. If A has rank $r \leq k = \min(m, n)$ then the $k - r$ smallest singular values will be negligible and the pseudo inverse of A can be obtained as $Z = P\Sigma^{-1}Q^T$ as described in Section 2. If the rank of A is not known in advance it can be estimated from the singular values (see Section 2.2 in the f04 Chapter Introduction). For large sparse matrices, leading terms in the singular-value decomposition can be computed using functions from Chapter f12.

3.2 Matrix Factorizations


Each of these functions serves a special purpose required for the solution of sets of simultaneous linear equations or the eigenvalue problem. For further details you should consult Sections 3 or 4 in the f02 Chapter Introduction or Sections 3 or 4 in the f04 Chapter Introduction.



For the factorization of sparse matrices, see `nag_sparse_nsym_fac` (f11dac), `nag_superlu_lu_factorize` (f11mec) and `nag_sparse_sym_chol_fac` (f11jac). These functions should be used only when A is **not** banded and when the total number of non-zero elements is less than 10% of the total number of elements. In all other cases either the band functions or the general functions should be used.


4 Decision Trees

The decision trees show the functions in this chapter and in Chapter f04 that should be used for inverting matrices of various types. Functions marked with an asterisk (*) only perform part of the computation – see Section 3.1 for further advice.

Tree 1



Tree 2: Inverse of a real n by n matrix of full rank**Tree 3: Inverse of a complex n by n matrix of full rank**

Tree 4: Pseudo-inverses

Note 1: the inverse of a band matrix A does not in general have the same shape as A , and no functions are provided specifically for finding such an inverse. The matrix must either be treated as a full matrix, or the equations $AX = B$ must be solved, where B has been initialized to the identity matrix I . In the latter case, see the decision trees in Section 4 in the f04 Chapter Introduction.

5 Index

Matrix Transformations,

complex Hermitian positive-definite matrix, UU^H factorization	nag_complex_cholesky (f01bnc)
complex m by n matrix ($m \leq n$), QR factorization nag_complex_qr (f01rcc)
complex matrix,	
apply orthogonal matrix nag_complex_apply_q (f01rdc)
form unitary matrix nag_complex_form_q (f01rec)
real band symmetric positive-definite matrix,	
variable bandwidth, LDL^T factorization nag_real_cholesky_skyline (f01mcc)
real m by n matrix ($m \leq n$), QR factorization nag_real_qr (f01qcc)
real matrix,	
apply orthogonal matrix nag_real_apply_q (f01qdc)
form orthogonal matrix nag_real_form_q (f01qec)

6 Functions Withdrawn or Scheduled for Withdrawal

None.

7 References

Golub G H and Van Loan C F (1996) *Matrix Computations* (3rd Edition) Johns Hopkins University Press, Baltimore

Wilkinson J H (1965) *The Algebraic Eigenvalue Problem* Oxford University Press, Oxford

Wilkinson J H (1977) Some recent advances in numerical linear algebra *The State of the Art in Numerical Analysis* (ed D A H Jacobs) Academic Press

Wilkinson J H and Reinsch C (1971) *Handbook for Automatic Computation II, Linear Algebra* Springer-Verlag