

Update 2 of: Electrophilicity Index

Pratim Kumar Chattaraj*, Santanab Giri, and Soma Duley

Department of Chemistry and Center for Theoretical Studies, Indian Institute of Technology, Kharagpur 721302, India

This is a Chemical Reviews Perennial Review. The root paper of this title was published in Chem. Rev. 2006, 106, 2065–2091, DOI: 10.1021/cr040109f; Published (Web) May 25, 2006. Update 1 of this title was published in Chem. Rev. 2007, 107, PR46–PR74, DOI: 10.1021/cr078014b; Published (Web) September 12, 2007. Updates to the text appear in red type.

Received May 17, 2010

Contents

1. Introduction: Electrophilic and Nucleophilic Processes—Reaction Mechanisms, Thermodynamics, and Kinetics	PR43
2. Global Electrophilicity Index	PR44
2.1. Genesis	PR44
2.2. Prescription	PR45
3. Local Extensions and Site Selectivity	PR46
3.1. Local Electrophilicity	PR46
3.2. Site Selectivity	PR48
4. Electrophilicity Scales	PR48
4.1. Global Approach	PR48
4.2. Local Approach	PR50
5. Intramolecular and Intermolecular Reactivity Patterns	PR51
5.1. Periodicity	PR51
5.2. Excited States	PR51
5.3. Spherical Confinement	PR51
5.4. Chemical Processes	PR52
5.5. Solvent Effects	PR54
5.6. External Field Effects	PR55
5.7. Biological Activity and Toxicity	PR55
6. Variation of the Electrophilicity Index during Physicochemical Processes Such as Vibrations, Internal Rotations, and Chemical Reactions	PR58
6.1. Molecular Vibrations	PR58
6.2. Molecular Internal Rotations	PR58
6.3. Chemical Reactions	PR58
7. Dynamical Variants	PR62
7.1. Quantum Fluid Density Functional Theory	PR62
7.2. Atom—Field Interactions	PR62
7.3. Ion—Atom Collisions	PR62
7.4. Chemical Kinetics	PR62
8. Spin Dependent Generalizations	PR63
8.1. $\{N, N_s, \nu(\vec{r})\}$ Representation	PR63
8.2. $\{N_\alpha, N_\beta, \nu(\vec{r})\}$ Representation	PR64
9. Conclusions	PR65
10. Abbreviations and Symbols of Some Important Subjects/Quantities	PR66
11. Acknowledgments	PR67
12. References	PR67

1. Introduction: Electrophilic and Nucleophilic Processes—Reaction Mechanisms, Thermodynamics, and Kinetics

Chemistry is the science of bond making and bond breaking. A thorough knowledge of these processes in the course of the chemical reaction lies at the heart of any reaction mechanism. In the heterolytic cleavage of a bond, the electron pair lies with one of the fragments, which becomes electron rich, while the other fragment becomes electron deficient. An electron-rich reagent gets attracted to the center of the positive charge and forms a bond with an electron-deficient species by donating electrons. The electron-rich species is known as a nucleophile, and the electron-deficient one, as an electrophile.^{1–4}

Free radicals are generated through a corresponding homolytic process where an equal share of one electron is obtained by each fragment. Even radicals are designated as electrophilic/nucleophilic depending on their tendency to attack the reaction sites of relatively higher/lower electron density. Moreover, nucleophiles (electrophiles) are Lewis bases (acids) as well as reducing (oxidizing) agents since they donate (accept) electrons, implying a connection among electrophile–nucleophile chemistry, acid–base chemistry, and oxidation–reduction chemistry. Since the majority of the reactions can be analyzed through the electrophilicity/nucleophilicity of various species involved, a proper understanding of these properties becomes essential. Some related reviews on specific types of reactions are available,^{5–13} albeit without a rigorous definition of electrophilicity.

The most important types of reactions we often encounter are substitution, addition (including pericyclic reactions), elimination (the opposite of addition reactions), and rearrangements. All these reactions are analyzed using thermodynamic and kinetic data. While the former determines how far a reaction will go (a large decrease in Gibbs free energy implies a large value of the equilibrium constant, K), the latter determines how fast it will take place (a smaller free energy of activation value implies a larger rate constant, k , at a given temperature, i.e., a faster reaction). Although the electrophilicity (nucleophilicity) and Lewis acidity (basicity) are related, the former is traditionally assumed to be a kinetic quantity and, hence, is estimated by relative k values whereas the latter is a thermodynamic quantity and is measured by relative K values.

The concept of electrophilicity has been known for several decades, although there has not been a rigorous definition of it until recently, when, inspired by the experimental findings of Maynard et al.,¹³ Parr et al.¹⁴ proposed a definition

* To whom correspondence should be addressed. E-mail: pkc@chem.iitkgp.ernet.in. Phone: 91-03222-283304 (office); 91-03222-283305 (residence). Fax: 91-3222-255303.



After obtaining his B.Sc. and M.Sc. degrees from Burdwan University and his Ph.D. degree from Indian Institute of Technology (Bombay), Pratik Kumar Chattaraj joined the faculty of Indian Institute of Technology (Kharagpur). He is now a professor and the Head of the Department of Chemistry and also the Convener of the Center for Theoretical Studies there. In the meantime, he visited the University of North Carolina (Chapel Hill) as a postdoctoral research associate and several other universities throughout the world as a visiting professor. Apart from teaching, Professor Chattaraj is involved in research on density functional theory, the theory of chemical reactivity, aromaticity in metal clusters, ab initio calculations, quantum trajectories, and nonlinear dynamics. He has been invited to deliver special lectures at several international conferences and to contribute chapters to many edited volumes. Professor Chattaraj is a member of the editorial board of *J. Mol. Struct. (Theochem)* and *J. Chem. Sci.* among others. He is a council member of the Chemical Research Society of India and a Fellow of the Indian Academy of Sciences (Bangalore), the Indian National Science Academy (New Delhi), the National Academy of Sciences, India (Allahabad), and the West Bengal Academy of Science and Technology. He is a J. C. Bose National Fellow. He has edited three books, *Chemical reactivity theory: A density functional view*, *Aromaticity and metal clusters*, and *Quantum trajectories*, published by Taylor and Francis Books, Inc./CRC Press, Boca Raton, FL, and a special issue each of *J. Chem. Sci.* (on "Chemical Reactivity") and *J. Mol. Struct. (Theochem)* (with Professor A. J. Thakkar, on "Conceptual aspects of electron densities and density functionals"). He is the Editor of the book series on "Atoms, Molecules, and Clusters: Structure, Reactivity, and Dynamics" published by Taylor and Francis Books, Inc./CRC Press, Boca Raton, FL, USA.



Santanab Giri received his B.Sc. (Chemistry Honors) and M.Sc. (Chemistry) degrees from the Vidyasagar University, West Bengal. Then he joined the research group of Professor P. K. Chattaraj for his Ph.D. degree in the Department of Chemistry, Indian Institute of Technology, Kharagpur, as an Institute Fellow. At present he is a Senior Research Fellow in a CSIR (Government of India) scheme. He is a recipient of two national awards (Professor Santi Ranjan Palit award and Sir P. C. Ray award).

based on the energy lowering associated with a maximum amount of electron flow between two species. There resulted an upsurge of interest in understanding the utility of this quantity in analyzing several diverse areas of chemistry. Little



Soma Duley received her B.Sc. degree (Chemistry Honors) from the Vidyasagar University and M.Sc. degree (Chemistry) from Indian Institute of Technology, Kharagpur. Then she joined the research group of Professor P. K. Chattaraj for her Ph.D. degree in the Department of Chemistry, Indian Institute of Technology, Kharagpur, as a CSIR (Government of India) Fellow. She received a certificate of merit in the Acharya Prafulla Chandra Ray Memorial Symposium.

did they¹⁴ realize when they proposed this simple-looking index that it contains an enormous potential to connect the major facets of chemical sciences. It has been shown that electrophilicity possesses adequate information regarding structure, stability, reactivity, toxicity, bonding, interactions, and dynamics. Only the concept of electrophilicity index provided by Parr et al.,¹⁴ its usefulness, and its various possible extensions will be reviewed in this article. Citations of the works related to the Maynard–Parr electrophilicity index are intended to be as exhaustive as possible with mention of and necessary comparison with all other known popular electrophilicity measures. Several papers related to this index appeared after the final submission of this article. They are cited in appropriate places in the References section. Section 2 introduces this global index, and its local variants are described in section 3. Existing electrophilicity scales are reported in section 4. Section 5 describes the use of the electrophilicity index in analyzing the reactivity patterns in various intramolecular and intermolecular physicochemical processes. Variation of this quantity during molecular vibrations, internal rotations, and chemical reactions is mentioned in section 6. Sections 7 and 8 report the dynamical and spin dependent variants of this index, respectively. Finally, section 9 contains some concluding remarks.

2. Global Electrophilicity Index

2.1. Genesis

Popular qualitative chemical concepts such as electronegativity^{15,16} (χ) and hardness^{17,18} (η) have been provided with rigorous definitions within the purview of conceptual density functional theory^{19–28} (DFT). Electronegativity is the negative of chemical potential defined²⁹ as follows for an N -electron system with total energy E and external potential $v(\vec{r})$,

$$\chi = -\mu = -\left(\frac{\partial E}{\partial N}\right)_{v(\vec{r})} \quad (1)$$

μ is the Lagrange multiplier associated with the normalization constraint of DFT.^{19,27,28} In DFT, the electron density ($\rho(\vec{r})$) is the basic variable instead of the many-particle wave function ($\Psi(\vec{x}_1, \vec{x}_2, \dots, \vec{x}_N)$).

Hardness (η) is defined³⁰ as the corresponding second derivative,

$$\eta = \left(\frac{\partial^2 E}{\partial N^2} \right)_{v(\vec{r})} = \left(\frac{\partial \mu}{\partial N} \right)_{v(\vec{r})} \quad (2)$$

Sometimes a $1/2$ factor is included in the above definition. Softness (S) is the reciprocal of hardness; $S = 1/\eta$. **The above definitions suffer from the derivative discontinuity problem.³¹ A grand canonical ensemble picture may remedy this problem.^{31–33}**

Complete characterization of an N -particle wave function needs only N and $v(\vec{r})$. While χ and η measure the response of the system when N varies at constant $v(\vec{r})$, the behavior of the system for a change in $v(\vec{r})$ at fixed N is given by the linear density response function.¹⁹ The linear response of the electronic cloud of a chemical species to a weak external electric field is measured in terms of the static electric dipole polarizability (α). The electric dipole polarizability is a measure of the linear response of the electron density in the presence of an electric field F , and it represents a second-order variation in energy,

$$\alpha_{a,b} = - \left(\frac{\partial^2 E}{\partial F_a \partial F_b} \right); \quad a, b = x, y, z \quad (3)$$

The polarizability (α) is then calculated as follows

$$\langle \alpha \rangle = 1/3 (\alpha_{xx} + \alpha_{yy} + \alpha_{zz}) \quad (4)$$

These reactivity parameters are better appreciated by various associated electronic structure principles. According to the electronegativity equalization principle,^{29,34–36} “All the constituent atoms in a molecule have the same electronegativity value given by the geometric mean of the electronegativities of the pertinent isolated atoms”. Two hardness related principles are the hard–soft acids and bases (HSAB) principle^{17,30,37–43} and the maximum hardness principle.^{44–51} While the former states that, “Hard acids prefer to coordinate with hard bases and soft acids to soft bases for their thermodynamic and kinetic properties”, the statement of the latter is, “There seems to be a rule of nature that molecules arrange themselves so as to be as hard as possible”. On the basis of the inverse relationship, $\eta \propto 1/\alpha^{1/3}$,^{52–55} between hardness and polarizability, a minimum polarizability principle^{56–59} has been proposed which states that,⁵⁶ “The natural direction of evolution of any system is toward a state of minimum polarizability”. An inverse relationship between hardness and magnetizability has also been shown,^{52b,c} which leads to a minimum magnetizability principle. A soft molecule is both polarizable and magnetizable. **Like electronegativity and hardness, electrophilicity also gets equalized during a chemical reaction.⁶⁰ The molecular electrophilicity may be roughly expressed as the geometric mean of the corresponding isolated atom values.⁶⁰**

Using a finite difference method, working equations for the calculation of χ and η may be given as¹⁹

$$\chi = \frac{I + A}{2} \quad (5)$$

$$\eta = I - A \quad (6)$$

where I and A are the ionization potential and electron affinity, respectively. If ϵ_{HOMO} and ϵ_{LUMO} are the energies of

the highest occupied and lowest unoccupied molecular orbitals, respectively, then the above equations can be rewritten⁶¹ using Koopmans’ theorem⁶² as

$$\chi = - \frac{\epsilon_{\text{HOMO}} + \epsilon_{\text{LUMO}}}{2} \quad (7)$$

$$\eta = \epsilon_{\text{LUMO}} - \epsilon_{\text{HOMO}} \quad (8)$$

Maynard and co-workers⁶³ have shown that the reaction rates from the fluorescence decay studies on the human immunodeficiency virus type 1 (HIV-1) nucleocapsid protein p7 (NCp 7) interacting with several electrophilic agents correlate strongly with the square of the electronegativity divided by its chemical hardness ($\chi^2/2\eta$). The $1/2$ factor arises due to the definition of hardness used here (eq 2). The quantity $\chi^2/2\eta$ is considered^{13,63} to be related to the capacity of an electrophile to promote a soft (covalent) reaction. Prompted by this work of Maynard and co-workers,^{13,63} an electrophilicity index (ω) has been defined as $\chi^2/2\eta$ by Parr et al.¹⁴ They have shown¹⁴ that ω measures the second-order energy of an electrophile when it gets saturated with electrons.

2.2. Prescription

To propose an electrophilicity index, Parr et al.¹⁴ assumed a sea of free electron gas at zero temperature and zero chemical potential. When an electrophilic system (atom, molecule, or ion) would get immersed into the sea, there would be an electron flow of amount ΔN from the sea to the system until the chemical potential of the system becomes zero (cf. Sanderson’s principle,^{29,34–36}). The resulting energy change (up to second order) associated with the electron-transfer process is

$$\Delta E = \mu \Delta N + \frac{1}{2} \eta (\Delta N)^2 \quad (9)$$

Considering the situation when the system becomes saturated by soaking up the maximum amount of electrons, ΔN_{max} , they set¹⁴ ($\Delta E/\Delta N$) to be zero, implying

$$\Delta E = - \frac{\mu^2}{2\eta} \quad (10)$$

and

$$\Delta N_{\text{max}} = - \frac{\mu}{\eta} \quad (11)$$

In eq 10, the numerator (μ^2) is quadratic and, hence, positive and the denominator (2η) is positive due to the convexity of the energy, and hence, ΔE is negative: the charge transfer is an energetically favorable process. Parr et al.¹⁴ defined ω as

$$\omega = \frac{\mu^2}{2\eta} = \frac{\chi^2}{2\eta} \quad (12)$$

as a measure of electrophilicity of the ligand, just as was suggested by Maynard and co-workers.⁶³

This quantity ω is called¹⁴ the “electrophilicity index” and is considered to be a measure of electrophilic power, just as, in classical electrostatics, power = V^2/R , and μ and η serve the purpose of potential (V) and resistance (R),




Figure 1. Correlation between electrophilicity index and electron affinity of some neutral atoms and simple molecules in the ground-state parabola model. Reprinted with permission from ref 14. Copyright 1999 American Chemical Society.




Figure 2. Correlation between electronegativities and electrophilicity indices for the functional groups XY_3 ($X = C, Si, Ge, Sn, Pb$, element 114; $Y = CH_3, H, F, Cl, Br, I, At$). Reprinted with permission from ref 64. Copyright 2005 American Chemical Society.




Figure 3. Correlation between electron affinities and electrophilicity indices for the functional groups XY_3 ($X = C, Si, Ge, Sn, Pb$, element 114; $Y = CH_3, H, F, Cl, Br, I, At$). Reprinted with permission from ref 64. Copyright 2005 American Chemical Society.

respectively. It is transparent from Figures 1–3 that ω and A are not equal but they are correlated¹⁴ and ω is more tightly correlated⁶⁴ with A than with χ , though all these quantities measure the propensity of electron intake. Since ω depends on both I and A , it is expected that A can provide similar qualitative trends as ω whenever the variation in I is not

very significant. This is commonly observed for the elements belonging to the same group in the periodic table⁶⁴ and the functional groups containing them. It may be noted that A takes care of the energy change due to the addition of a single electron whereas the energy lowering associated with maximal electron flow is characterized by ω . However, for the systems like dianions that have negative electronegativity (χ) values, the electrophilicity (ω) remains high although the system does not like to take electrons. This may be considered to be a drawback of the quadratic appearance⁶⁵ of χ in the expression of ω (eq 12). Considering the discontinuity in the E vs N curve,^{63b} separate definitions for electrophilicity have been proposed^{63c} for the charge donation and charge acceptance processes, respectively, as $\omega^\pm = (\mu^\pm)^2/2\eta$, where μ^\pm are the corresponding left and right derivatives obtained in the spirit of the Perdew, Parr, Levy, and Balduz prescription.^{63b}

In the present review we will restrict ourselves to various ramifications of ω only. An update is provided here as a Perennial Review to the original article, *Chem. Rev.* **2006**, *106*, 2065, and Update 1 of the original, *Chem. Rev.* **2007**, *107*, PR46.

3. Local Extensions and Site Selectivity

3.1. Local Electrophilicity

Global reactivity descriptors such as electronegativity, chemical potential, hardness, polarizability, and electrophilicity as introduced in the last section are defined for the system as a whole. To describe the site selectivity in a molecule, local descriptors of reactivity have also been proposed. An equivalent definition of hardness has been given by^{66,67}

$$\eta = \frac{1}{N} \int \int \eta(\vec{r}, \vec{r}') f(\vec{r}') \rho(\vec{r}) d\vec{r} d\vec{r}' \quad (13)$$

or its other variants^{68–70} where $f(\vec{r})$ is the Fukui function^{71–75} and the hardness kernel can be written as

$$\eta(\vec{r}, \vec{r}') = \frac{\delta^2 F[\rho]}{\delta \rho(\vec{r}) \delta \rho(\vec{r}')} \quad (14)$$

where $F[\rho]$ is the Hohenberg–Kohn–Sham^{27,28} universal functional.

The Fukui function is by far the most important local reactivity index. It is defined as⁷⁴

$$f(\vec{r}) \equiv (\partial \rho(\vec{r}) / \partial N)_{v(\vec{r})} = (\delta \mu / \delta v(\vec{r}))_N \quad (15)$$

Because of the discontinuities in slope of the $\rho(\vec{r})$ versus N curve,^{76,63b} three types of Fukui functions can be written as follows⁷⁴

for nucleophilic attack

$$f^+(\vec{r}) = \left(\frac{\partial \rho(\vec{r})}{\partial N} \right)_{v(\vec{r})}^+ \approx \rho_{N+1}(\vec{r}) - \rho_N(\vec{r}) \approx \rho_{\text{LUMO}}(\vec{r}) \quad (16a)$$

for electrophilic attack

$$f^-(\vec{r}) = \left(\frac{\partial \rho(\vec{r})}{\partial N} \right)_{v(\vec{r})}^- \approx \rho_N(\vec{r}) - \rho_{N-1}(\vec{r}) \approx \rho_{\text{HOMO}}(\vec{r}) \quad (16b)$$

for radical attack

$$f^0(\vec{r}) = \frac{1}{2}[f^+(\vec{r}) + f^-(\vec{r})]$$

which capture the essence of Fukui's frontier orbital theory. A gradient correction method^{77,78} and a variational approach^{79,80} for the calculation of $f(\vec{r})$ are known.

Other important local reactivity descriptors include $\nabla \rho(\vec{r})$ and $\nabla^2 \rho(\vec{r})$ (as analyzed by Bader and co-workers),^{81,82} the molecular electrostatic potential,^{83–86} and the quantum potential.^{87,88} The quantum potential has been defined as

$$V_{\text{qu}}(\vec{r}, t) = -\frac{1}{2} \frac{\nabla^2 \rho^{1/2}(\vec{r}, t)}{\rho^{1/2}(\vec{r}, t)} \quad (17)$$

Two useful theories based on V_{qu} are quantum fluid dynamics (QFD)⁸⁷ and quantum theory of motion (QTM).⁸⁸ In the former, the dynamics of a quantum system is mapped onto that of a probability fluid under the guidance of a classical potential augmented by a quantum potential, while in QTM it is represented in terms of an ensemble of particle motions governed by forces originating from both classical and quantum potentials. In section 7 we will report the chemical reactivity dynamics by making use of these concepts.

Another important local reactivity descriptor is the electron localization function (ELF), which has been defined^{89–92} for a single determinantal wave function in terms of various kinetic energy densities (Kohn–Sham, Weizsäcker and Thomas–Fermi) or, equivalently, the related local temperatures⁹² as

$$\text{ELF} = \frac{1}{1 + (t/t_{\text{F}})^2} \quad (18a)$$

where

$$t = \frac{1}{2} \sum_i |\nabla \Psi_i|^2 - \frac{1}{8} \frac{|\nabla \rho|^2}{\rho} \quad (18b)$$

and

$$t_{\text{F}} = \frac{3}{10} (3\pi^2)^{2/3} \rho^{5/3} \quad (18c)$$

Sometimes it becomes difficult to analyze site selectivity using these local \vec{r} dependent quantities. To tackle this problem, the related condensed-to-atom variants are written for the atomic site k of the molecule. For example, the corresponding Fukui functions (f_k^α , $\alpha = +, -, 0$) can be written⁹³ by replacing the associated electron densities by the respective electron populations (q_k); viz.,

$$\text{for nucleophilic attack} \\ f_k^+ = q_k(N + 1) - q_k(N) \quad (19a)$$

for electrophilic attack

$$f_k^- = q_k(N) - q_k(N - 1) \quad (19b)$$

for radical attack

$$f^0(\vec{r}) = \frac{1}{2}[f_k^+ + f_k^-] \quad (19c)$$

To tackle the hard–soft interactions better, local softnesses have been defined as^{19,93}

$$S^a(\vec{r}) = S f^a(\vec{r}) \quad (20a)$$

and

$$S_k^\alpha = S f_k^\alpha \quad (20b)$$

where $\alpha = +, -, \text{ and } 0$ refer to nucleophilic, electrophilic, and radical reactions, respectively. A local version of the HSAB principle has been proposed^{94,95} in terms of these quantities.

On the other hand, a local electrophilicity has been introduced to analyze the electrophile–nucleophile reactions better. It is defined as^{96,97}

$$\omega_k = \frac{\mu^2}{2} s_k^+ = \frac{\mu^2 S}{2} f_k^+ = \omega f_k^+ \quad (21)$$

A generalized version of this quantity has been termed as philicity, which has been defined through the resolution of the identity associated with the normalization of the Fukui function as⁹⁸

$$\omega = \omega \int f(\vec{r}) d\vec{r} = \int \omega f(\vec{r}) d\vec{r} = \int \omega(\vec{r}) d\vec{r} \quad (22a)$$

where

$$\omega(\vec{r}) = \omega f(\vec{r}) \quad (22b)$$

Note that $\omega(\vec{r})$ can provide both ω (using eq 22a) and $f(\vec{r})$ (using eq 22b along with ω obtained from eq 22a) but $f(\vec{r})$ needs an explicit knowledge of ω to give $\omega(\vec{r})$. Moreover, $\omega(\vec{r})$ can provide $s(\vec{r})$, S , and η with an input of μ . The corresponding condensed-to-atom variants may be defined as⁹⁸

$$\omega_k^\alpha = \omega f_k^\alpha; \quad \alpha = +, -, 0 \quad (22c)$$

Equations 20 and 21 dictate eq 22b to be the natural choice. In the place of $f(\vec{r})$, one may use other normalized-to-one quantities such as the shape function, $\sigma(\vec{r}) = \rho(\vec{r})/N$, which, however, may not be a better descriptor due to the lack of information regarding electron addition/removal. Prominent shell structures in the radial distribution of the philicity of halogen atoms are depicted in Figure 4. A strong rationale for the above definition of philicity has been provided recently by Gázquez et al.^{63c} They have started^{63c} from a functional Taylor expansion of the energy functional followed by the Ayers–Parr type minimization procedure⁴⁸ based on the Chattaraj–Cedillo–Parr variational principle⁷⁹ to obtain the above definition of philicity⁹⁸ and its generalization, $\omega_k^\pm = \omega^\pm f_k^\pm$, strengthening the original definition obtained through the resolution of identity associated with the normalization of $f(\vec{r})$. Of course, the two-parabola model^{63c} may not be appropriate in all cases. It may be noted




Figure 4. Radial distribution of philicity in the ground states of halogen atoms. Reprinted with permission from ref 98. Copyright 2003 American Chemical Society.

that the condition^{63c} $\omega^+ = \omega^- = \omega$ is valid^{63d} when there is a transfer of a fraction (noninteger number) of an electron, as is the case¹⁴ with the electrophilicity as opposed to the electron affinity, which involves exactly one electron; hence the former does not suffer^{63d} from the derivative discontinuity problem.^{63b}

The local electrophilicity, $\omega(\vec{r})$, obeys an exact cusp condition given by^{24c}

$$\lim_{r \rightarrow 0} \frac{\vec{r} \cdot \nabla \omega(\vec{r})}{\omega(\vec{r})} = -2Z \quad (23)$$

It is important to note that the Fukui function and the related quantities such as s_k^α and ω_k^α may not provide proper reactivity trends for hard–hard interactions,^{99–101} as was long ago pointed out by Klopman.¹⁰² Hard–hard interactions are charge-controlled since they are ionic in nature, whereas soft–soft interactions are frontier-controlled because of their covalent nature. The charge-based descriptors would be better suited to tackle the hard–hard interactions.^{99–101} Although ambiguous,^{68,69} a local hardness has been shown⁷⁰ to be a better descriptor of hard–hard reactions than the Fukui function.

3.2. Site Selectivity

An analysis of philicity (ω_k^α) provides the local information of a particular atomic site in a molecule being prone to electrophilic, nucleophilic, or radical attack.⁹⁸ The global electrophilicity of a molecule is determined by these local properties,^{96,98,103} as was suggested by Legon^{104,105} in his electrostatic model.

The local reactivity of various atomic sites in a molecule can be understood equivalently by f_k^α , s_k^α , or ω_k^α (or its variants proposed by Gázquez et al.^{63c}) because S and ω (or ω^\pm) remain the same except for the cases where the molecule is undergoing an intramolecular process such as vibration, internal rotation, rearrangement, and/or interaction with a solvent or an external field where both the local and global descriptors change during the physicochemical process.

Philicity and local softness, respectively, would be better intermolecular reactivity indices (because they are products of global and local indices) than the Fukui function for analyzing electrophile–nucleophile interactions and hard–soft interactions. Recently, some of these aspects have been numerically verified.^{106,107} Philicity and local softness es-

entially provide the same information that is provided by the Fukui function regarding intramolecular reactivity trends except for the intramolecular processes where ω and/or S also changes along with $f(\vec{r})$. However, for analyzing the intermolecular reactivity, f_k^α would be inadequate and s_k^α (or ω_k^α) should be used to compare the hard–soft (electrophilic–nucleophilic) behavior of a given atomic site in one molecule with that of another atomic site in another molecule. For the same molecule, f_k^α is adequate. While f_k^α will take care of reactivity variations on all atoms present in a molecule (and will not be able to differentiate between the reactivities of different molecules), the global quantities (S , ω , ω^\pm , etc.) present in the definition will differentiate between the global reactivity behavior of two (or more) molecules, and hence, the product may be ideal for describing the intermolecular reactivity trends. Moreover, the global reactivity stems from the local reactivities since $X = \sum_k X_k^\alpha = \sum_k X f_k^\alpha = X \sum_k f_k^\alpha$; $X = S$, ω , ω^\pm , etc., which implies that a simple addition of reactivities at all atomic sites will provide the global reactivity. In most of the molecules, a single (or a few) atom(s) is (are) overwhelmingly more reactive (with comparatively very large f_k^α values) than the rest and hence will provide an estimate of the global reactivity. Of course, the domain of applicability of ω_k^α (or its variants proposed by Gázquez et al.^{63c}) and s_k^α as intermolecular descriptors needs to be analyzed in detail. Moreover, the numerical values of any condensed-to-atom quantity and the resulting trends should be analyzed with caution as they are empirical in nature owing to their dependence on the density partitioning scheme used.

During an electrophile–nucleophile interaction process, when two reactants approach each other from a large distance, they feel only the effect of the global electrophilicity of each other and not its local counterpart. The molecule with the larger ω value will act as an electrophile, and the other will behave as the nucleophile. The preferred interaction will be through the most electrophilic site of the former and the most nucleophilic site of the latter. The atom with the largest local electrophilicity value in the electrophile may not necessarily have this value larger than that of the nucleophile, especially when there is more than one active electro(nucleo)philic site present in a molecule. Similar situations arise during an analysis of the corresponding local and global softnesses, and the HSAB principle^{37–43} can be at variance with its local counterpart^{94,95} in those cases.

4. Electrophilicity Scales

4.1. Global Approach

Ever since Ingold¹⁰⁸ proposed an electrophilicity scale, various experimental and theoretically calculated quantities have been made use of in analyzing the electrophilicity behavior of a group of molecules. Both electrophilicity and nucleophilicity have been estimated^{104,105} through hydrogen-bond stretching force constants measured from the rotational spectra of various hydrogen-bonded dimers. The hydrogen-bond strength given by k is related to nucleophilicities (N) and electrophilicities (E) as

$$k_\sigma = CNE \quad (24)$$

where C is a proportionality constant. It has been shown that $N \propto 1/E$ for a fixed value of k_σ , as expected. Here, the force constant is considered to be a measure of the binding strength




Figure 5. Correlation between experimental electrophilicity (E) and theoretical electrophilicity (ω) of a series of benzene diazonium ion and its derivatives containing a large variety of electron-releasing and electron-withdrawing groups in the *ortho*- and *para*-positions. Reprinted with permission from ref 133. Copyright 2003 American Chemical Society.

between an electrophile and a nucleophile and is an alternative to the corresponding bond dissociation energy, which is conventionally used to describe the electrophilic power. A comparison of E and ω has been made.¹⁰³ Activation hardness¹⁰⁹ and protonation energies¹¹⁰ have been used to analyze electrophilic aromatic substitution reactions.

One of the most popular electrophilicity scales has been proposed and used to explain diverse types of reactions by Mayr and co-workers.^{111–125} They^{111–125} have demonstrated through studies on a series of electrophile–nucleophile combination reactions that the absolute rate constants of these reactions follow the following linear free energy relationship,

$$\log k(20^\circ\text{C}) = s(N + E) \quad (25)$$

where E and N are respectively the electrophilicity and nucleophilicity parameters and s is a nucleophile-specific slope parameter. This scale has been argued^{126,127} to be the generalization of the Ritchie's scale^{128,129} (eq 26) and is applicable to a larger domain

$$\log(k/k_0) = N_+ \quad (26)$$

where k_0 and N_+ are electrophile- and nucleophile-dependent parameters, respectively. Mayr's scale is used¹³⁰ to analyze the HSAB principle. It has been used by other researchers as well.^{96,131–133} Figure 5 shows¹³³ the linear correlation between E and ω for diazonium ions and their reactivity during interactions with π -nucleophiles.

The carbene-philicity scale (m_{CXY}) proposed by Moss^{134–139} on the basis of kinetic data has been shown to be useful in categorizing carbenes^{134–143} into electrophilic, ambiphilic, and nucleophilic varieties. There exists a linear free energy relationship between the carbene-philicity and the related Taft substituent parameters. These parameters have been made use of^{144,145} in analyzing the connection between skin sensitization and electrophilicity. A comparison between m_{CXY} and ω has been made recently.¹⁴⁶

The Swain–Scott free energy relationship¹⁴⁷ along with Legon's^{104,105} electrophilicity scale have been made use of in analyzing¹⁴⁸ the kinetics and mechanism of oxidation by halogens and interhalogens vis-à-vis their electrophilicity trends.

As discussed in the previous section, a theoretical electrophilicity index (ω) has been proposed by Parr et al.,¹⁴ prompted by a qualitative finding by Maynard et al.^{13,63,149} As electrophilicity and nucleophilicity are physically inverse of each other, the multiplicative inverse ($1/\omega$) (in the spirit of the definition of softness ($S = 1/\eta$)) and an additive inverse ($1 - \omega$) have been proposed¹⁵⁰ as possible definitions of nucleophilicity. Of course, C/ω and $(B - \omega)$, where C and B are constants (B may be zero also) for a given series of molecules, may be considered as well. Similar inverse behaviors have been proposed by others.^{104,105,151,152} The electrophilicity index (ω) obtained from several models for the charge dependence of the energy¹⁵³ within a broad framework of valence state atoms in molecules^{154–156} has been reviewed recently.¹⁵³ In an interesting study on the nucleophilic substitution reaction of carbonyl compounds, it has been shown¹⁵⁷ that the concerted reaction mechanism will be associated with a large electrophilicity/nucleophilicity gap whereas the corresponding small gap will imply a stepwise reaction mechanism.

Several electrophilicity scales based on different physicochemical properties have been proposed prior to the introduction of the electrophilicity index (ω) rigorously defined by Parr et al.¹⁴ In addition to the Ingold prescription,^{108,158} solution phase ionization potentials,¹⁵⁹ ^{13}C NMR chemical shifts,^{160–162} IR absorption frequencies,^{160,161} charge decompositions,¹⁶³ LUMO energies,^{164–167} ionization potentials,¹⁶⁸ redox potentials,¹⁶⁹ scaled electrophilicity parameter,¹⁶⁹ HPLC,¹⁷⁰ solid-state syntheses,¹⁷¹ K_e values,¹⁷² isoelectrophilic windows,¹⁷³ and the harmonic oscillator models of the aromaticity (HOMA) index¹⁷⁴ are some of the related quantities and subjects that have been used to understand the electrophilic/nucleophilic characteristics of chemical systems. Other related reactivity descriptors include an empirical nucleophilicity index in terms of the gas-phase ionization potential,^{175–180} nucleofugality,¹⁸¹ leaving group quality,¹⁸² potential philicity (phobicity),¹⁸³ charge philicity (phobicity),¹⁸⁴ etc. A net electrophilicity ($\Delta\omega^\pm$) (electroaccepting power (ω^+) relative to electrodonating power (ω^-))¹⁸⁵ has also been introduced.

The various electrophilicity descriptors reported in this section may be broadly classified into three categories: viz., the kinetic descriptor that measures the rate at which an electrophilic attack takes place, the thermodynamic descriptor which measures the ease of such an attack, and a combination of these two approaches through a linear free energy relationship. Among the kinetic scales, the most important is that of Mayr and co-workers.^{111–125} They have rank ordered various nucleophiles and electrophiles in terms of their N and E parameters respectively obtained from the associated experimental rate constants. Ritchie's parameters^{128,129} or Swain–Scott parameters¹⁴⁷ are similar in spirit. Various quantum chemical and spectroscopic quantities such as LUMO energy,^{164–167} ^{13}C NMR chemical shift, ν_{CO} frequencies,^{160–162} and charge decomposition¹⁶³ have been correlated with the respective rates of the reaction. Thermodynamic electrophilic descriptors are based on the strength of the bond formed between the electrophilic and nucleophilic sites. They include the HOMA index as a measure of aromaticity¹⁷⁴ or other aromaticity indices,²⁷¹ bond force constants,^{104,105} covalent bonding interaction via an HPLC assay,¹⁷⁰ solid-state synthesis,¹⁷¹ K_e parameters,¹⁷² LUMO energies,^{164,172} hydrophobicities,¹⁷³ redox potentials,¹⁶⁹ maximum acceptor superdelocalizabilities,¹⁶⁴ and solution phase ionization po-

tentials.¹⁵⁹ Uses of Hammett or Taft parameters in various linear free energy relationships have also been reported.^{13,160,376,377} A linear correlation between the ionization and activation energies for electrophilic additions is also known.¹⁶⁸ Most of these descriptors are empirical in nature, and it is not always possible to transfer the parameters for one class of compounds to the other. The Maynard–Parr electrophilicity index is based on a firm footing. Originally, it was introduced by Maynard et al.¹³ as $\omega = \chi^2/2\eta$, when they noticed that, compared to other descriptors, this quantity had a stronger correlation with the logarithm of the experimental rates of the reactions between HIV-1 nucleocapsid p7 zinc finger thiolates and different electrophilic ligands. It has been shown by Parr et al.¹⁴ that the energy change associated with the process of an atom or a molecule in the gas phase becoming saturated with electrons from the environment is given by this quantity and, hence, can be legitimately considered to be a definition for electrophilicity. It may be noted that the basis of the definition given by Maynard et al. is kinetic in nature whereas the interpretation by Parr et al. is thermodynamic in nature.

4.2. Local Approach

The majority of the empirical electrophilicity scales introduced so far are global in nature. Their local variants were developed only recently, and most of them are based on the Fukui function.^{71–75,186} An elegant recent review is obtainable in ref 20. Applications of these descriptors in understanding the substituent effects on the electrophilic processes^{187–189} and related studies^{190–193} highlight the power of these indices. Radical charge-transfer Fukui functions have been used¹⁹⁴ within an atoms-in-molecules (AIM) framework¹⁹⁵ for recognizing the electrophilic and nucleophilic centers in a molecule.

Relative electrophilicity and relative nucleophilicity are defined as s_k^+/s_k^- and s_k^-/s_k^+ , respectively.¹⁹⁶ Although they perform better^{106,107,196–198} than the Fukui function or local softness on certain occasions, they suffer from various drawbacks.^{96,186,191,192,199} In light of the local HSAB principle,^{94,95} a softness matching index has been defined^{200–202} for analyzing the regioselectivity as

$$\Delta_{ij}^{kl} = (s_i^- - s_k^+)^2 + (s_j^- - s_l^+)^2 \quad (27)$$

when atoms i and j of a nucleophile form a cycloadduct through the atoms k and l of an electrophile. The corresponding philicity⁹⁸ related quantity was also reported recently.²⁰³

Novel reactivity and selectivity indices have been proposed as the integral²⁰⁴ between the electrophilic Fukui function on one reactant and the nucleophilic Fukui function on the other or a difference²⁰⁵ between two such functions. A similar integral index appears in quantum similarity studies also.²⁰⁶ The effect of excess nucleophilicity over electrophilicity or vice versa (in a group sense) has been analyzed in the context of all-metal aromaticity/antiaromaticity and a possible molecular electronics.²⁰⁵

Local softness^{19,93} and local electrophilicity^{96–98} also perform well in analyzing regioselectivity. Figures 6 and 7 depict the beautiful correlations⁹⁶ between the experimental electrophilicity and respectively the global and local electrophilicities of a series of benzhydryl cations.

Li and Evans^{207,208a} have restated the local HSAB principle^{94,95} as, “For the soft–soft reactions the site with




Figure 6. Correlation between the experimental electrophilicity (E) and the theoretical relative electrophilicity ($\Delta\omega$) of a series of benzhydryl cations. Reprinted with permission from ref 96. Copyright 2002 American Chemical Society.




Figure 7. Correlation between the experimental electrophilicity (E) and the theoretical relative local electrophilicity ($\Delta\omega_c$) of a series of benzhydryl cations. Reprinted with permission from ref 96. Copyright 2002 American Chemical Society.

the maximum value of the Fukui function is preferred and the preferred site for the hard–hard interactions is that with the corresponding minimum value”. The ambiguity in local hardness is felt in the local version of the HSAB principle.^{94,95a} Minimization of a local grand potential is ad hoc.^{95b} Even the minimization of energies of individual atoms does not necessarily lead to the minimization of the overall energy of the molecule. While the Gázquez–Méndez^{94,95a} version considers similar Fukui function (FF) values for preferred interactions, the Li–Evans²⁰⁷ principle dictates the preference of soft–soft interactions in the maximum FF site whereas the minimum FF site is the best for the hard–hard interactions. However, it has been argued^{99,100} that the site with the minimum FF is the least reactive and, in the absence of a proper local hardness, hard reactions are to be analyzed in a different fashion at the local level. Of course the minimum Fukui function condition may be used as a “tiebreaker” in some specific strongly electrostatically controlled reactions.^{208b–d} Since hard interactions are electrostatic in nature, the Fukui function has been argued to be a poor descriptor for these reactions.⁹⁹ Charges^{99–101,209} or associated quantities^{70,210} such as molecular electrostatic potentials and local hardnesses are supposed to provide a better description for hard reactions.




Figure 8. Variation of μ , η , γ , ω , and $\partial\omega/\partial N$ for neutral atoms from He to Kr. Reprinted with permission from ref 227. Copyright 2003 American Chemical Society.

Other quantities used for this purpose include the 1s electron energy of nitrogen in the substituted aniline,²¹¹ the average local ionization energy,^{212,213} pair-site nonlocal interactions,²¹⁴ electron localization functions,^{215,216} etc. Nuclear Fukui functions^{217–219} and electrofugality²²⁰ have also been introduced. Local nucleophilicity^{221,222} and local hypersoftness²²³ have been used recently. Local polarizability is also treated as a descriptor to rationalize the enantiomeric excess in certain asymmetric hydrogenation reactions.²²⁴

5. Intramolecular and Intermolecular Reactivity Patterns

5.1. Periodicity

The concepts of atomic shell structure and chemical periodicity as proposed by Mendeleev form the cornerstone of chemical education. A periodic law may be stated as:²²⁵ “The properties of chemical elements and their compounds are periodic functions of the atomic numbers of the elements”. Atoms with completely filled shells and subshells are often relatively more stable and less reactive when compared with their open-shell counterparts. As expected from the principles of maximum hardness (MHP)^{44–51} and minimum polarizability (MPP),^{56–59} hardness increases along a period and decreases along a group whereas polarizability decreases along a period and increases along a group.^{51,226,227} Alkali metals are the softest and the most polarizable while noble gases are the hardest and the least polarizable.^{226,227} Electrophilicity also exhibits characteristic periodic oscillations with maxima on halogens, which are most electronegative and least nucleophilic as well.^{226,227} As shown in Figure 8, the variation of $[\partial\omega/\partial N]$ for neutral atoms mimics²²⁷ that of μ because of the small values²²⁸ of $\gamma = 1/3[\partial\eta/\partial N]_{v(\vec{r})}$.

5.2. Excited States

Through the excited-state DFT calculations on atoms and molecules in various electronic states which happen to be the lowest state of given symmetry^{229–231} or in ensembles of states^{232–235} (along with the related penalty-function-based formalisms²³⁶), it has been shown^{237–240} that, “A system is harder and less polarizable in its ground state than in any of its excited states and an increase in the excited state contribution in a two state ensemble makes the system softer




Figure 9. Plot of electrophilicity index (ω) versus cutoff radius (R_c) for atoms (He, Li, Be, B, C, N, O, F, Ne) confined in a spherical box. Reprinted with permission from ref 250. Copyright 2003 American Chemical Society.

and more polarizable”. This fact is in conformity with MHP and MPP since an atom or a molecule is generally more reactive in its excited state. For example, the S (α) values (in au) of He atom in different electronic states are as follows:²³⁸ (1S , 1.51 (1.86); 1P , 6.89 (117.92); 1D , 13.09 (728.71); 1F , 21.95 (3536.36)). And the $\alpha(\eta)$ values (in eV) of HF molecule are as follows:²⁴⁰ ($\sigma^2\pi^4$, $^1\Sigma^+$, 5.86 (10.8); $\sigma^2\pi^3\sigma^1$, $^3\Pi$, 38.4 (4.97); $\sigma^2\pi^3\sigma^1$, $^1\Pi$, 39.5 (3.91)). Even in time dependent situations^{43,241–245} involving excited states including Rydberg states,^{246–248} this fact remains valid.

Any system is generally less electronegative in its excited state.¹⁵⁰ The behavior of electrophilicity in the excited state in comparison to that in the ground state, however, will depend completely on the relative variations in χ and η upon electronic excitation, although both of them often decrease. **Electrophilicity is generally less in the excited state than in the ground state, as expected from a minimum electrophilicity principle.**²⁴⁸ For example, the $\chi(\eta, \omega)$ values (in au) of He atom in different electronic states at the beginning of the process are as follows:¹⁵⁰ ($^1S(1s^2)$, 0.2591 (0.3920, 0.0856); $^1P(1s2p)$, 0.2044 (0.1315, 0.1589)).

5.3. Spherical Confinement

The concept of confined quantum mechanical systems stems from the idea of simulating the effect of pressure on atoms or molecules by confining them in impenetrable spherical boxes.²⁴⁹ Numerical Hartree–Fock calculations with Dirichlet boundary conditions of various global reactivity descriptors of several atoms and ions have revealed^{250–252} that all systems become harder and less polarizable with an increase in pressure. The inverse relationship $\eta \propto 1/\alpha^{1/3}$ remains valid when the confinement volume is decreased. Electrophilicity is not very sensitive²⁵⁰ (Figures 9 and 10) to confinement except for very small cutoff radii, where it increases abruptly. Variations of ω with the atomic number (section 5.1) and degree of ionization (more positive charge implies more electrophilic) remain unaltered for all possible extents of confinement.




Figure 10. Plot of electrophilicity index (ω) versus cutoff radius (R_c) for ions (C^{n+} ; $n = 1, 2, 3, 4$) confined in a spherical box. Reprinted with permission from ref 250. Copyright 2003 American Chemical Society.

5.4. Chemical Processes

Both global and local electrophilicities have been found to be helpful in analyzing the reactivity and selectivity behavior of various chemical compounds as well as the reaction mechanisms of diverse classes of chemical processes. A plethora of systems have been studied including pentazolato complexes of the first row transition metals,²⁵³ diazonium ions,¹³³ carbonyl carbons,²⁵⁴ fluorine substituted disilanes,²⁵⁵ carbenes,^{146,256} Fischer-type chromium–carbene complexes,²⁵⁷ copper clusters,²⁵⁸ zeolites,²⁵⁹ group-14 elements and related functional groups,⁶⁴ aliphatic amines,²⁶⁰ alkanes,²⁶¹ silylenes and germylenes,¹⁵² cobalt porphyrins and related aza derivatives,²⁶² highly hindered polyanionic chelating ligands,^{263,264} organorhenium²⁶⁵ and organoneodymium complexes,²⁶⁶ and thiadiazolium salts.²⁰³ Figure 11 clearly reveals¹⁵² the power of global and local electrophilicities through beautiful linear variation of the reaction energy with these quantities associated with the complexation reactions of silylenes and germylenes with ammonia.




Figure 11. Plots of global (ω) and local (ω_{Si} and ω_{Ge}) electrophilicities of silylenes and germylenes versus their reaction energies (ΔE) with a Lewis base such as NH_3 . Reprinted with permission from ref 152. Copyright 2005 American Chemical Society.

Intermolecular reactivity of carbonyl compounds has also been studied²⁶⁷ using the group philicity. The importance of a theoretical analysis of the philicity/electrophilicity behavior in providing an effective synthetic protocol has been highlighted.^{263,264} The connection between the electrophilicity^{141,142} index and aromaticity^{268–273} as well as superacidity^{274–279} (superelectrophilicity) has been analyzed. Global electrophilicity values need not always correlate²⁷³ with the nucleus independent chemical shift values,²⁸⁰ which characterize the magnetic aspects of aromaticity. **Stability and reactivity of cyclic alkali clusters like Na_6^{281} and K_6^{281} and cyclic trigonal dianion metal clusters like Be_3^{2-} , Mg_3^{2-} , Zn_3^{2-} , and their different types of sandwich complexes,^{282–284} stabilization of direct $Zn-Zn$,^{282,285} $Be-Be$,²⁸⁶ and $Mg-Mg$ ²⁸⁷ bonds by Be_3^{2-} and Mg_3^{2-} rings, multidecker sandwich complexes,²⁸⁸ etc. have been successfully analyzed by electrophilicity and its local variants. Minimum electrophilicity principle has been shown to be operative in the cases of the all-metal aromatic systems²⁷³ as well as in the magic clusters.²⁸⁹**

The stability and reactivity of azametallocenes have been studied²⁵³ in terms of their global electrophilicity (ω) values, which corroborate the fact that the charge transfer from the ligand to the metal makes the aromatic pentazolato ligand more electrophilic. Similar studies on aza derivatives of cobalt porphyrin have revealed²⁶² that an increase in the number of aza-N atoms renders an electrophilicity enhancement at the cobalt center of the cobalt porphyrin complex. The presence of electron-withdrawing substituent groups on the carbonyl carbon atom in phenyl acetates has been shown²⁵⁴ to drive the nucleophilic attack at those carbon centers through the enhancement of electrophilicity. Studies on Fischer-type chromium–carbene complexes have highlighted²⁵⁷ that their electrophilicity is reduced due to the presence of π -donor substituents because the acceptor orbital in carbene gets occupied by π -donation. Similar behavior has been reported for silylenes and germylenes as well.¹⁵² Among the pyridyl substituted bis-coumarins, the para-compound is the most electrophilic whereas the ortho-isomer is the least electrophilic.²⁶⁶ In the corresponding neodymium complexes, the carbonyl oxygen is the most favorable site for the electrophilic attack in comparison to the hydroxyl oxygen, the lactone oxygen, or the nitrogen atoms.²⁶⁶ However, for metal coordination in the double-deprotonated compound, both carbonyl and hydroxyl oxygen sites become favorable sites for electrophilic attack.²⁶⁶ Similar analysis on organorhenium complexes²⁶⁵ shows that 2,2'-azobis(5-chloropyrimidine) ligand (L) is a better π -acceptor than the 2,2'-azobispyridine ligand (Cl) in the dinuclear radical anion complexes: $\{(\mu-L)[Re(CO)_3Cl]_2\}$. Electrophilicity remains more or less constant for most of the fluorine substituted disilanes.²⁵⁵ Neutral copper clusters, on the other hand, exhibit odd–even oscillations,²⁵⁸ as evidenced by experiments and theoretical calculations. Odd clusters are more electrophilic, are softer, and have the capacity to attain a closed-shell configuration by accepting electrons. Adsorption of small molecules and cracking of hydrocarbons in zeolites are properly accounted for by the philicity.²⁵⁸ Theoretical calculation of ω using spin–orbit interactions for halogens and group 14 atoms can reproduce the experimental trend of a monotonic decrease in electrophilicity by going down the group.⁶⁴ Global and local electrophilicities can properly reproduce the experimental electrophilicity/nucleophilicity patterns of diazonium ions,¹⁵³ aliphatic amines,²⁶⁰ carbonyl compounds,^{254,260} thiadiazolium salts,²⁰³

etc. This is also true in most of the other cases described above. As is demonstrated in the case of highly hindered polyanionic chelating ligands, sometimes the theoretical values even suggest^{263,264} possible efficient synthetic protocols.

Electrophilicity index and its variants have been found to be useful in understanding spectral shifts in acetone,²⁹⁰ excited-state properties of organic dyes,²⁹¹ α -pinene isomerization reaction,²⁹² application of 8-hydroxyquinoline metal complexes in organic light-emitting diodes,^{293,294} internal rotation barrier in alkanes,²⁹⁵ acceptor properties of quinone-derivatized polypyridinic ligands,²⁹⁶ reactivity of exohedral fullerenes $C_{64}X_4$,²⁹⁷ cyclopolymerization reactions of diallyl monomers,²⁹⁸ properties of pyridine-derived N-heterocyclic carbenes,²⁹⁹ hydrolytic deamination of a cytosine derivative in a protic medium,³⁰⁰ reactivity of pyridine derived N-heterocyclic germynes,³⁰¹ radical stability in terms of bond dissociation enthalpies,³⁰² chemical reactivity of pyrrole derivatives,³⁰³ chemical explosivity,³⁰⁴ thermal stability of nitroaromatic compounds,³⁰⁵ properties of phenol compounds,³⁰⁶ nucleophilic behavior of benzofused thieno [3,2-b] furans,³⁰⁷ hydride affinity of quinines,³⁰⁸ stability of N-heterocyclic silylenes,³⁰⁹ reactivity of coinage metal clusters,³¹⁰ modulation of N_2 activation in triamidoamine—Mo complexes,³¹¹ reactivity of nickel-modified platinum cluster,³¹² active site in bis(imino) pyridyl iron catalyst,³¹³ etc. Several other popular electronic concepts like electronegativity,³¹⁴ charge transfer descriptor,³¹⁵ electron-acceptor characteristics,³¹⁶ reactivity of paracyclophanes,³¹⁷ etc. have been connected with electrophilicity.

The main classes of reactions for which the mechanisms have been analyzed in terms of global and local electrophilicities comprise 1,3-dipolar cycloaddition reactions,^{318–324} specific cycloaddition reactions such as Diels–Alder reactions^{199,325–332} with varied diene–dienophile pairs, and other cycloaddition reactions.^{333–338} A typical Diels–Alder reaction will follow a nonpolar pericyclic mechanism if the electrophilicity difference between the diene and the dienophile is small whereas a polar mechanism will be followed for a large value of this difference. **Philicity also properly accounts for these reactions.**³³⁹

Cycloaddition reactions constitute the most widely studied pericyclic processes where two π -components approach to form two new σ -bonds within a cyclic framework. In the Diels–Alder-type cycloaddition reactions, a diene and a dienophile interact to form a six-membered ring product. Considerable enhancement in the rates of these reactions has been noticed in the diene–dienophile pair with an electron-withdrawing substituent in one and an electron-releasing substituent in the other. It is expected that the difference in their global electrophilicity values will provide important insights into the associated reaction mechanism. Pericyclic processes with a nonpolar mechanism are characterized by a small electrophilicity difference of diene and dienophile whereas a large difference leads to a polar mechanism in the corresponding ionic processes. Use of the Maynard–Parr electrophilicity index and its local variant in understanding these aspects has been attempted mainly by Domingo and co-workers. In the normal electron demand reactions, the presence of electron-withdrawing groups in the dienophile increases the reaction rates of the associated Diels–Alder reactions. In these reactions, the charge transfer is from the diene–nucleophile to the dienophile–electrophile. However, in the inverse electron demand-type reactions,

an electron-withdrawing substituent is present in the diene and/or an electron-releasing substituent is present in the dienophile, so that the charge gets transferred from the dienophile to the diene. Rank ordering of various dienes and dienophiles as electrophiles or nucleophiles has been done so that the reaction mechanism for a given pair may be ascertained at the beginning.³³² Cycloaddition reactions with large ionic character involving large electrophilicity differences include the reactions between 2-methylfuran and a masked *o*-benzoquinone,³³⁴ substituted butadienes and ethylenes,¹⁹⁹ Lewis acid coordinated 2-(trimethylsilyloxy)-acrolein and furan,³³⁵ etc., whereas those with a nonpolar mechanism include the concerted [4 + 2] process between 2-azadiene and cyclopentene/propene,³²⁹ the concerted [3 + 2] process between benzonitrile oxide and ethynyl/propynylboronate,³²² etc. This analysis allows one to devise a strategy so that the changes in the nature of the substituents in the dienes/dienophiles or changes of the reaction conditions, including the presence of a Lewis acid catalyst or a polar solvent, may change a nonpolar concerted process to a polar stepwise process. This causes an enhancement in the rate of the reaction and, in turn, of the yield of the corresponding kinetically controlled products. Several related experimental trends are understood through the electrophilicity analysis. They include Lewis acid catalyzed [4 + 2] and [4 + 3] cycloadditions between cyclopentadiene and arylidenoxazolones,³³⁷ 1,3-butadienes and dimethyl acetylenedicarboxylate,³³⁸ *N*-acyl-1-aza-1,3-butadienes and vinylamines,³³⁰ nitroalkenes and aluminum derivatives of vinyl ethers,³³¹ butadiene derivatives and acetone,³²⁷ and cyclopentadiene and cyanoethylenes.³²⁶ Problems associated with this analysis in explaining inverse electron demand Diels–Alder reactions have also been reported.³²⁸ It has been argued³²⁸ that they are due to the inadequacy of the frontier molecular orbital theory.

Similar electrophilicity-based analysis has been extended to a variety of 1,3-dipolar cycloaddition reactions with a variety of dipole/dipolarophile pairs. Larger electrophilicity differences correspond to faster reactions.³²¹ Polar regioselective reactions between nitrile/azomethine ylide and methyl acrylate/propiolate have been properly accounted for by the relative global and local electrophilicities between dipoles and dipolarophiles.³²⁴ The regio- and stereoselectivity of various 1,3-dipolar cycloaddition reactions, such as that between hindered thiocarbonyl ylides and tetracyanoethylene,³²³ nitrile oxide and anthracene/acridine,³³⁶ *C*-(methoxy carbonyl)-*N*-methyl nitron and methyl acrylate/vinyl acetate,³²⁰ 5-ethoxy-3-*p*-(*S*)-tolylsulfanyl furan-2(5H)-ones and diazoalkanes,³¹⁹ etc., have been shown to be in conformity with those predicted by the relative electrophilicity patterns. This approach is found to be more reliable³³⁶ than the frontier molecular orbital theory.

Polar cycloaddition reactions between electron-rich dienes and electron-deficient heteroarenes are also properly analyzed using electrophilicity.³⁴⁰ Apart from highlighting the importance of the superelectrophilicity concept as well as that of the σ -complexation, a connection between the S_NAr reactivity and the feasibility of the related σ -complexation process is properly delineated.

Rank ordering of various dienes–dienophiles/dipoles–dipolarophiles may be summarized as in the following scheme.^{321,332}

Large ω	Small ω
Strong electrophile (weak nucleophile): <i>electron-poor dienophile (e.g. acrolein) and dipolarophile (e.g. nitroethylene) in a normal electron-demand (NED) reaction</i>	Weak electrophile (strong nucleophile): <i>electron-rich diene (e.g. 1-methoxy-1,3-butadiene) and dipole (e.g. azides) in an NED reaction</i>
<i>For inverse-electron-demand (IED) reactions: Local philicity (ω_i) or a group philicity would be a better descriptor to identify the electron releasing (withdrawing) substituent on the dienophile (diene)/ dipolarophile (dipole).</i>	

The following scheme provides a guideline toward the possible reaction mechanism associated with a given Diels–Alder/1,3-dipolar cycloaddition reaction.^{102,321,332}

$\Delta\omega$ (diene/ dienophile or dipole/ dipolarophile interacting pair)	
Small	Large
<i>Non polar, pericyclic, concerted process, frontier-controlled, soft-soft interactions e.g. Ethylene/1,3-butadiene reaction</i>	<i>Polar, ionic, stepwise process, charge-controlled, hard-hard interactions e.g. N-methylmethylen ammonium cation/1,3-butadiene reaction</i>
<i>For Lewis-acid catalyzed reactions: Increased ω of the electron-poor component: dienophile (NED reaction)/ diene (IED reaction)</i>	

Apart from these major types of reactions, the electrophilicity concept at both the global and the local levels has been used to analyze a wide variety of chemical processes. They include oxidation of thiophenes,³⁴¹ catalytic olefination of carbonyl compounds³⁴¹ and polyhaloalkanes,³⁴² reduction of formylchromones,³⁴³ hydride transfer reaction in 1-methyl nicotinamide-lumiflavine,³⁴⁴ formaldehyde decomposition,^{345a} intermolecular ligand exchange in alkyltin trihalides,^{345b} dissociation of HS^- (HCN),^{345c} discoloration of titanyl-porphyrin,^{345d} nucleophilic addition to carbon–carbon double bonds,³⁴⁶ Friedel–Crafts benzylation and acylation reactions,³⁴⁷ metalation of oxazolines,³⁴⁸ oxidation of thiols,³⁴⁹ alkaline hydrolysis of *N*-phenylacetamides,³⁵⁰ ene reactions of nitroso compounds,³⁵¹ aminolysis of thiocarbonates,³⁵² etc. The main theme of these studies is to identify a reactant which will act as an electrophile (larger ω value) or another to behave as a nucleophile (smaller ω value). Apart from the thermodynamic information content of ω as shown in Figure 11, it has been shown to possess enough kinetic information as well.^{346,347,352}

Global ω values of nitroso compounds suggest that during their reactions, such as ene reactions, they behave³⁵¹ as good nucleophiles similar to singlet oxygen and triazolidiones. Their electrophilicity stems from the fact that their HOMOs are formed through high energy antibonding combination of lone pairs of N and O centers and are orthogonal to their low energy π^* LUMO. The energy change associated with the rate-determining step of the alkaline hydrolysis of *N*-phenylacetamides has been shown to decrease with an increase in their electrophilicity values.³⁵⁰ An electrophilicity analysis of oxidation of thiols by cobalt N_4^- complexes reveals that azaporphyrins are very good electron acceptors during the oxidation process, which gets improved in the presence of polar solvents.³⁴⁹ The behavior of chloromethyl derivatives toward metalation can be rationalized.³⁴⁸ Quantitative linear relationships between the experimental rate

constants and the Maynard–Parr electrophilicity index and/or its local counterpart have been obtained for the Friedel–Crafts reactions,³⁴⁷ nucleophilic addition involving the $\text{C}=\text{C}$ double bond,³⁴⁶ and aminolysis of thiocarbonates and dithiocarbonates.³⁵² The reactivity patterns associated with formaldehyde decomposition,³⁴⁵ intermolecular ligand exchange in alkyltin trihalides,³⁴⁵ hydride ion transfer,³⁴⁴ etc. have been properly analyzed in terms of the electrophilicity index or its variants. **The electrophilicity and nucleophilicity have been considered to be two extremes of the same scale,³⁵³ but there is no such strong theoretical footing for nucleophilicity.^{354,355} The electrofugality³⁵⁵ may also be identified with the negative³⁵⁴ of the electrophilicity.**

5.5. Solvent Effects

To understand the effect of a solvent on electrophilicity, a variation of ω (eq 12) up to first order has been written by Pérez et al.³⁵⁶ as

$$\Delta\omega(1 \rightarrow \epsilon) = (\mu/\eta)\Delta\mu - \frac{1}{2}(\mu/\eta)^2\Delta\eta = \Delta\omega^{(1)} + \Delta\omega^{(2)} \quad (28)$$

where ϵ is the dielectric constant of the medium and $\Delta\mu$ and $\Delta\eta$, respectively, describe the variation in μ and η when the system goes from the gas phase to the solution.

The insertion energy, ΔE_{ins} , of the solute going to solvent is defined as twice^{357–359} the energy of solvation, ΔE_{solv} , i.e.,

$$\Delta\omega^{(1)}(1 \rightarrow \epsilon) = \left(\frac{\Delta E}{\Delta N}\right)_{v(\vec{r})} \left(\frac{\Delta N}{\Delta \mu}\right)_{v(\vec{r})} \Delta\mu \approx \Delta E_{\text{ins}} = E(\epsilon) - E(1) = 2\Delta E_{\text{solv}} \quad (29)$$

The second term of eq 28 becomes

$$\Delta\omega^{(2)}(1 \rightarrow \epsilon) = \frac{\mu}{\eta\Delta N} E_{\text{solv}} \quad (30)$$

A linear relation³⁵⁶ between $\Delta\omega(1 \rightarrow \epsilon)$ and ΔE_{solv} for a series of both neutral and charged electrophilic ligands as shown in Figure 12 highlights the authenticity of this approach.

A somewhat similar and related analysis has been presented in refs 360 and 361. In these papers, the effects of solvent on two intramolecular rearrangement reactions, viz., $\text{trans-N}_2\text{H}_2 \rightarrow \text{cis-N}_2\text{H}_2$ and $\text{F}_2\text{S}_2 \rightarrow \text{FSSF}$, have been studied.^{360,361} It has been demonstrated that solvation makes the reactions more favorable both thermodynamically and kinetically. As shown in Figure 13 for the N_2H_2 rearrangement, ω passes through³⁶¹ an extremum at the transition state (TS) for both the reactions and both the phases although the dipole moment does not always pass through an extremum at the TS. In both the cases, the solvation decreases the electrophilicity.

The reactivity of several systems has been studied to gain insights into the solvation effects on both global and local electrophilicities. Charged peroxides,³⁶² cyclopropane ring-opening in duocarmycin SA derivatives,³⁶³ various electron donors,¹⁵⁹ aliphatic amines,²⁶⁰ different organometallic compounds,³⁶⁴ carbonyl compounds,³⁶⁵ dye–redox mediator reactions,³⁶⁶ etc. have been studied for this purpose. In general, the reactions become easier to perform in the solution phase, with some exceptions.³⁶⁷ The effects of the




Figure 12. Plots of electrophilicity changes versus solvation energy of a series of both charged and neutral electrophilic ligands. Reprinted with permission from ref 356. Copyright 2001 American Chemical Society.

solvent are more pronounced for the global electrophilicity than for its local variant.

5.6. External Field Effects

The reactivity of a chemical system changes drastically in the presence of an external field. This field may be an explicit external electromagnetic field or it may arise due to the presence of another molecule (reagent/reactant) and/or a solvent. The nuclear Coulomb potential in the parent molecule being experienced by its electrons would get augmented by the potentials generated due to the electrons and nuclei present in the reactant/reagent/solvent molecules in addition to the possible electron transfer between them. All these effects, including any external electromagnetic field, can be simulated by the presence of an external generic field. An explicit dynamical study on the variation of chemical reactivity in the presence of an external electric field will be discussed in section 7.

Changes in molecular reactivity and selectivity due to the presence of an applied uniform electric field are analyzed.³⁶⁸ The internal electrostatic field of the molecule plays a crucial role in determining the chemical reactivity when the strength of the external field is low. However, at larger field strengths, significant variation in reactivity has been observed.³⁶⁸

Since any quantum system can be completely characterized by its number of electrons (N) and the external potential ($v(\vec{r})$), variation in any property of that system, including its chemical reactivity, may be analyzed by changing N and $v(\vec{r})$. For example, the change in chemical potential may be written as¹⁹

$$d\mu = \eta dN + \int f(\vec{r}) dv(\vec{r}) d\vec{r} \quad (31)$$

For a uniform electric field $E(\vec{r})$ it may be written as

$$d\mu = \eta dN - \int f(\vec{r}) E(\vec{r}') d\vec{r} d\vec{r}' \quad (32)$$

Therefore, the changes in chemical potential would be directly proportional to the strength of the external field when there is no charge transfer. It has been shown³⁶⁸ that μ and ω get significantly altered when the external field strength is increased. The effect of the external field is only marginal in the case of η , which implies that the second-order variation in the energy due to external perturbation is less than the corresponding first-order variation. A dynamical variant of this will be analyzed in section 7.

Local reactivity indices such as the Fukui function and the philicity have been shown³⁶⁸ to change drastically in the presence of the external field. It may be noted that the variation of f_k^α and ω_k^α are not similar because the global electrophilicity also changes in this case. All these changes become more pronounced as the number of atoms in the molecule and the external field strength increase.

The effects of the external potential variation on reactivity and regioselectivity have been analyzed within variational³⁶⁹ as well as perturbative^{370,371} frameworks. A cooperative effect of the solvent and surface together to increase the reactivity has also been reported.³⁷²

5.7. Biological Activity and Toxicity

There has been a recent upsurge of interest in unraveling the connection between electrophilicity and biological activity, especially toxicity, mutagenicity, and carcinogenicity in different chemical, biological, and biochemical systems,^{373,374} to broaden the applicability of the associated quantitative structure–activity relationships (QSAR). Most of these analyses are qualitative in nature with more interpretive power and relatively less predictive potential. Based on these newly acquired ideas, strategies for rational drug designs have been developed. Relationships have been shown between electrophilicity and allergic contact dermatitis,³⁷⁵ including skin sensitization,^{376,377} the activity of phase 2 enzyme and glutathione in protecting mammalian cells from malignancy,³⁷⁸ the toxicity of organic chemicals to *Tetrahymena pyriformis*¹⁶⁴ and *Chlorella vulgaris*,³⁷⁹ glutathione S-transferase induction by *tert*-butylhydroquinone,³⁸⁰ the mutagenicity and DNA damaging potential of *N*-acyloxy-*N*-alkoxymides,^{381a} DNA alkylation of duocarmycins,^{381b} the antioxidant activity of quercetin^{382a} and catechi,^{382b} the antitumor and antibacterial activity of kinamycins and lomaiviticin A,^{383a} the antileukemic activity of phenol,^{383b} etc.

Similar studies have been performed in showing the importance of the electrophilicity concept in analyzing the role of methyldiene imidazolone as an electrophilic prosthetic group,³⁸⁴ aquatic toxicity,^{165,166} polymeric surfactants as glutathione transferase mimics,^{380,385} charged phenyl radical activity toward nucleic acid components,³⁸⁶ in vitro trypano-




Figure 13. Variation of the electrophilicity index and the dipole moment in a vacuum and solvent along the reaction path of the intramolecular rearrangement reaction: $trans\text{-N}_2\text{H}_2 \rightarrow cis\text{-N}_2\text{H}_2$. Reprinted with permission from ref 361. Copyright 2001 American Chemical Society.

cid activities of some typical heterocyclic quinones,³⁸⁷ the importance of the standard-helix motif in biological hydrolysis,³⁸⁸ the function of the Michaelis complex of pI258 arsenate reductase,³⁸⁹ the behavior of zinc fingers related to retroviral activity,¹⁶⁹ suppression of breast cancer,³⁹⁰ HIV-1 nucleocapsid protein p7,^{63,149} *Escherichia coli* Ada protein,¹⁴⁹ etc.

The toxicity of polychlorinated biphenyls has been studied^{391,392} through the profiles of electrophilicity and philicity in both gas and solution phases. Figure 14 depicts the variation of these quantities for 2,2',5,5'-tetrachlorobiphenyl as a function of the torsional angle. High rotational energy barriers do not allow the toxin to rotate freely in a real environment so that it can interact with the cellular component of a living system.^{391–394} Therefore, toxicity is related to the low rotational barrier. A comparison between the rotational energy profile and those of the hardness and the polarizability clearly delineates^{391,392} that the high toxicity of PCBs is related to the minimum η value and the maximum α value, as expected from the MHP and the MPP. The

electrophilicity is often maximum in those conformations. The most active toxic sites are identified through the philicity profiles.

The biological activities of various testosterone derivatives in terms of relative binding affinity (RBA), androgenic potency, relative androgenic activity, therapeutic index, TeBG affinity, relative competition indices, binding affinity for rat ventral prostate receptor protein, and myotrophic to androgenic potency in temporal as well as some estrogen derivatives quantified in terms of their RBA values have been shown³⁹⁵ to correlate strongly with the electrophilicity index, suggesting it to be a suitable descriptor of the biological activity of these systems.

The toxicity of polychlorinated dibenzofurans (PCDFs) and dibenzo-*p*-dioxins (PCDDs) has been correlated with the electrophilicity index. The correlation is reasonably good.³⁹⁶ However, a combination of electrophilicity and philicity drastically improves the situation,³⁹⁷ as is authenticated by the analysis of the toxicity of various electron-donor- and electron-acceptor-type toxins, measured by their pIGC₅₀ data, toward *Tetrahymena pyriformis*. It also highlights the




Figure 14. Variation of relative energy, electronegativity, global electrophilicity index, and local electrophilic power of 2,2',5,5'-tetrachlorobiphenyl with the torsional angle. Reprinted with permission from ref 391. Copyright 2003 American Chemical Society.

importance of charge transfer between a toxin and a biosystem for an overall understanding of toxicity. Experimental toxicity values (pIC_{50}) of a variety of polyaromatic hydrocarbons³⁹⁸ such as PCDFs, PCDDs, and polychlorinated biphenyls (PCBs) as well as those of several aliphatic amines³⁹⁷ correlate well with the corresponding toxicity values calculated using the HF energy along with the global and local electrophilicities.

Electrophilicity index has also been made use of in analyzing antitumoral activity of synthetic chalcones,³⁹⁹ cis-platinum complexes,^{400,401} testosterone derivatives,⁴⁰² synaptosomal toxicity of 4-hydroxyl-2-nonenal,⁴⁰³ toxicity of PCDDs,⁴⁰⁴ hemolytic activity of methacrylates,⁴⁰⁵ activation of AhR by some biphenyls,⁴⁰⁶ metal-binding specificity of porphyrins,⁴⁰⁷ effect of plant growth regulators on chloro-acetanilide herbicides,⁴⁰⁸ reactivity of porphyrin pincers,⁴⁰⁹ mutagenic activity of dinitrobenzo[a]pyrene isomers,⁴¹⁰ activities of nonbenzodiazepine compounds,⁴¹¹ radical-scavenging activity of flavones,⁴¹² cancer-cell membrane lysis properties of peptides like PNC- 27,⁴¹³ different sulfonamides as carbonic anhydrase inhibitors against isozyme, CA-II,⁴¹⁴ skin sensitization potential of alkenes,^{415,416} arsenic toxicity,⁴¹⁷ toxicity of electrophilic sugars,⁴¹⁸ pyridodindoles,⁴¹⁹ etc. Biological activities of *N*-phenyl benzamides

(antimicrobial agents)⁴²⁰ and nifurtimox analogues (antiparasitic agents)⁴²¹ are also correlated with electrophilicity and its local variants.

To avoid collinearity and overfitting, pIC_{50} values of several electron-acceptor toxins such as PCDFs and PCBs are correlated with only one parameter, ω , and for the donor toxins such as aliphatic amines and amino alcohols, the related pIGC_{50} values are correlated with the related maximum local nucleophilicity values to obtain beautiful correlations.^{397b} The regression model is more robust^{397b} for acceptors than for donors. For example, the regression equations for 171 acceptors comprising saturated alcohols, diols and halogenated alcohols, mono- and diesters, carboxylic and halogenated acids, aldehydes, and ketones and for 81 donors comprising unsaturated, α -acetylenic and amino alcohols and amines with toxicity toward *Tetrahymena pyriformis* are as follows:^{347c}

Acceptors:

$$\begin{aligned} \text{predicted}(\text{pIGC}_{50}) &= 1.000(0.020) \times \text{observed}(\text{pIGC}_{50}) - \\ &\quad 1.708 \times 10^{-11}(0.019) \\ R^2 &= 0.937, R_{\text{CV}}^2 = 0.936, \text{SD} = 0.241, N = 171 \end{aligned}$$

Donors:

$$\text{predicted}(\text{pIGC}_{50}) = 1.000(0.036) \times \text{observed}(\text{pIGC}_{50}) - 1.044 \times 10^{-10}(0.039)$$

$$R^2 = 0.904, R_{\text{CV}}^2 = 0.899, \text{SD} = 0.232, N = 81$$

The number of carbon (non-hydrogenic) atoms, N_{C} (N_{NH}), may be considered to be a crude alternative^{397c} to the *n*-octanol/water partition coefficient (corresponding $\log P$, also correlated^{397d} to $N_{\text{C}}/N_{\text{NH}}$) in developing QSAR/QSTR models which can be further improved^{395b,397c} by adding ω and/or its local counterpart (ω_k^α) as well as charges on the relevant atoms (q_k). Of course, $N_{\text{C}}/N_{\text{NH}}$ is a lot easier to obtain than $\log P$. Several recent publications highlight the importance of electrophilicity in understanding biological processes and obtaining QSAR/QSTR models.^{395,397a–k} Although $N_{\text{C}}/N_{\text{NH}}/\log P$ can be used for a broad spectrum of molecules, $\omega/\omega_k^\alpha/q_k$ depends on the electronic environment and hence should be applied to a group of congener molecules,^{397l} as was done^{397a} for nine acceptor groups and four donor groups separately. Various physicochemical properties like charge transfer capability,⁴²² boiling point of alcohol,⁴²³ enthalpy of vaporization of PCBs,⁴²³ glass transition temperature,⁴²⁴ electro-acceptor characteristics,⁴²⁵ electronegativity, etc. correlate reasonably with $N_{\text{C}}/N_{\text{NH}}$. Electrophilicity and its local variants improve the correlation when used as additional descriptors.

In this section, the electrophilicity patterns related to chemical periodicity, the excited state reactivity, confined systems, and various intermolecular and intramolecular processes, including solvent and external field effects and biological activity, have been reviewed. The Maynard–Parr electrophilicity index and its local variants not only support the trends expected from chemical intuition but also provide new directions in analyzing reaction mechanisms of a diverse class of chemical reactions.

6. Variation of the Electrophilicity Index during Physicochemical Processes Such as Vibrations, Internal Rotations, and Chemical Reactions

6.1. Molecular Vibrations

It is important to know how the chemical reactivity of a molecule changes when it undergoes vibration, internal rotation, or chemical reaction. To analyze specifically the behavior of the electrophilicity index (ω , eq 12) in this regard, one starts from a first-order derivative of it as follows^{227,426}

$$\frac{\partial \omega}{\partial \lambda} = \frac{\mu}{\eta} \left(\frac{\partial \mu}{\partial \lambda} \right) - \frac{1}{2} \left(\frac{\mu}{\eta} \right)^2 \left(\frac{\partial \eta}{\partial \lambda} \right) \quad (33)$$

where λ may be a bond length (stretching), bond angle (bending), dihedral angle (internal rotation), or reaction coordinate (chemical reaction).

The extremal behavior of ω results from that of μ and η . If both μ and η are extrema (also constants or having inflection points), ω will be an extremum, and at that point the following condition will be satisfied:

$$\frac{\partial \mu}{\partial \lambda} = \frac{\mu}{2\eta} \left[\frac{\partial \eta}{\partial \lambda} \right] \quad (34)$$

So the extremum of electrophilicity will occur when the slopes of the changes in μ and η are of opposite signs,




Figure 15. Profile of the maximum charge transferred during the internal rotation of the hydroxylic group of the enol form of guanine. Reprinted with permission from ref 427. Copyright 2003 American Chemical Society.

because $\mu < 0$ and $\eta > 0$, owing to the convexity in energy. Therefore, ω will be a minimum (maximum) when both μ and η are maxima (minima).^{227,361,426}

Various global reactivity descriptors of water, ammonia, and ethane have been calculated⁴²⁶ for their equilibrium geometries as well as the distorted geometries originating from displacements along normal coordinates of vibration. The extremal analysis made above has been found to be true. For H_2O and NH_3 , the minimum energy–maximum hardness–minimum electrophilicity criteria for the equilibrium geometry have been shown⁴²⁶ to be true when compared with corresponding quantities for displacements along all allowed normal modes. For C_2H_6 , however, it is not true for a few normal modes because μ does not possess the maximum value where η is maximum in those cases.

6.2. Molecular Internal Rotations

It has been observed⁴²⁷ during the internal rotation of the hydroxylic group of the enol form of guanine that the stable conformations are associated with minimum energy, maximum hardness, and minimum electrophilicity values, as would have been expected from the above analysis. The corresponding transition states have been shown to be the most electrophilic. This fact is confirmed⁴²⁶ for the internal rotations of formamide, which is not obvious for hydrogen peroxide. Figure 15 depicts the profile of the maximum amount of charge transferred⁴²⁷ during the internal rotation of the enol group of guanine, which mimics the corresponding electrophilicity behavior, as expected. As shown¹⁵² in Figure 11, energy and electrophilicity show a strong linear correlation in this case as well. Internal rotation of the phenyl group induces the reactivity in *trans*- and *cis*-phenyldiazene. For both isomers, the planar conformers are the most electrophilic and transition states are the least electrophilic (the *cis*-isomer has a maximum ω value in one of the TSs), which accounts for the extra electronic delocalization in the transition states.⁴²⁸ This unexpected behavior may be rationalized⁴²⁶ by the fact that both μ and η are maxima at the TS.

6.3. Chemical Reactions

In the intramolecular rearrangement reactions, $\text{trans-N}_2\text{H}_2 \rightarrow \text{cis-N}_2\text{H}_2$ and $\text{F}_2\text{S}_2 \rightarrow \text{FSSF}$, it has been shown that (μ , η , ω) values at the TS are, respectively, (maximum, minimum,




Figure 16. Variation of the chemical potential, hardness, and electrophilicity along the reaction path of the $\text{CNH} \rightarrow \text{HCN}$ isomerization reaction. Reprinted with permission from ref 227. Copyright 2003 American Chemical Society.




Figure 17. Variation of the chemical potential, hardness, and electrophilicity along the reaction path of the symmetric oxygen to oxygen proton-transfer reaction in the $\text{HO}-\text{C}(=\text{O})-\text{C}(=\text{S})-\text{OH}$ system. Reprinted with permission from ref 227. Copyright 2003 American Chemical Society.




Figure 18. Variation of the chemical potential, hardness, and electrophilicity along the reaction path of the sulfur to oxygen proton transfer in the $\text{HS}-\text{C}(=\text{O})-\text{C}(=\text{S})-\text{OH}$ system. Reprinted with permission from ref 227. Copyright 2003 American Chemical Society.

minimum) and (minimum, minimum, maximum), as depicted in Figure 13. The behavior of the latter reaction is as expected. To gain further insights into the former reaction where μ is maximum and η is minimum, one needs to analyze²²⁷ the following second derivative

$$\frac{\partial^2 \omega}{\partial \lambda^2} = \frac{1}{\eta} \left(\frac{\partial \mu}{\partial \lambda} \right)^2 + \frac{\mu^2}{\eta^3} \left(\frac{\partial \eta}{\partial \lambda} \right)^2 - 2 \frac{\mu}{\eta^2} \left(\frac{\partial \mu}{\partial \lambda} \right) \left(\frac{\partial \eta}{\partial \lambda} \right) + \frac{\mu}{\eta} \left(\frac{\partial^2 \mu}{\partial \lambda^2} \right) - \frac{1}{2} \frac{\mu^2}{\eta^2} \left(\frac{\partial^2 \eta}{\partial \lambda^2} \right) \quad (35)$$

Because the first derivatives are zero at the extremal points, the exact nature of the extremal (maximum or minimum) in ω at the TS would be governed by the relative magnitudes of the last two terms of the above equation for this type of reaction or where μ is minimum and η is maximum at the TS.

Figures 16–18 represent²²⁷ some more interesting cases, respectively, in the $\text{CNH} \rightarrow \text{HCN}$ isomerization reaction, oxygen to oxygen proton transfer in $\text{HO}-\text{C}(=\text{O})-\text{C}(=\text{S})-\text{OH}$, and oxygen to sulfur proton transfer in $\text{HS}-\text{C}(=\text{O})-\text{C}(=\text{S})-\text{OH}$ thioxalic acid derivatives, highlighting the




Figure 19. Profiles of f_{Fa}^- ($f = f, s, \omega$), f_{Fb}^- ($f = f, s, \omega$), the bond order (BO), and the energy along the reaction path of the gas-phase S_N2 substitution: $F_a^- + CH_3-F_b \rightarrow F_a-CH_3 + F_b^-$. Reprinted with permission from ref 433. Copyright 2005 American Chemical Society.




Figure 20. Profiles of $f_{Fa}^- + f_{Fb}^-$ ($f = f, s, \omega$), $f_{Fa}^+ + f_{Fb}^+$ ($f = f, s, \omega$), and the bond order (BO) along the reaction path of the gas-phase S_N2 substitution: $F_a^- + CH_3-F_b \rightarrow F_a-CH_3 + F_b^-$. Reprinted with permission from ref 433. Copyright 2005 American Chemical Society.

validity of the above analysis. Similar findings on 1,3-intramolecular proton-transfer reactions in $HXNY \rightarrow XNYH$ ($X, Y = O, S$) have also been reported.^{429a}

If μ^2 does not change significantly (or decreases) and η increases (cf. MHP) the minimum electrophilicity ($\omega = \mu^2/2\eta$) criterion may be attached to a favorable chemical process. In many exchange reactions, one of the products is the least electrophilic (among the reactants and the products) and the average product electrophilicity is less than the average reactant electrophilicity,^{52d} which may be termed as the minimum electrophilicity principle as suggested by the preceding analysis.^{227,361,426} Even for different isomers, the hardest one would be the least electrophilic,^{429b} when the μ^2 behavior is as mentioned above, and hence will be the most stable. This analysis^{227,426} has been recently revisited by Morell et al.⁴³⁰ Chaquin⁴³¹ has used a classical electrostatics analogy for this purpose.

A thorough study has been performed to analyze the profiles of global and local reactivity descriptors during vibration, internal rotation, and chemical reaction.⁴³² Most of the characteristics discussed in this section are found to be valid. Variations in the Fukui functions and the atomic charges along the reaction path have also been studied.¹⁰¹ This analysis¹⁰¹ highlights the failure of the Fukui function and the usefulness of the charges in explaining hard–hard interactions.^{99–102}

Figures 19 and 20 present the profiles of various local reactivity descriptors,⁴³³ including ω_{Fa}^- and ω_{Fb}^- , along the reaction path (IRC) of the gas-phase S_N2 substitution: $F_a^- + CH_3-F_b \rightarrow F_a-CH_3 + F_b^-$. In ref 433, the Mulliken charges were made use of. The corresponding populations are used in Figures 19 and 20. Local reactivity descriptors pass through a point of inflection in the transition state. This inflection point coincides with the saddle point of the reaction




Figure 21. Profiles of energy (E) and different global reactivity descriptors (η , ω , and $\Delta\omega^\pm$) along the paths of the gas-phase thermoneutral (a and b), endothermic (c), exothermic (d), and (4 + 2)-cycloaddition (e) reactions. Reprinted with permission from ref 185. Copyright 2009 American Chemical Society.

and provides a link between bonding and reactivity. For a thermoneutral reaction, similar profiles for the bond-making and the bond-breaking processes intersect at the transition state. The importance of these local descriptors is vindicated through a comparison with the profiles of standard indicators such as energy and bond order. Free F_a^- is more reactive to start with, and it gradually becomes less reactive as it becomes bonded. On the other hand, bonded F_b^- is the least reactive at the beginning, and it becomes more and more

reactive as it is released during the course of the reaction to have the most reactive free F_b^- . At the transition state, both F_a^- and F_b^- are equally reactive, as expected for a thermoneutral reaction.⁴³³ Figure 21 depicts the profiles of net electrophilicity along the reaction paths of (a, b) thermoneutral, $F_a + HF_b \rightarrow HF_a + F_b$ and $HO_a + HO_bH \rightarrow HO_b + HO_aH$; (c) endothermic, $HNO \rightarrow HON$; (d) exothermic, $H_2OO \rightarrow HOOH$; and (e) (4 + 2)-cycloaddition, $CH_2=CH-CH=CH_2 + CH_2=CH_2 \rightarrow C_6H_{10}$, reactions. A minimum net

electrophilicity principle appears to be valid.¹⁸⁵ Much work is needed in obtaining a more transparent view of how bonding, reactivity, and dynamics complement one another.

7. Dynamical Variants

7.1. Quantum Fluid Density Functional Theory

As discussed in section 3, two important time dependent density-based quantum mechanical theories are quantum fluid dynamics (QFD)⁸⁷ and the quantum theory of motion (QTM).⁸⁸ The quantum domain behavior of classically chaotic systems has been studied by using these theories.^{434,435} Time dependent density functional theory (TDDFT)^{436,437} strengthens them. According to TDDFT, which offers a time dependent extension to the original Hohenberg–Kohn theorems, the mapping between the time dependent external potential, $v(\vec{r}, t)$, and the density, $\rho(\vec{r}, t)$, is uniquely invertible up to an additive trivial time dependent function in the potential. This implies that all the properties of a system are functionals of $\rho(\vec{r}, t)$ and the current density $j(\vec{r}, t)$, whose time evolution for any many-electron system is governed by two basic QFD equations:^{436,437} viz., the equation of continuity,

$$\frac{\partial \rho}{\partial t} + \nabla \cdot \vec{J} = 0 \quad (36a)$$

and a Euler-type equation of motion

$$\frac{\partial \vec{J}}{\partial t} = P[\rho(\vec{r}, t), \vec{J}(\vec{r}, t)] \quad (36b)$$

where P is a functional whose form cannot be ascertained from TDDFT. To have an approximate form for P , a quantum fluid density functional theory (QFDFT)^{242,244,438–441} has been proposed via an amalgamation of TDDFT and QFD. The basic equation in QFDFT is a generalized nonlinear Schrödinger equation (GNLSE) as follows (in a.u.),

$$\left[-\frac{1}{2}\nabla^2 + v_{\text{eff}}(\vec{r}, t) \right] \Phi(\vec{r}, t) = i \frac{\partial \Phi(\vec{r}, t)}{\partial t} \quad (37a)$$

where the effective potential may be written as

$$v_{\text{eff}}(\vec{r}, t) = \frac{\partial T_{\text{NW}}}{\partial \rho} + \frac{\partial E_{\text{xc}}}{\partial \rho} + \int \frac{\rho(\vec{r}', t)}{|\vec{r} - \vec{r}'|} d\vec{r}' + v_{\text{ext}}(\vec{r}, t) \quad (37b)$$

where T_{NW} and E_{xc} are the non-Weizsäcker part of the kinetic energy and the exchange-correlation energy functional, respectively. The density and the current density are related to $\Phi(\vec{r}, t)$ as follows

$$\rho(\vec{r}, t) = |\Phi(\vec{r}, t)|^2 \quad (38a)$$

and

$$\vec{J}(\vec{r}, t) = [\Phi_{\text{re}} \nabla \Phi_{\text{im}} - \Phi_{\text{im}} \nabla \Phi_{\text{re}}] \quad (38b)$$

This GNLSE has been alternatively derived via a stochastic quantization⁴³⁸ and has been solved to study ion–atom collisions^{438,439} and atom–field interactions.^{440,441} As discussed in section 5, these two processes may be considered to mimic the external perturbation a molecule experiences during a chemical reaction.

7.2. Atom–Field Interactions

The external potential, $v_{\text{ext}}(\vec{r}, t)$, in GNLSE (eq 37b) of this problem has been written^{150,440,441} for an atom in its ground and excited electronic states interacting with a z-polarized laser field of varying intensities and colors. Figure 22 presents the time dependence of the external field for three different field intensities for monochromatic and bichromatic pulses. The time evolution of μ and η shows that the in-phase oscillations are observed only when the external field intensity becomes appreciable. To start with, the electron density will have a spherical distribution due to the central nature of the nuclear Coulomb field. A tug-of-war between this and an axial laser field will begin once the latter is switched on. It has been shown¹⁵⁰ that η is less sensitive than μ . Only when the strength of the external laser field is large enough to overcome the effect of the nuclear field do the in-phase oscillations in the reactivity parameters start. The electron density becomes cylindrical, and an oscillating dipole results. Figures 23 and 24 clearly delineate¹⁵⁰ these aspects in the plots of time dependent ω and $1/\omega$, respectively. At the very large field intensity, the difference in response of μ and η is manifested in these plots. A corresponding analysis on the Rydberg states of hydrogen and helium atoms provides important insights into their chaotic ionization.^{246–248}

7.3. Ion–Atom Collisions

For the ion–atom collision problem, $v_{\text{ext}}(\vec{r}, t)$ comprises the electron–nuclear attraction potentials originating from the target and the projectile nuclei.^{43,58,59,241–245,438,439} Figures 25–27 present the time evolution⁴³ of μ , η , and α , respectively, for a collision between a proton and an X-atom/ion ($X = \text{He}, \text{Li}^+, \text{Be}^{2+}, \text{B}^{3+}, \text{C}^{4+}$) in various electronic states. The dynamic μ profile has been shown to divide the whole collision process into three distinct regimes: viz., approach, encounter, and departure. In the encounter regime, the actual reaction takes place where η maximizes and α minimizes, showing the validity of the MHP and MPP in a dynamical situation. The HSAB principle also has revealed itself in action, and the associated regioselectivity of a reaction has been analyzed.^{43,242} The dynamic ω profile resembles that of μ for different projectile velocities and impact parameters in both ground and excited states²⁵² and hence is not shown here.

7.4. Chemical Kinetics

Ever since Maynard et al.¹³ pointed out the linear relationship between the logarithm of the rate coefficients and the electrophilicity index associated with the reactions between HIV-1 nucleocapsid protein p7 and several electrophilic reagents, there has been a renewed interest in analyzing this behavior for a variety of reactions. Hydrolysis of carbenium ions,⁴⁴² Friedel–Crafts reactions,^{5,347} aminolysis of thiocarbonates,³⁵² etc. have been studied for this purpose. Figure 28 depicts a representative plot for a series of thiolcarbonates reacting with piperidine.³⁵²

In general, these correlations are obtained for a set of molecules of similar chemical reactivity and often the outliers are to be removed for obtaining a meaningful correlation.^{347b} A linear correlation between the rate constant and ω has also been proposed.²⁵⁴ Both logarithmic^{352,443} and linear^{273,442} relations between the Hammett constant and the electrophil-




Figure 22. Time evolution of the external electric field with different colors and intensities. Reprinted with permission from ref 150. Copyright 2001 American Chemical Society.

licity index have been reported. The experimental Hammett substituent constant correlates well⁴⁴³ with its theoretically calculated electronic contribution for a series of substituted alkenes, as shown in Figure 29.

Some other related studies include those on reactivities of carbon-centered radicals,⁴⁴⁴ dienophiles,³²⁸ carbon–carbon double bonds,³⁴⁶ metal polypyridyl complexes,³⁶⁶ and some radicals toward hydrogen abstraction.⁴⁴⁵ A slightly different version of Parr et al.'s definition has been used for the calculation of absolute rates of atom abstractions by radicals^{446,447} as well as ultrafast excited-state proton transfer.⁴⁴⁸ The quality of the leaving groups has been quantified^{220a} in terms of electrofugality and nucleofugality, which help in analyzing the reactivities associated with electrophilic and nucleophilic processes. Electro- (nucleo)-philicity may be considered to be a kind of inverse of nucleo- (electro)fugality.^{123,220} Although there have been some^{220a–g} attempts in this regard, it needs to be analyzed whether we are comparing thermodynamic and/or kinetic properties.^{1–3} It is also to be studied whether a good electrophile or a poor nucleophile will be a good nucleofuge and vice versa. As was pointed out,^{220a} it has been numerically verified that the definition is not always useful.^{220f}

8. Spin Dependent Generalizations

8.1. $\{N, N_s, v(\vec{r})\}$ Representation

The reactivity and selectivity descriptors defined so far are inadequate in the study of reactions involving changes

in spin multiplicity, including spin catalysis. A spin polarized version of DFT has been developed for this purpose.^{449–454} For a system with N_α and N_β numbers of α and β spins, respectively, with the corresponding densities ρ_α and ρ_β , the electron density, $\rho(\vec{r})$, the spin density, $\rho_s(\vec{r})$, the electron number, N , and the spin number, N_s , are defined as^{449–454}

$$\rho(\vec{r}) = \rho_\alpha(\vec{r}) + \rho_\beta(\vec{r}) \quad (39a)$$

$$\rho_s(\vec{r}) = \rho_\alpha(\vec{r}) - \rho_\beta(\vec{r}) \quad (39b)$$

$$N = N_\alpha + N_\beta \quad (39c)$$

$$N_s = N_\alpha - N_\beta \quad (39d)$$

The corresponding chemical potential (μ_N) and spin potential (μ_s) are given by^{449–454}

$$\mu_N = \left(\frac{\partial E}{\partial N} \right)_{N_s, v(\vec{r})} \quad (40a)$$

$$\mu_s = \left(\frac{\partial E}{\partial N_s} \right)_{N, v(\vec{r})} \quad (40b)$$

The related hardness parameters are defined as^{449–454}

$$\eta_{NN} = \left(\frac{\partial \mu_N}{\partial N} \right)_{N_s, v(\vec{r})} \quad (41a)$$




Figure 23. Time evolution of electrophilicity index (ω) of a helium atom in ground and excited states in the presence of external electric fields with different colors and intensities. Reprinted with permission from ref 150. Copyright 2001 American Chemical Society.

$$\eta_{\text{NS}} = \left(\frac{\partial \mu_{\text{N}}}{\partial N_{\text{s}}} \right)_{N_{\text{s}}, v(\vec{r})} = \left(\frac{\partial \mu_{\text{s}}}{\partial N} \right)_{N_{\text{s}}, v(\vec{r})} = \eta_{\text{SN}} \quad (41\text{b})$$

$$\eta_{\text{SS}} = \left(\frac{\partial \mu_{\text{s}}}{\partial N_{\text{s}}} \right)_{N_{\text{s}}, v(\vec{r})} \quad (41\text{c})$$

Specific combinations⁴⁵⁵ of the above quantities in the spirit of eq 12 will yield various spin dependent generalizations of the electrophilicity index. Spin philicity and spin donicity have been defined by Pérez et al.,⁴⁵⁶ extended by Melin et al.,⁴⁵⁷ and revised by Olah et al.^{458,459} in this connection.

Spin dependent philicities can also be obtained⁴⁵⁵ by knowing the above quantities and the various Fukui functions in this representation as described below:

$$f_{\text{NN}}(\vec{r}) = \left(\frac{\partial \rho(\vec{r})}{\partial N} \right)_{N_{\text{s}}, v(\vec{r})} = \left(\frac{\partial \mu_{\text{N}}}{\partial v(\vec{r})} \right)_{N_{\text{s}}, N_{\text{s}}} \quad (42\text{a})$$

$$f_{\text{NS}}(\vec{r}) = \left(\frac{\partial \rho(\vec{r})}{\partial N_{\text{s}}} \right)_{N_{\text{s}}, v(\vec{r})} = \left(\frac{\partial \mu_{\text{s}}}{\partial v(\vec{r})} \right)_{N_{\text{s}}, N_{\text{s}}} \quad (42\text{b})$$

and similarly for $f_{\text{SN}}(\vec{r})$ and $f_{\text{SS}}(\vec{r})$. Some of these local quantities have been calculated⁴⁶⁰ in analyzing the regioselectivity in the [2 + 2]-photocycloaddition of enones to substituted alkenes.

8.2. $\{N_{\alpha}, N_{\beta}, v(\vec{r})\}$ Representation

An alternative representation may be developed⁴⁵⁵ by defining various global and local reactivity descriptors

explicitly for the spin-up and spin-down cases as follows:

Chemical potentials

$$\mu_{\alpha} = \left(\frac{\partial E}{\partial N_{\alpha}} \right)_{N_{\beta}, v(\vec{r})}; \mu_{\beta} = \left(\frac{\partial E_{\beta}}{\partial N_{\beta}} \right)_{N_{\alpha}, v(\vec{r})} \quad (43\text{a})$$

Hardness

$$\eta_{\alpha\alpha} = \left(\frac{\partial \mu_{\alpha}}{\partial N_{\alpha}} \right)_{N_{\beta}, v(\vec{r})}; \eta_{\beta\beta} = \left(\frac{\partial \mu_{\beta}}{\partial N_{\beta}} \right)_{N_{\alpha}, v(\vec{r})} \quad (43\text{b})$$

$$\eta_{\alpha\beta} = \left(\frac{\partial \mu_{\alpha}}{\partial N_{\beta}} \right)_{N_{\alpha}, v(\vec{r})} = \left(\frac{\partial \mu_{\beta}}{\partial N_{\alpha}} \right)_{N_{\beta}, v(\vec{r})} = \eta_{\beta\alpha} \quad (43\text{c})$$

Fukui functions

$$f_{\alpha\alpha}(\vec{r}) = \left(\frac{\partial \rho_{\alpha}(\vec{r})}{\partial N_{\alpha}} \right)_{N_{\beta}, v(\vec{r})}; f_{\beta\beta}(\vec{r}) = \left(\frac{\partial \rho_{\beta}(\vec{r})}{\partial N_{\beta}} \right)_{N_{\alpha}, v(\vec{r})} \quad (44\text{a})$$

$$f_{\alpha\beta}(\vec{r}) = \left(\frac{\partial \rho_{\alpha}(\vec{r})}{\partial N_{\beta}} \right)_{N_{\alpha}, v(\vec{r})}; f_{\beta\alpha}(\vec{r}) = \left(\frac{\partial \rho_{\beta}(\vec{r})}{\partial N_{\alpha}} \right)_{N_{\beta}, v(\vec{r})} \quad (44\text{b})$$




Figure 24. Time evolution of nucleophilicity index ($1/\omega$) of a helium atom in ground and excited states in the presence of external electric fields with different colors and intensities. Reprinted with permission from ref 150. Copyright 2001 American Chemical Society.




Figure 25. Dynamical chemical potential profile during a collision process between an X-atom/ion ($X = \text{He}, \text{Li}^+, \text{Be}^{2+}, \text{B}^{3+}, \text{C}^{4+}$) in its ground state and a proton. Reprinted with permission from ref 43. Copyright 2003 American Chemical Society.

Making use of the above quantities, the spin dependent variants of the electrophilicity index and the philicity may be easily derived.⁴⁵⁵ The representation to be used would depend⁴⁵⁵ on the nature of the spin multiplicity change in the given spin polarized process.

9. Conclusions

The tremendous power of the electrophilicity index proposed by Parr, Szentpaly, and Liu provides insight into almost every arena of chemistry and encompasses informa-




Figure 26. Dynamical hardness profile during a collision process between an X-atom/ion ($X = \text{He}, \text{Li}^+, \text{Be}^{2+}, \text{B}^{3+}, \text{C}^{4+}$) in various electronic states and a proton. Reprinted with permission from ref 43. Copyright 2003 American Chemical Society.




Figure 27. Dynamical polarizability profile during a collision process between an X-atom/ion ($X = \text{He}, \text{Li}^+, \text{Be}^{2+}, \text{B}^{3+}, \text{C}^{4+}$) and a proton. Reprinted with permission from ref 43. Copyright 2003 American Chemical Society.




Figure 28. Plot of the experimental nucleophilic rate coefficient (k_N) for a series of thiolcarbonates reacting with piperidine, versus electrophilicity index. Reprinted with permission from ref 352. Copyright 2005 American Chemical Society.

tion about the structure, properties, stability, reactivity, interactions, bonding, toxicity, and dynamics of many-electron systems in ground and excited electronic states. The




Figure 29. Plot of the experimental Hammett constant (σ_p) of a series of substituted ethylenes versus its theoretical counterpart ($\sigma_p(\omega)$). Reprinted with permission from ref 443. Copyright 2003 American Chemical Society.

recurring theme of this review has been the electrophilicity concept in general, with the common thread being the Maynard–Parr electrophilicity index. The whole gamut of the conceptual density functional theory lends support toward the electrophilicity index and helps it realize its full potential. As is the case for most of the conceptual DFT-based reactivity/selectivity descriptors, the global and local electrophilicities possess strong interpretive power, which itself is important in understanding a diverse class of biophysics-chemical processes. However, their predictive capacity has yet to be assessed. We conclude with the optimistic note that the electrophilicity will exhibit its tremendous predictive potential, which, along with its existing interpretive characteristics, together with those of the other descriptors, will be adequate in developing a complete theory of chemical reactivity.

10. Abbreviations and Symbols of Some Important Subjects/Quantities

A	electron affinity
α	polarizability
AIM	atoms-in-molecules
B3LYP	Becke three-parameter Lee–Yang–Parr functional
$[\partial\eta/\partial N]$	variation of hardness with electron number
ΔE_{ins}	insertion energy
ΔE_{solv}	energy of solvation
DFT	density functional theory
Δ_{ij}^{kl}	softness matching index
DNA	DNA
$[\partial\omega/\partial N]$	variation of electrophilicity with electron number
$\varepsilon_{\text{HOMO}}$	highest occupied molecular orbital energy
ELF	electron localization function
$\varepsilon_{\text{LUMO}}$	lowest unoccupied molecular orbital energy
E_{xc}	exchange–correlation energy functionals
FF	Fukui function
f_k^α	condensed Fukui function
$F[\rho]$	Hohenberg–Kohn–Sham universal functional
$f(\vec{r})$	Fukui function
η	hardness

GNLSE	generalized nonlinear Schrödinger equation
$\eta(\vec{r}, \vec{r}')$	hardness kernel
HF	Hartree–Fock
HIV	human immunodeficiency virus
HOMA	harmonic oscillator model of aromaticity
HOMO	highest occupied molecular orbital
HSAB	hard and soft acids and bases
I	ionization potential
IC ₅₀	50% inhibitory concentration
IGC ₅₀	50% inhibitory growth concentration
IRC	intrinsic reaction coordinate
$j(\vec{r}, t)$	current density
K	equilibrium constant
k	rate constant
k_{σ}	hydrogen bond strength
LUMO	lowest unoccupied molecular orbital
μ	chemical potential
m_{CXY}	carbene-philicity scale
MHP	maximum hardness principle
MPP	minimum polarizability principle
μ_s	spin potential
N	number of electrons
$v_{\text{ext}}(\vec{r}, t)$	time dependent external potential
NICS	nucleus independent chemical shift
NMR	nuclear magnetic resonance
NPA	natural population analysis
$v(\vec{r})$	external potential
N_s	spin number
PCB	polychlorinated biphenyl
PCDD	polychlorinated dibenzo- <i>p</i> -dioxin
PCDF	polychlorinated dibenzofuran
QFD	quantum fluid dynamics
QSA	quantitative structure–activity relationship
QTM	quantum theory of motion
RBA	relative binding affinity
$\rho(\vec{r})$	electron density
ρ_s	spin density
S	softness
s_k^{α}	condensed softness
$S_{\text{N}2}$	substitution nucleophilic bimolecular
$s(\vec{r})$	local softness
TDDFT	time dependent density functional theory
TeBG	testosterone–estrogen binding globulin
T_{F}	Thomas–Fermi kinetic energy density
T_{NW}	non-Weizsäcker part of the kinetic energy
TS	transition state
V_{qu}	quantum potential
$V_{\text{qu}}(r, t)$	time dependent quantum potential
$\Phi(r, t)$	3-D hydrodynamical wave function
χ	electronegativity
$\Psi(\vec{x}_1, \vec{x}_2, \dots, \vec{x}_N)$	many particle wave function for an N -electron system
ω	electrophilicity index
ω^{\pm}	directional electrophilicity index
ω_k^{α}	condensed philicity
$\omega(\vec{r})$	philicity
$\omega^{\pm}(\vec{r})$	directional philicity
ω^{+}	electroaccepting power
ω^{-}	electrodonating power
$\Delta\omega^{\pm}$	net electrophilicity

11. Acknowledgments

One of us (P.K.C.) would like to express his deep sense of gratitude to three of his teachers, Professors S. C. Rakshit, B. M. Deb, and R. G. Parr, for kindly introducing him to the worlds of quantum mechanics and formal and conceptual density functional theory, respectively. Several of his teachers, collaborators, coauthors, friends, and students from different countries such as Belgium, Canada, Chile, China, **France**, Germany, India, Mexico, Spain,

U.K., U.S.A., etc. have been involved in his quest for understanding the behavior of atoms and molecules. He thanks all of them including Professors, Drs., and Drs.-to-be F. Aparicio, E. M. Arnett, T. V. S. Arun Murthy, P. Ayers, **S. Banerjee**, M. Bhattacharjee, S. Bhattacharya, P. Bultinck, A. Cedillo, **A. Chakraborty**, D. Chakraborty, E. Chamorro, K. Chitra, R. Contreras, M. P. Das, **R. Das**, **S. Dasadhikari**, **D. Deb**, **A. A. Denopoulos**, F. De Proft, **J. Dinda**, M. Elango, P. Fuentealba, M. Galván, J. Garza, P. Geerlings, S. K. Ghosh, B. Gomez, N. Gonzalez-Rivas, **P. Goyal**, **A. Gupta**, K. Gupta, S. Gutierrez-Oliva, S. Haque, M. K. Harbola, **R. Islas**, **M. Jaccob**, P. Jaque, **L. Jhulki**, **D. Jouvenot**, **M. Khatua**, S. Khatua, S. R. Koneru, **B. S. Krishnamoorthy**, **R. P. S. A. Kumar**, R. Lalitha, C. Lee, H. Lee, G. H. Liu, S. Liu, **F. Loiseau**, B. Maiti, **A. K. Mandal**, **G. Merino**, **S. Mondal**, J. Martinez-Magadan, M. Matus, J. Melin, S. Mishra, A. Mitra, **P. Mitra**, A. Mukherjee, S. Mukherjee, P. K. Nandi, G. K. Narayanan, S. Nath, J. Padmanabhan, N. Pal, S. Pal, K. Panchanatheswaran, **A. Pandith**, R. Parthasarathi, **R. G. Pearson**, P. Pérez, A. Poddar, **B. Rana**, **P. Ravinder**, **D. R. Roy**, **G. S. Roymahapatra**, A. M. Sabeelullah, **T. Samanta**, A. B. Sannigrahi, J. Santos, U. Sarkar, **B. Sateesh**, P. v. R. Schleyer, K. L. Sebastian, S. Sengupta, Y. Simon-Manso, H. Singh, **K. Srinivasu**, **H. K. Srivastava**, V. Subramanian, A. Tanwar, **A. J. Thakkar**, A. Toro-Labbé, M. Torrent-Sucarrat, S. Vandamme, R. Vargas, N. S. Venkata-Subramaniyan, **P. Venuvanalingam**, **J. L. Vigneresse**, **R. Vijayaraj**, **D. Z. Wang**, J. Zevallos, and Z. Zhou. Special thanks are due to Professors W. Kohn, R. G. Pearson, W. Yang, and P. Geerlings for constant encouragement and Professor R. F. W. Bader for helpful discussion. **One of us (P.K.C.) would like to thank the Department of Science and Technology, New Delhi, for the J. C. Bose Fellowship and Professor J. Michl for kindly inviting him to write this update.** P.K.C. also thanks his wife Samhita and daughter Saparya for their help in various ways. Financial assistance from CSIR, New Delhi, and **Indo–EU project HYPOMAP** is gratefully acknowledged.

12. References

- (1) Carey, F. A.; Sundberg, R. J. *Advanced Organic Chemistry; Part B: Reactions and Synthesis*, 4th ed.; Kluwer Academic/Plenum Publishers: New York, 2001.
- (2) Lowry, T. H.; Richardson, K. S. *Mechanism and Theory in Organic Chemistry*, 3rd ed.; Harper & Row: New York, 1987.
- (3) Smith, M. B.; March, J. *Advanced Organic Chemistry: Reactions, Mechanisms, and Structure*, 5th ed.; John Wiley & Sons: New York, 2001.
- (4) Sykes, P. *A Guidebook to Mechanism in Organic Chemistry*, 6th ed.; Orient Longman Limited: New Delhi, 1970.
- (5) Olah, G. A. *Acc. Chem. Res.* **1971**, 4, 240.
- (6) Kane-Maguire, L. A. P.; Honig, E. D.; Sweigart, D. A. *Chem. Rev.* **1984**, 84, 525.
- (7) Smit, W. A.; Caple, R.; Smoliakova, I. P. *Chem. Rev.* **1994**, 94, 2359.
- (8) Holmes, R. R. *Chem. Rev.* **1996**, 96, 927.
- (9) (a) Lal, G. S.; Pez, G. P.; Syvret, R. G. *Chem. Rev.* **1996**, 96, 1737. (b) Umemoto, T. *Chem. Rev.* **1996**, 96, 1757.
- (10) Chen, J.-T. *Coord. Chem. Rev.* **1999**, 190, 1143.
- (11) Sander, W.; Kötting, C.; Hübert, R. *J. Phys. Org. Chem.* **2000**, 13, 561.
- (12) Makosza, M.; Wojciechowski, K. *Chem. Rev.* **2004**, 104, 2631.
- (13) Maynard, A. T.; Huang, M.; Rice, W. G.; Covell, D. G. *Proc. Natl. Acad. Sci. U.S.A.* **1998**, 95, 11578.
- (14) Parr, R. G.; Szentpály, L. v.; Liu, S. *J. Am. Chem. Soc.* **1999**, 121, 1922.
- (15) Pauling, L. *The Nature of the Chemical Bond*, 3rd ed.; Cornell University Press: Ithaca, NY, 1960.
- (16) *Structure and Bonding*, Vol. 66: *Electronegativity*; Sen, K. D., Jorgenson, C. K., Eds.; Springer: Berlin, 1987.

- (17) Pearson, R. G. *Chemical Hardness: Applications from Molecules to Solids*; Wiley-VCH: Weinheim, Germany, 1997.
- (18) *Structure and Bonding*, Vol. 80: *Chemical Hardness*; Sen, K. D., Mingos, D. M. P., Eds.; Springer: Berlin, 1993.
- (19) Parr, R. G.; Yang, W. *Density Functional Theory of Atoms and Molecules*; Oxford University Press: Oxford, U.K., 1989.
- (20) Geerlings, P.; De Proft, F.; Langenaeker, W. *Chem. Rev.* **2003**, *103*, 1793.
- (21) (a) Chermette, H. *J. Comput. Chem.* **1999**, *20*, 129. (b) Gázquez, J. L. *J. Mex. Chem. Soc.* **2008**, *52*, 3. (c) Roos, G.; Geerlings, P.; Messens, J. *J. Phys. Chem. B* **2009**, *113*, 13465. (d) Liu, S. *Acta Phys. Chim. Sin.* **2009**, *25*, 590.
- (22) Chattaraj, P. K.; Parr, R. G. Density Functional Theory of Chemical Hardness. In *Chemical Hardness, Structure and Bonding*; Sen, K. D., Mingos, D. M. P., Eds.; Springer-Verlag: Berlin, 1993; Vol. 80, pp 11–25.
- (23) Chattaraj, P. K.; Poddar, A.; Maiti, B. Chemical Reactivity and Dynamics within a Density-Based Quantum Mechanical Framework. In *Reviews in Modern Quantum Chemistry: A Celebration of the Contributions of Robert Parr*; Sen, K. D., Eds.; World Scientific: Singapore, 2002; pp 871–935.
- (24) (a) Chattaraj, P. K.; Nath, S.; Maiti, B. Reactivity Descriptors. In *Computational Medicinal Chemistry for Drug Discovery*; Tollenaere, J., Bultinck, P., Winter, H. D., Langenaeker, W., Eds.; Marcel Dekker: New York, 2003; Chapter 11, pp 295–322. (b) Chattaraj, P. K., Ed. *Chemical Reactivity Theory: A Density Functional View*; Taylor and Francis: Boca Raton, FL, 2009. (c) Chattaraj, P. K.; Giri, S. *Annu. Rep. Prog. Chem., Sect. C* **2009**, *105*, 13. (d) Chattaraj, P. K., Thakkar, A. J., Eds. Special Issue on Conceptual aspects of electron densities and density functionals. *THEOCHEM* **2010**.
- (25) Chattaraj, P. K. *J. Ind. Chem. Soc.* **1992**, *69*, 173. [See also the Special Issue of *J. Chem. Sci.* on Chemical Reactivity, 2005, Vol. 117, Guest Editor: Chattaraj, P. K. There are several related papers in this issue, which include the following: Melin, J.; Ayers, P. W.; Ortiz, J. V. *J. Chem. Sci.* **2005**, *117*, 387. Chamorro, E.; Duque, M.; Cardenas, C.; Santos, J.; Tiznado, W.; Fuentealba, P. *J. Chem. Sci.* **2005**, *117*, 419. Liu, S. *J. Chem. Sci.* **2005**, *117*, 477. Korchowiec, J. *J. Chem. Sci.* **2005**, *117*, 491. Tanwar, A.; Pal, S. *J. Chem. Sci.* **2005**, *117*, 497. Cardenas-Jiron, G. I.; Espinoza-Leyton, F.; Sordo, T. L. *J. Chem. Sci.* **2005**, *117*, 515. For some recent applications, see: Gomez, B.; Likhanova, N. V.; Dominguez-Aguilar, M. A.; Martinez-Palou, R.; Vela, A.; Gázquez, J. L. *J. Phys. Chem. B* **2006**, *110*, 8928. Tian, S. X.; Yang, J. *Angew. Chem., Int. Ed.* **2006**, *45*, 2069. DELETED (Now 320b)] A philicity based site activation model has been adopted in understanding the Markovnikov rule (Padmanabhan, J.; Parthasarathi, R.; Subramanian, V.; Chattaraj, P. K. *THEOCHEM* **2007**, *804*, 17). by extending a similar local softness based analysis (Aizman, A.; Contreras, R.; Galván, M.; Cedillo, A.; Santos, J. C.; Chamorro, E. *J. Phys. Chem. A* **2002**, *106*, 7844).
- (26) Kohn, W.; Becke, A. D.; Parr, R. G. *J. Phys. Chem.* **1996**, *100*, 12974.
- (27) Hohenberg, P.; Kohn, W. *Phys. Rev. B* **1964**, *136*, 864.
- (28) Kohn, W.; Sham, L. J. *Phys. Rev. A* **1965**, *140*, 1133.
- (29) Parr, R. G.; Donnelly, R. A.; Levy, M.; Palke, W. E. *J. Chem. Phys.* **1978**, *68*, 3801.
- (30) Parr, R. G.; Pearson, R. G. *J. Am. Chem. Soc.* **1983**, *105*, 7512. It has been shown here that the charge transfer not only depends on the electronegativity difference but is also inversely proportional to the hardness sum.
- (31) Cohen, M. H.; Wasserman, A. *J. Phys. Chem. A* **2007**, *111*, 2229. See also: Ayers, P. W. *J. Math. Chem.* **2008**, *43*, 285.
- (32) Chattaraj, P. K.; Cedillo, A.; Parr, R. G. *Chem. Phys.* **1996**, *204*, 429.
- (33) Cedillo, A.; Chattaraj, P. K.; Parr, R. G. *Int. J. Quantum Chem.* **2000**, *77*, 403.
- (34) Sanderson, R. T. *Science* **1951**, *114*, 670. [See Bultinck, P.; Vanholme, R.; Popelier, P. L. A.; De Proft, F.; Geerlings, P. *J. Phys. Chem. A* **2004**, *108*, 10359. to know the present day relevance of this principle in the charge calculation and. Glasser, L.; von Szentpaly, L. *J. Am. Chem. Soc.* **2006**, *128*, 12314. for a generalized Born–Haber–Fajans cycle based on that].
- (35) Sanderson, R. T. *Science* **1955**, *121*, 207.
- (36) Sanderson, R. T. *J. Chem. Educ.* **1954**, *31*, 238.
- (37) Pearson, R. G. *Hard and Soft Acids and Bases*; Dowden, Hutchinson & Ross: Stroudsburg, PA, 1973.
- (38) Pearson, R. G. *Coord. Chem. Rev.* **1990**, *100*, 403.
- (39) (a) Chattaraj, P. K.; Lee, H.; Parr, R. G. *J. Am. Chem. Soc.* **1991**, *113*, 1855. See ref 33 to see the external potential effects on this principle. (b) Pearson, R. G.; Chattaraj, P. K. *Chemtracts–Inorg. Chem.* **2009**, *21*, 1.
- (40) Ayers, P. W. *J. Chem. Phys.* **2005**, *122*, 141102. [See Chattaraj, P. K.; Ayers, P. W. *J. Chem. Phys.* **2005**, *123*, 086101 for a connection between MHP and HSABP.] Note that the arithmetic mean principle for softness and MHP together imply HSABP. Corresponding proof for the half-reactions is available in Ayers, P. W.; Parr, R. G.; Pearson, R. G. *J. Chem. Phys.* **2006**, *124*, 194107. An MHP variant is also attempted. Chattaraj, P. K.; Ayers, P. W.; Melin, J. *Phys. Chem. Chem. Phys.* **2007**, *9*, 3853.
- (41) Chattaraj, P. K.; Schleyer, P. v. R. *J. Am. Chem. Soc.* **1994**, *116*, 1067. Chattaraj, P. K.; Nath, S. *Ind. J. Chem. A* **1994**, *33A*, 842. See refs 39 and 41 for a connection between MHP and HSABP.
- (42) Chattaraj, P. K.; Gomez, B.; Chamorro, E.; Santos, J.; Fuentealba, P. *J. Phys. Chem. A* **2001**, *105*, 8815.
- (43) Chattaraj, P. K.; Maiti, B. *J. Am. Chem. Soc.* **2003**, *125*, 2705.
- (44) Pearson, R. G. *J. Chem. Educ.* **1987**, *64*, 561.
- (45) Parr, R. G.; Chattaraj, P. K. *J. Am. Chem. Soc.* **1991**, *113*, 1854.
- (46) Chattaraj, P. K.; Liu, G. H.; Parr, R. G. *Chem. Phys. Lett.* **1995**, *237*, 171.
- (47) Chattaraj, P. K. *Proc. Indian Natl. Sci. Acad., Part A* **1996**, *62*, 513 (review).
- (48) Ayers, P. W.; Parr, R. G. *J. Am. Chem. Soc.* **2000**, *122*, 2010.
- (49) Chattaraj, P. K.; Fuentealba, P.; Gomez, B.; Contreras, R. *J. Am. Chem. Soc.* **2000**, *122*, 348.
- (50) Pearson, R. G. *Acc. Chem. Res.* **1993**, *26*, 250.
- (51) Parr, R. G.; Zhou, Z. *Acc. Chem. Res.* **1993**, *26*, 256.
- (52) (a) Politzer, P. *J. Chem. Phys.* **1987**, *86*, 1072. (b) Tanwar, A.; Pal, S.; Roy, D. R.; Chattaraj, P. K. *J. Chem. Phys.* **2006**, *125*, 056101. (c) Chattaraj, P. K.; Arun Murthy, T. V. S.; Giri, S.; Roy, D. R. *THEOCHEM* **2007**, *813*, 63. (d) For chemical reactions, see Chattaraj, P. K.; Roy, D. R.; Giri, S. *Comput. Lett.* **2007**, *3*, 223 (A. D. Buckingham Special issue).
- (53) Ghanty, T. K.; Ghosh, S. K. *J. Phys. Chem.* **1993**, *97*, 4951.
- (54) Fuentealba, P.; Reyes, O. *THEOCHEM* **1993**, *282*, 65.
- (55) Simon-Manso, Y.; Fuentealba, P. *J. Phys. Chem. A* **1998**, *102*, 2029.
- (56) Chattaraj, P. K.; Sengupta, S. *J. Phys. Chem.* **1996**, *100*, 16126.
- (57) Ghanty, T. K.; Ghosh, S. K. *J. Phys. Chem.* **1996**, *100*, 12295.
- (58) Chattaraj, P. K.; Sengupta, S. *J. Phys. Chem. A* **1997**, *101*, 7893.
- (59) Chattaraj, P. K.; Fuentealba, P.; Jaque, P.; Toro-Labbé, A. *J. Phys. Chem. A* **1999**, *103*, 9307.
- (60) Chattaraj, P. K.; Giri, S.; Duley, S. *J. Phys. Chem. Lett.* **2010**, *1*, 1064.
- (61) (a) Pearson, R. G. *Inorg. Chem.* **1988**, *27*, 734. (b) Pearson, R. G. *Proc. Natl. Acad. Sci. U.S.A.* **1986**, *83*, 8440. (c) Cohen, A. J.; Mori-Sanchez, P.; Yang, W. T. *Phys. Rev. B* **2008**, *77*, 115123.
- (62) Koopmans, T. A. *Physica* **1933**, *1*, 104.
- (63) (a) Huang, M.; Maynard, A.; Turpin, J. A.; Graham, L.; Janini, G. M.; Covell, D. G.; Rice, W. G. *J. Med. Chem.* **1998**, *41*, 1371. (b) Perdew, J. P.; Parr, R. G.; Levy, M.; Balduz, J. L., Jr. *Phys. Rev. Lett.* **1982**, *49*, 1691. (c) Gázquez, J. L.; Cedillo, A.; Vela, A. *J. Phys. Chem. A* **2007**, *111*, 1966. (d) Ayers, P. W.; De Proft, F.; Borgoo, A.; Geerlings, P. *J. Chem. Phys.* **2007**, *126*, 224107. (e) Sablon, N.; De Proft, F.; Ayers, P. W.; Geerlings, P. *J. Chem. Phys.* **2007**, *126*, 224108.
- (64) Gijū, K. T.; De Proft, F.; Geerlings, P. *J. Phys. Chem. A* **2005**, *109*, 2925.
- (65) Chattaraj, P. K.; Duley, S. *J. Chem. Eng. Data* **2010**, *55*, 1882. Ground state analysis of metastable anions/dianions (with negative electronegativity values) in gas phase becomes problematic. Even nonmetallic elements sometimes exhibit negative electron affinities; see Pearson, R. G. *Inorg. Chem.* **1991**, *30*, 2856.
- (66) Berkowitz, M.; Ghosh, S. K.; Parr, R. G. *J. Am. Chem. Soc.* **1985**, *107*, 6811.
- (67) Ghosh, S. K.; Berkowitz, M. *J. Chem. Phys.* **1985**, *83*, 2976.
- (68) (a) Harbola, M. K.; Chattaraj, P. K.; Parr, R. G. *Isr. J. Chem.* **1991**, *31*, 395. (b) Chattaraj, P. K.; Roy, D. R.; Geerlings, P.; Torrent-Sucarrat, M. *Theor. Chem. Acc.* **2007**, *118*, 923.
- (69) Ghosh, S. K. *Chem. Phys. Lett.* **1990**, *172*, 77.
- (70) Langenaeker, W.; De Proft, F.; Geerlings, P. *J. Phys. Chem.* **1995**, *99*, 6424.
- (71) Fukui, K. *Theory of Orientation and Stereoselection*; Springer: Berlin, 1973; p 134.
- (72) Fukui, K. *Science* **1982**, *218*, 747.
- (73) Fukui, K.; Yonezawa, Y.; Shingu, H. *J. Chem. Phys.* **1952**, *20*, 722.
- (74) Parr, R. G.; Yang, W. *J. Am. Chem. Soc.* **1984**, *106*, 4049.
- (75) Ayers, P. W.; Levy, M. *Theor. Chem. Acc.* **2000**, *103*, 353.
- (76) DELETED [Now 59b].
- (77) Chattaraj, P. K.; Cedillo, A.; Parr, R. G. *J. Chem. Phys.* **1995**, *103*, 10621.
- (78) Pacios, L. F. *Chem. Phys. Lett.* **1997**, *276*, 381.
- (79) Chattaraj, P. K.; Cedillo, A.; Parr, R. G. *J. Chem. Phys.* **1995**, *103*, 7645.
- (80) De Proft, F.; Geerlings, P.; Liu, S.; Parr, R. G. *Pol. J. Chem.* **1999**, *72*, 1737.
- (81) Bader, R. F. W.; MacDougall, P. J.; Lau, C. D. H. *J. Am. Chem. Soc.* **1984**, *106*, 1594.
- (82) Bader, R. F. W. *Atoms in Molecules: a Quantum Theory*; Oxford University Press: Oxford, 1990.

- (83) Naray-Szabo, G.; Ferenczy, G. G. *Chem. Rev.* **1995**, *95*, 829.
- (84) Gadre, S. R.; Kulkarni, S. A.; Srivastava, I. H. *J. Chem. Phys.* **1992**, *96*, 5253.
- (85) Murray, J. S.; Politzer, P. *Theor. Chim. Acta* **1987**, *72*, 507.
- (86) Koster, A. M.; Kolle, C.; Jug, K. J. *Chem. Phys.* **1993**, *99*, 1244.
- (87) Madelung, E. Z. *Phys.* **1926**, *40*, 322.
- (88) (a) Holland, P. R. *The Quantum Theory of Motion*; Cambridge University Press: Cambridge, 1993. (b) Wyatt, R. E. *Quantum Dynamics with Trajectories: Introduction to Quantum Hydrodynamics*; Springer: New York, 2007. (c) Chattaraj, P. K., Ed. *Quantum Trajectories*; Taylor & Francis/CRC Press: Boca Raton, FL, 2010 (in press).
- (89) Becke, A. D.; Edgecombe, K. E. *J. Chem. Phys.* **1990**, *92*, 5397.
- (90) Savin, A.; Becke, A. D.; Flad, J.; Nesper, R.; Preuss, H.; Von Schnering, H. G. *Angew. Chem.* **1991**, *103*, 421. Silvi, B.; Savin, A. *Nature* **1994**, *371*, 683. Chesnut, D. B.; Savin, A. *J. Am. Chem. Soc.* **1999**, *121*, 2335. Fuentealba, P.; Savin, A. *J. Phys. Chem. A* **2001**, *105*, 11531.
- (91) Fuster, F.; Silvi, B. *J. Phys. Chem. A* **2000**, *104*, 947.
- (92) Chattaraj, P. K.; Chamorro, E.; Fuentealba, P. *Chem. Phys. Lett.* **1999**, *314*, 114. [See also Ghosh, S. K.; Berkowitz, M.; Parr, R. G. *Proc. Natl. Acad. Sci. U.S.A.* **1984**, *81*, 8028. Ayers, P. W.; Parr, R. G.; Nagy, A. *Int. J. Quantum Chem.* **2002**, *90*, 309.] A condensed-to-atom variant of this local temperature is being explored (Chakraborty, D.; Giri, S.; Roy, D. R.; Bultinck, P.; Chattaraj, P. K. Unpublished work). **Although the condensed local temperature correlates well with several physico-chemical properties, the condensation becomes mathematically inexact due to the nonlinearity of the kernel. A mathematically inexact and ambiguous local hardness has been proposed [Saha, S.; Roy, R. K. *J. Phys. Chem. B* **2007**, *111*, 9664 (Erratum: *J. Phys. Chem. B* **2008**, *112*, 1884)]. It may be noted that $\int [f(x)]^n dx \neq [\int f(x) dx]^n$. A similar definition in the case of the condensed Fukui function is justified (see ref 93) due to the fact that "the related parameter differentiation and integration do commute". Accordingly it is also valid for the condensed local softness and the philicity because in these cases a product of the Fukui function with a global property is considered. An approximate condensed-to-atom hardness may be constructed by integrating the local hardness over an atomic basin, provided the other points are taken into consideration (refs 68 and 69). See also: Toufar, H.; Nulens, K.; Janssens, G. O. A.; Mortier, W. J.; Schoonheydt, R.; De Proft, F.; Geerlings, P. *J. Phys. Chem.* **1996**, *100*, 15383.**
- (93) Yang, W.; Mortier, W. J. *J. Am. Chem. Soc.* **1986**, *108*, 5708.
- (94) Mendez, F.; Gázquez, J. L. *J. Am. Chem. Soc.* **1994**, *116*, 9298. See also Meza, R.; Gordillo, B.; Galván, M. *Int. J. Quantum Chem.* **2005**, *104*, 29.
- (95) (a) Gázquez, J. L.; Mendez, F. *J. Phys. Chem.* **1994**, *98*, 4591. (b) R. F. W. Bader, private communication.
- (96) Pérez, P.; Toro-Labbé, A.; Aizman, A.; Contreras, R. *J. Org. Chem.* **2002**, *67*, 4747.
- (97) Chamorro, E.; Chattaraj, P. K.; Fuentealba, P. *J. Phys. Chem. A* **2003**, *107*, 7068.
- (98) Chattaraj, P. K.; Maiti, B.; Sarkar, U. *J. Phys. Chem. A* **2003**, *107*, 4973.
- (99) Chattaraj, P. K. *J. Phys. Chem. A* **2001**, *105*, 511.
- (100) Melin, J.; Aparicio, F.; Subramanian, V.; Galván, M.; Chattaraj, P. K. *J. Phys. Chem. A* **2004**, *108*, 2487.
- (101) Hocquet, A.; Toro-Labbé, A.; Chermette, H. *THEOCHEM* **2004**, 686, 213.
- (102) Klopman, G. In *Chemical Reactivity and Reaction Paths*; Klopman, G., Ed.; Wiley: New York, 1974; p 1.
- (103) Pérez, P.; Aizman, A.; Contreras, R. *J. Phys. Chem. A* **2002**, *106*, 3964.
- (104) Legon, A. C.; Millen, D. J. *J. Am. Chem. Soc.* **1987**, *109*, 356.
- (105) Legon, A. C. *Angew. Chem., Int. Ed.* **1999**, *38*, 2686. The inverse behavior between electrophilicity and nucleophilicity is highlighted in some recent papers, e.g., Moncada, J. L.; Toro-Labbe, A. *Chem. Phys. Lett.* **2006**, *429*, 161. Ebrahimi, A.; Roohi, H.; Habibi, M.; Hasannejad, M. *Chem. Phys.* **2006**, *327*, 368. See also Jaramillo, P.; Fuentealba, P.; Perez, P. *Chem. Phys. Lett.* **2006**, *427*, 421. Jaramillo, P.; Perez, P.; Contreras, R.; Tiznado, W.; Fuentealba, P. *J. Phys. Chem. A* **2006**, *110*, 8181. Cedillo, A.; Contreras, R.; Galván, M.; Aizman, A.; Andres, J.; Safont, V. S. *J. Phys. Chem. A* **2007**, *111*, 2442. De Vleeschouwer, F.; Van Speybroeck, V.; Waroquier, M.; Geerlings, P.; De Proft, F. *Org. Lett.* **2007**, *9*, 2721.
- (106) Roy, R. K.; Usha, V.; Paulovici, J.; Hirao, K. *J. Phys. Chem. A* **2005**, *109*, 4601.
- (107) Roy, R. K. *J. Phys. Chem. A* **2004**, *108*, 4934. Although it was claimed in this paper that the local electrophilicity is a more reliable descriptor than its global counterpart, in two recent publications [Roy, R. K.; Usha, V.; Patel, B. K.; Hirao, K. *J. Comput. Chem.* **2006**, *27*, 773. Roy, R. K.; Bagaria, P.; Naik, S.; Kavala, V.; Patel, B. K. *J. Phys. Chem. A* **2006**, *110*, 2181], based on the numerical calculation on a different set of molecules, the authors have prescribed just the opposite. Where both fail, other factors such as steric effects are mentioned to be the plausible reasons. Even the analysis based on the Hirshfeld population scheme was questioned recently. [Bultinck, P.; Carbó-Dorca, R. *J. Phys. Chem. A* **2007**, *111*, 2640. Bultinck, P. *Faraday Discuss.* **2007**, *135*, 244] **Fukui function does become negative in some cases: Bultinck, P.; Dias, S.; Van Alsenoy, C.; Ayers, P. W.; Carbo-Dorca, R. *J. Chem. Phys.* **2007**, *127*, 034102. Ayers, P. W. *Phys. Chem. Chem. Phys.* **2006**, *8*, 3387. Melin, J.; Ayers, P. W.; Ortiz, J. V. *J. Phys. Chem. A* **2007**, *111*, 10017.** We feel that both global and local descriptors, other charge-based descriptors, and, if possible, other effects, such as steric, solvent, and entropy effects, are to be considered in order to have better insights into the chemical reactivity and selectivity. Of course, a few of them will dominate over the rest and some will be dependent on the others. We hope that it would be possible to identify the appropriate descriptor(s) for a given class of molecules/reactions prior to the computation and not after comparing the numerical results with the corresponding experimental trends.
- (108) Ingold, C. K. *J. Chem. Soc.* **1933**, 1120. Ingold, C. K. *Chem. Rev.* **1934**, *15*, 225.
- (109) Zhou, Z.; Parr, R. G. *J. Am. Chem. Soc.* **1990**, *112*, 5720.
- (110) Bader, R. F. W.; Chang, C. J. *Phys. Chem.* **1989**, *93*, 5095.
- (111) Kuhn, O.; Rau, D.; Mayr, H. *J. Am. Chem. Soc.* **1998**, *120*, 900. Gotta, M. F.; Mayr, H. *J. Org. Chem.* **1998**, *63*, 9769.
- (112) Mayr, H.; Muller, K.-H.; Ofial, A. R.; Buhl, M. *J. Am. Chem. Soc.* **1999**, *121*, 2418.
- (113) Lucius, R.; Mayr, H. *Angew. Chem., Int. Ed.* **2000**, *39*, 1995.
- (114) Mayr, H.; Bug, T.; Gotta, M. F.; Hering, N.; Irrgang, B.; Janker, B.; Kempf, B.; Loos, R.; Ofial, A. R.; Remennikov, G.; Schimmel, H. *J. Am. Chem. Soc.* **2001**, *123*, 9500.
- (115) Mayr, H.; Lang, G.; Ofial, A. R. *J. Am. Chem. Soc.* **2002**, *124*, 4076.
- (116) Schindele, C.; Houk, K. N.; Mayr, H. *J. Am. Chem. Soc.* **2002**, *124*, 11208.
- (117) Mayr, H.; Fichtner, C.; Ofial, A. R. *J. Chem. Soc., Perkin Trans.* **2002**, *2*, 1435.
- (118) Mayr, H.; Schimmel, H.; Kobayashi, S.; Kotani, M.; Prabakaran, T. R.; Sipos, L.; Faust, R. *Macromolecules* **2002**, *35*, 4611.
- (119) Bug, T.; Hartnagel, M.; Schlierf, C.; Mayr, H. *Chem.—Eur. J.* **2003**, *9*, 4068.
- (120) Ofial, A. R.; Ohkubo, K.; Fukuzumi, S.; Lucius, R.; Mayr, H. *J. Am. Chem. Soc.* **2003**, *125*, 10906.
- (121) Lemek, T.; Mayr, H. *J. Org. Chem.* **2003**, *68*, 6880.
- (122) Remennikov, G. Y.; Kempf, B.; Ofial, A. R.; Polborn, K.; Mayr, H. *J. Phys. Org. Chem.* **2003**, *16*, 431. Lakhdar, S.; Goumont, R.; Terrier, F.; Boubaker, T.; Dust, J. M.; Buncel, E. *Org. Biomol. Chem.* **2007**, *5*, 1744.
- (123) (a) Denegri, B.; Minegishi, S.; Kronja, O.; Mayr, H. *Angew. Chem., Int. Ed.* **2004**, *43*, 2302. (b) Denegri, B.; Streiter, A.; Jurić, S.; Ofial, A. R.; Kronja, O.; Mayr, H. *Chem.—Eur. J.* **2006**, *12*, 1648.
- (124) Tokuyasu, T.; Mayr, H. *Eur. J. Org. Chem.* **2004**, 2791.
- (125) Minegishi, S.; Loos, R.; Kobayashi, S.; Mayr, H. *J. Am. Chem. Soc.* **2005**, *127*, 2641.
- (126) Minegishi, S.; Mayr, H. *J. Am. Chem. Soc.* **2003**, *125*, 286.
- (127) Bug, T.; Mayr, H. *J. Am. Chem. Soc.* **2003**, *125*, 12980.
- (128) Ritchie, C. D. *Acc. Chem. Res.* **1972**, *5*, 348.
- (129) Ritchie, C. D. *J. Am. Chem. Soc.* **1975**, *97*, 1170.
- (130) Tishkov, A. A.; Mayr, H. *Angew. Chem., Int. Ed.* **2005**, *44*, 142.
- (131) Netz, A.; Müller, T. J. *J. Tetrahedron* **2000**, *56*, 4149.
- (132) Böttger, G. M.; Fröhlich, R.; Würthwein, E.-U. *Eur. J. Org. Chem.* **2000**, 1589.
- (133) Pérez, P. *J. Org. Chem.* **2003**, *68*, 5886.
- (134) Moss, R. A. *Acc. Chem. Res.* **1980**, *13*, 58.
- (135) Moss, R. A.; Pérez, L. A.; Wlostowska, J.; Guo, W.; Krogh-Jespersen, K. *J. Org. Chem.* **1982**, *47*, 4177.
- (136) Moss, R. A.; Kmiecik-Lawrynowicz, G.; Krogh-Jespersen, K. *J. Org. Chem.* **1986**, *51*, 2168.
- (137) Moss, R. A.; Wlostowski, M.; Terpinski, J.; Kmiecik-Lawrynowicz, G.; Krogh-Jespersen, K. *J. Am. Chem. Soc.* **1987**, *109*, 3811.
- (138) Moss, R. A.; Shen, S.; Hadel, L. M.; Kmiecik-Lawrynowicz, G.; Wlostowska, J.; Krogh-Jespersen, K. *J. Am. Chem. Soc.* **1987**, *109*, 4341.
- (139) Moss, R. A. *Acc. Chem. Res.* **1989**, *22*, 15.
- (140) Kötting, C.; Sander, W. *J. Am. Chem. Soc.* **1999**, *121*, 8891.
- (141) Sander, W.; Kötting, C. *Chem.—Eur. J.* **1999**, *5*, 24.
- (142) Sander, W.; Kötting, C.; Hübert, R. *J. Phys. Org. Chem.* **2000**, *13*, 561.
- (143) Strassner, T. *Top. Organomet. Chem.* **2004**, *13*, 1.
- (144) Roberts, D. W.; York, M.; Basketter, D. A. *Contact Dermatitis* **1999**, *41*, 14.
- (145) Roberts, D. W.; Patlewicz, G. *SAR QSAR Environ. Res.* **2002**, *13*, 145.

- (146) Pérez, P. J. *Phys. Chem. A* **2003**, *107*, 522.
- (147) Swain, C. G.; Scott, C. B. *J. Am. Chem. Soc.* **1953**, *75*, 141.
- (148) Jia, Z.; Salaita, M. G.; Margerum, D. W. *Inorg. Chem.* **2000**, *39*, 1974.
- (149) Maynard, A. T.; Covell, D. G. *J. Am. Chem. Soc.* **2001**, *123*, 1047.
- (150) Chattaraj, P. K.; Maiti, B. *J. Phys. Chem. A* **2001**, *105*, 169. Nucleophilicity as the multiplicative inverse of electrophilicity, as proposed in this paper, as well as a local variant of the same exhibit excellent correlations with the associated experimental results in the cases of several arenes and heteroarenes. Pratihar, S.; Roy, S. *J. Org. Chem.* **2010**, *75*, 4957.
- (151) Tielemans, M.; Areschkhia, V.; Colomer, J.; Promel, R.; Langenaeker, W.; Geerlings, P. *Tetrahedron* **1992**, *48*, 10575.
- (152) Olah, J.; De Proft, F.; Veszpremi, T.; Geerlings, P. *J. Phys. Chem. A* **2005**, *109*, 1608.
- (153) Szentpály, L. v. *Int. J. Quantum Chem.* **2000**, *76*, 222.
- (154) Gardner, D. O. N.; Szentpály, L. v. *J. Phys. Chem. A* **1999**, *103*, 9313.
- (155) Donald, K. J.; Mulder, W. H.; Szentpály, L. v. *J. Phys. Chem. A* **2004**, *108*, 595.
- (156) Szentpály, L. v.; Gardner, D. O. N. *J. Phys. Chem. A* **2001**, *105*, 9467. For the calculation of ω for a large system, see: Fan, W. J.; Zhang, R. Q.; Liu, S. *J. Comput. Chem.* **2007**, *28*, 967.
- (157) Campodonico, P.; Santos, J. G.; Andres, J.; Contreras, R. *J. Phys. Org. Chem.* **2004**, *17*, 273.
- (158) Ingold, C. K. *Structure and Mechanism in Organic Chemistry*; Cornell University Press: Ithaca, NY, 1953.
- (159) Contreras, R.; Andres, J.; Safont, V. S.; Campodonico, P.; Santos, J. G. *J. Phys. Chem. A* **2003**, *107*, 5588. See also: Meneses, L.; Tiznado, W.; Contreras, R.; Fuentealba, P. *Chem. Phys. Lett.* **2004**, *383*, 181. DELETED (Now ref 204d).
- (160) Neuvonen, H.; Neuvonen, K.; Koch, A.; Kleinpeter, E.; Pasanen, P. *J. Org. Chem.* **2002**, *67*, 6995.
- (161) Neuvonen, H.; Neuvonen, K. *J. Chem. Soc., Perkin Trans. 2* **1999**, 1497.
- (162) Epstein, D. M.; Meyerstein, D. *Inorg. Chem. Commun.* **2001**, *4*, 705.
- (163) Deubel, D. V.; Frenking, G.; Senn, H. M.; Sundermeyer, J. *Chem. Commun.* **2000**, *24*, 2469.
- (164) Cronin, M. T. D.; Manga, N.; Seward, J. R.; Sinks, G. D.; Schultz, T. W. *Chem. Res. Toxicol.* **2001**, *14*, 1498.
- (165) Ren, S. *Toxicol. Lett.* **2003**, *144*, 313. See also: Pasha, F. A.; Srivastava, H. K.; Singh, P. P. *Bioorg. Med. Chem.* **2005**, *13*, 6823.
- (166) Ren, S.; Schultz, T. W. *Toxicol. Lett.* **2002**, *129*, 151.
- (167) Lippa, K. A.; Roberts, A. L. *Environ. Sci. Technol.* **2002**, *36*, 2008.
- (168) Saethre, L. J.; Thomas, T. D.; Svensson, S. *J. Chem. Soc., Perkin Trans. 2* **1997**, 749.
- (169) Topol, I. A.; McGrath, C.; Chertova, E.; Dasenbrock, C.; Lacourse, W. R.; Eissenstat, M. A.; Burt, S. K.; Henderson, L. E.; Casas-Finet, J. R. *Protein Sci.* **2001**, *10*, 1434. See also: Moens, J.; Geerlings, P.; Roos, G. *Chem.—Eur. J.* **2007**, *13*, 8174. Moens, J.; Roos, G.; Jaque, P.; De Proft, F.; Geerlings, P. *Chem.—Eur. J.* **2007**, *13*, 9331. Moens, J.; Jaque, P.; De Proft, F.; Geerlings, P. *J. Phys. Chem. A* **2008**, *112*, 6023.
- (170) Morris, S. J.; Thurston, D. E.; Nevell, T. G. *J. Antibiot.* **1990**, *43*, 1286.
- (171) Dronskowski, R.; Hoffmann, R. *Adv. Mater.* **1992**, *4*, 514.
- (172) Benigni, R.; Cotta-Ramusino, M.; Andreoli, C.; Giuliani, A. *Carcinogenesis* **1992**, *13*, 547.
- (173) Mekenyan, O. G.; Veith, G. D. *SAR QSAR Environ. Res.* **1993**, *1*, 335.
- (174) Mrozek, A.; Karolak-Wojciechowska, J.; Amiel, P.; Barbe, J. *THEOCHEM* **2000**, *524*, 159.
- (175) Jaramillo, P.; Pérez, P.; Fuentealba, P. *J. Phys. Chem. A* **2009**, *113*, 6812.
- (176) Jaramillo, P.; Domingo, L. R.; Chamorro, E.; Pérez, P. *THEOCHEM* **2008**, *865*, 68.
- (177) Campodonico, P. R.; Aizman, A.; Contreras, R. *Chem. Phys. Lett.* **2006**, *422*, 204.
- (178) Domingo, L. R.; Aurell, M. J.; Arnó, M. *Tetrahedron* **2009**, *65*, 3432.
- (179) Chamorro, E.; Duque-Noreña, M.; Peñez, P. *THEOCHEM* **2009**, *896*, 73.
- (180) Ghomri, A.; Mekelleche, S. M. *THEOCHEM* **2010**, *941*, 36.
- (181) Broeckaert, L.; Moens, J.; Roos, G.; De Proft, F.; Geerlings, P. *J. Phys. Chem. A* **2008**, *112*, 12164.
- (182) Anderson, J. S. M.; Liu, Y.; Thomson, J. W.; Ayers, P. W. *THEOCHEM* **2010**, *943*, 168.
- (183) Liu, S.; Li, T.; Ayers, P. W. *J. Chem. Phys.* **2009**, *131*, 114106.
- (184) Ayers, P. W.; Liu, S.; Li, T. *Chem. Phys. Lett.* **2009**, *480*, 318.
- (185) Chattaraj, P. K.; Chakraborty, A.; Giri, S. *J. Phys. Chem. A* **2009**, *113*, 10068. Feasibility of an electrophile–nucleophile interaction is studied (Chattaraj, P. K.; Chakraborty, A.; Giri, S. unpublished result) in terms of the electroaccepting power of the electrophile and the electrodonating power of the nucleophile.
- (186) Fuentealba, P.; Contreras, R. In *Reviews of Modern Quantum Chemistry*; Sen, K. D., Ed.; World Scientific: Singapore, 2002; pp 1013–1052.
- (187) Langenaeker, W.; Demel, K.; Geerlings, P. *THEOCHEM* **1991**, *234*, 329.
- (188) Russo, N.; Toscano, M.; Grand, A.; Mineva, T. *J. Phys. Chem. A* **2000**, *104*, 4017.
- (189) Chattaraj, P. K.; Gonzalez-Rivas, N.; Matus, M. H.; Galván, M. J. *Phys. Chem. A* **2005**, *109*, 5602. (a) See also: Lamsabhi, A. M.; Escobar, C. A.; Pérez, P. *J. Phys. Org. Chem.* **2005**, *18*, 1161. Senthilkumar, K.; Raman, M. S.; Kolandaivel, P. *THEOCHEM* **2006**, *758*, 107. Moraleda, D.; El Abed, D.; Pellissier, H.; Santelli, M. *THEOCHEM* **2006**, *760*, 113.
- (190) Kato, S. *Theor. Chem. Acc.* **2000**, *103*, 219.
- (191) Olah, J.; Van Alsenoy, C.; Sannigrahi, A. B. *J. Phys. Chem. A* **2002**, *106*, 3885.
- (192) Chatterjee, A.; Ebina, T.; Iwasaki, T. *J. Phys. Chem. A* **2001**, *105*, 10694.
- (193) Ireta, J.; Galván, M.; Kyeonjae, C.; Joannopoulos, J. D. *J. Am. Chem. Soc.* **1998**, *120*, 9771.
- (194) Korchowiec, J. *Comput. Chem.* **2000**, *24*, 259. Korchowiec, J. *Int. J. Quantum Chem.* **2005**, *101*, 714.
- (195) Nalewajski, R. F. *J. Am. Chem. Soc.* **1984**, *106*, 944. Nalewajski, R. F.; Korchowiec, J. *Charge Sensitivity Approach to Electronic Structure and Chemical Reactivity*; World Scientific: Singapore, 1997.
- (196) Roy, R. K.; Krishnamurti, S.; Geerlings, P.; Pal, S. *J. Phys. Chem. A* **1998**, *102*, 3746. See also: De, H. S.; Krishnamurti, S.; Pal, S. *J. Phys. Chem. C* **2009**, *113*, 7101; **2010**, *114*, 6690.
- (197) Chandrakumar, K. R. S.; Pal, S. *Int. J. Mol. Sci.* **2002**, *3*, 324, and references therein.
- (198) Tanwar, A.; Pal, S. *J. Phys. Chem. A* **2004**, *108*, 11838.
- (199) Domingo, L. R.; Aurell, M. J.; Pérez, P.; Contreras, R. *J. Phys. Chem. A* **2002**, *106*, 6871. and many other papers from this group. See also: Elango, M.; Parthasarathi, R.; Subramanian, V.; Chattaraj, P. K. *Int. J. Quantum Chem.* **2006**, *106*, 852. Roy, D. R.; Parthasarathi, R.; Padmanabhan, J.; Sarkar, U.; Subramanian, V.; Chattaraj, P. K. *J. Phys. Chem. A* **2006**, *110*, 1084.
- (200) Chandra, A. K.; Nguyen, M. T. *J. Chem. Soc., Perkin Trans. 2* **1997**, 1415. Damoun, S.; Van de Woude, G.; Méndez, F.; Geerlings, P. *J. Phys. Chem. A* **1997**, *101*, 886. Chandra, A. K.; Uchimaru, T.; Nguyen, M. T. *J. Chem. Soc., Perkin Trans. 2* **1999**, 2117.
- (201) Dal Pino, A., Jr.; Galván, M.; Arias, T. A.; Joannopoulos, J. D. *J. Chem. Phys.* **1993**, *98*, 1606.
- (202) Zhang, Y. L.; Yang, Z. Z. *THEOCHEM* **2000**, *496*, 139.
- (203) Espinosa, A.; Frontera, A.; García, R.; Soler, M. A.; Tárraga, A. *ARKIVOC* **2005**, *ix*, 415.
- (204) Clark, L. A.; Ellis, D. E.; Snurr, R. Q. *J. Chem. Phys.* **2001**, *114*, 2580.
- (205) (a) Morell, C.; Grand, A.; Toro-Labbé, A. *J. Phys. Chem. A* **2005**, *109*, 205. (b) Roy, D. R.; Subramanian, V.; Chattaraj, P. K. *Los Alamos Natl. Lab., Prepr. Arch., Condens. Matter* **2005**, 1–21, arXiv: cond-mat/0511441. Note that $\Delta\omega$ is a better intermolecular descriptor than Δf because of the global information in the former. A portion of this work was presented at the 231st annual meeting of the American Chemical Society in Atlanta, GA. (c) Roy, D. R.; Subramanian, V.; Chattaraj, P. K. *Ind. J. Chem. A* **2006**, *45A*, 2369. (d) Morell, C.; Grand, A.; Toro-Labbe, A. *Chem. Phys. Lett.* **2006**, *425*, 342. (e) It has been argued that $\Delta\omega(r)$ may be considered to be a measure of the difference between local and global (modulated by $f(r)$) reactivity variations when an electron is added or removed. Padmanabhan, J.; Parthasarathi, R.; Elango, M.; Subramanian, V.; Krishnamoorthy, B. S.; Gutierrez-Oliva, S.; Toro-Labbé, A.; Roy, D. R.; Chattaraj, P. K. *J. Phys. Chem. A* **2007**, *111*, 9130. (f) Size extensive nature of these descriptors or lack of it is being analyzed [Ayers, P. W.; Chattaraj, P. K. Unpublished results].
- (206) Geerlings, P.; Boon, G.; Van Alsenoy, C.; De Proft, F. *Int. J. Quantum Chem.* **2005**, *101*, 722.
- (207) Li, Y.; Evans, J. N. S. *J. Am. Chem. Soc.* **1995**, *117*, 7756.
- (208) (a) Li, Y.; Evans, J. N. S. *Proc. Natl. Acad. Sci. U.S.A.* **1996**, *93*, 4612. (b) Anderson, J. S. M.; Melin, J.; Ayers, P. W. *J. Chem. Theo. Comp.* **2007**, *3*, 358. (c) Anderson, J. S. M.; Melin, J.; Ayers, P. W. *J. Chem. Theo. Comp.* **2007**, *3*, 375. (d) Anderson, J. S. M.; Ayers, P. W. *Phys. Chem. Chem. Phys.* **2007**, *9*, 2371.
- (209) Boiani, M.; Cerecetto, M.; Gonzalez, M.; Piro, O. E.; Castellano, E. E. *J. Phys. Chem. A* **2004**, *108*, 11241.
- (210) Ritchie, J. P. *THEOCHEM* **1992**, *225*, 297.
- (211) Pollack, S. K.; Devlin, J. L.; Summerhays, K. D.; Taft, R. W.; Hehre, W. J. *J. Am. Chem. Soc.* **1977**, *99*, 4583.
- (212) Politzer, P.; Murray, J. S.; Concha, M. C. *Int. J. Quantum Chem.* **2002**, *88*, 19.

- (213) Sjöberg, P.; Murray, J. S.; Brinck, T.; Politzer, P. *Can. J. Chem.* **1990**, *68*, 1440.
- (214) Pérez, P.; Contreras, R.; Aizman, A. *Chem. Phys. Lett.* **1996**, *260*, 236.
- (215) Frison, G.; Mathey, F.; Sevin, A. *J. Organomet. Chem.* **1998**, *570*, 225.
- (216) Frison, G.; Mathey, F.; Sevin, A. *J. Phys. Chem. A* **2002**, *106*, 5653.
- (217) Cohen, M. H.; Ganduglia-Pirovano, M. V.; Kudrnovsky, J. *J. Chem. Phys.* **1994**, *101*, 8988.
- (218) Balawender, R.; Geerlings, P. *J. Chem. Phys.* **2001**, *114*, 682.
- (219) De Proft, F.; Liu, S.; Geerlings, P. *J. Chem. Phys.* **1998**, *108*, 7549.
- (220) (a) Ayers, P. W.; Anderson, J. S. M.; Rodríguez, J. I.; Jawed, Z. *Phys. Chem. Chem. Phys.* **2005**, *7*, 1918. (b) Jaramillo, P.; Domingo, L. R.; Pérez, P. *Chem. Phys. Lett.* **2006**, *420*, 95. (c) Roos, G.; Loverix, S.; Brosens, E.; Van Belle, K.; Wyns, L.; Geerlings, P.; Messens, J. *J. Chem. Bio. Chem.* **2006**, *7*, 981. (d) Campodonico, P. R.; Aizman, A.; Contreras, R. *Chem. Phys. Lett.* **2006**, *422*, 340. (e) Shaiyanyan, B. A. *Int. J. Quantum Chem.* **2007**, *107*, 189. (f) Giri, S.; Chattaraj, P. K. Unpublished work. (g) Campodonico, P. R.; Andres, J.; Aizman, A.; Contreras, R. *Chem. Phys. Lett.* **2007**, *439*, 177.
- (221) Deuri, S.; Phukan, P. *THEOCHEM* **2010**, *945*, 64. (b) Deuri, S.; Phukan, P. *Ind. J. Chem.* **2010**, *49A*, 1206.
- (222) Pefez, P.; Domingo, L. R.; Duque-Noreña, M.; Chamorro, E. *THEOCHEM* **2009**, *895*, 86.
- (223) Cárdenas, C.; Rabi, N.; Ayers, P. W.; Morell, C.; Jaramillo, P.; Fuentealba, P. *J. Phys. Chem. A* **2009**, *113*, 8660.
- (224) Giri, S.; Wang, D. Z.; Chattaraj, P. K. *Tetrahedron* **2010**, *66*, 4560.
- (225) (a) Huheey, J. *Inorganic Chemistry: Principles of Structure and Reactivity*, 4th ed.; Harper and Row: 1993. (b) *Inorganic Chemistry, Part-I Principles*; Dutta, R. L., Ed.; The New Book Stall: Kolkata, 2003.
- (226) Chattaraj, P. K.; Maiti, B. *J. Chem. Educ.* **2001**, *78*, 811.
- (227) Chamorro, E.; Chattaraj, P. K.; Fuentealba, P. *J. Phys. Chem. A* **2003**, *107*, 7068. The domain of validity of the minimum electrophilicity principle was discussed in this paper. For the validity of this principle in the case of the chemical reactions, see: ref 52d, and for the reactivity of [n]paracyclophanes using this principle, see: *THEOCHEM* **2007**, *820*, 1. See also: Noorizadeh, S. *J. Phys. Org. Chem.* **2007**, *20*, 514. For an exception, see: Correa, J. V.; Herrera, B.; Toro-Labbé, A. *THEOCHEM* **2007**, *817*, 111. A connection between this principle and the HSAB principle is available in Chattaraj, P. K.; Giri, S. *Ind. J. Phys.* **2007**, *81*, 871. **2008**, *82*, 467.
- (228) Fuentealba, P.; Parr, R. G. *J. Chem. Phys.* **1991**, *94*, 5559.
- (229) Gunnarson, O.; Lundqvist, B. I. *Phys. Rev. B* **1976**, *13*, 4274.
- (230) Ziegler, T.; Rauk, A.; Baerends, E. J. *Theor. Chim. Acta* **1977**, *43*, 261.
- (231) von Barth, U. *Phys. Rev. A* **1979**, *20*, 1693.
- (232) Theophilou, A. *J. Phys. C: Solid State Phys.* **1979**, *12*, 5419.
- (233) Hadjisavvas, N.; Theophilou, A. *Phys. Rev. A* **1985**, *32*, 720.
- (234) Kohn, W. *Phys. Rev. A* **1986**, *34*, 5419.
- (235) Gross, E. K. U.; Oliveira, L. N.; Kohn, W. *Phys. Rev. A* **1988**, *37*, 2805. Oliveira, L. N.; Gross, E. K. U.; Kohn, W. *Phys. Rev. A* **1988**, *37*, 2821.
- (236) Ayers, P. W. Private communication. Excited-state reactivity may be analyzed through Levy–Nagy [Levy, M.; Nagy, A. *Phys. Rev. Lett.* **1999**, *83*, 4361. Jor Görling [Göring, A. *Phys. Rev. A* **1993**, *47*, 2783] functionals.
- (237) Chattaraj, P. K.; Poddar, A. *J. Phys. Chem. A* **1998**, *102*, 9944.
- (238) Chattaraj, P. K.; Poddar, A. *J. Phys. Chem. A* **1999**, *103*, 1274.
- (239) Chattaraj, P. K.; Poddar, A. *J. Phys. Chem. A* **1999**, *103*, 8691.
- (240) Fuentealba, P.; Simon-Manso, Y.; Chattaraj, P. K. *J. Phys. Chem. A* **2000**, *104*, 3185.
- (241) Chattaraj, P. K. Nonlinear Chemical Dynamics. In *Symmetries and Singularity Structures: Integrability and Chaos in Nonlinear Dynamical Systems*; Lakshmanan, M., Daniel, M., Eds.; Springer-Verlag: Berlin, 1990; pp 172–182.
- (242) Chattaraj, P. K.; Sengupta, S.; Poddar, A. *Int. J. Quantum Chem.* **1998**, *69*, 279.
- (243) Chattaraj, P. K.; Maiti, B. *J. Phys. Chem. A* **2004**, *108*, 658.
- (244) Chattaraj, P. K.; Sengupta, S.; Poddar, A. *THEOCHEM* **2000**, *501*, 339.
- (245) Chattaraj, P. K.; Sarkar, U. Chemical Reactivity Dynamics in Ground and Excited Electronic States. In *Theoretical and Computational Chemistry, Vol. 12, Theoretical Aspects of Chemical Reactivity*; Toro-Labbé, A., Eds.; Elsevier: Netherlands, 2007; Chapter 13, pp 269–286.
- (246) Chattaraj, P. K.; Sengupta, S. *J. Phys. Chem. A* **1999**, *103*, 6122.
- (247) (a) Chattaraj, P. K.; Maiti, B. *Int. J. Mol. Sci.* **2002**, *3*, 338. (b) Sarkar, U.; Chattaraj, P. K. *Ind. J. Phys.* **2006**, *80*, 463.
- (248) Chattaraj, P. K.; Sarkar, U. *Int. J. Quantum Chem.* **2003**, *91*, 633.
- (249) Michels, A.; de Boer, J.; Bijl, A. *Physica* **1937**, *4*, 981.
- (250) Chattaraj, P. K.; Sarkar, U. *J. Phys. Chem. A* **2003**, *107*, 4877. See also: Sarkar, U.; Giri, S.; Chattaraj, P. K. *J. Phys. Chem. A* **2009**, *113*, 10759, for the effect of using different formulas vis-a-vis the use of Dirichlet boundary condition and the cut-off function method.
- (251) Chattaraj, P. K.; Sarkar, U. *Chem. Phys. Lett.* **2003**, *372*, 805.
- (252) Chattaraj, P. K.; Maiti, B.; Sarkar, U. *Proc. Indian Acad. Sci. (Chem. Sci.)* **2003**, *115*, 195.
- (253) Tsipis, A. C.; Chaviara, A. T. *Inorg. Chem.* **2004**, *43*, 1273.
- (254) Contreras, R.; Andrés, J.; Domingo, L. R.; Castillo, R.; Pérez, P. *Tetrahedron* **2005**, *61*, 417. See also: Castro, E. A.; Campodonico, P. R.; Contreras, R.; Fuentealba, P.; Santos, J. G.; Leis, J. R.; Garcia-Rio, L.; Saez, J. A.; Domingo, L. R. *Tetrahedron* **2006**, *62*, 2555. DELETED (Now ref 287c).
- (255) Valencia, F.; Romero, A. H.; Kiwi, M.; Ramírez, R.; Toro-Labbé, A. *Chem. Phys. Lett.* **2003**, *372*, 815.
- (256) Kostova, I.; Manolov, I.; Nicolova, I.; Danchev, N. *II Farmaco* **2001**, *56*, 707.
- (257) Cases, M.; Frenking, G.; Duran, M.; Sola, M. *Organometallics* **2002**, *21*, 4182.
- (258) Jaque, P.; Toro-Labbé, A. *J. Chem. Phys.* **2002**, *117*, 3208. Jaque, P.; Toro-Labbé, A. *J. Phys. Chem. B* **2004**, *108*, 2568.
- (259) Cuan, A.; Galván, M.; Chattaraj, P. K. *J. Chem. Sci.* **2005**, *117*, 541. Gomez, B.; Martinez-Magadan, J. M.; Sarkar, U.; Chattaraj, P. K. *THEOCHEM* **2005**, *755*, 99.
- (260) Chattaraj, P. K.; Sarkar, U.; Parthasarathi, R.; Subramanian, V. *Int. J. Quantum Chem.* **2005**, *101*, 690.
- (261) Thanikaivelan, P.; Subramanian, V.; Rao, J. R.; Nair, B. U. *Chem. Phys. Lett.* **2000**, *323*, 59.
- (262) Cárdenas-Jirón, G. I. *Int. J. Quantum Chem.* **2003**, *91*, 389. See also: Frenking, G.; Sola, M.; Vyboishchikov, S. F. *J. Organomet. Chem.* **2005**, *690*, 6178. Costas, M.; Ribas, X.; Poater, A.; Lopez Valbuena, J. M.; Xifra, R.; Company, A.; Duran, M.; Sola, M.; Llobet, A.; Corbella, M.; Uson, M. A.; Mahia, J.; Solans, X.; Shan, X.; Benet-Buchholz, J. *Inorg. Chem.* **2006**, *45*, 3569.
- (263) Belmer, J.; Zuñiga, C.; Jiménez, C.; Saavedra, J. *Bol. Soc. Chil. Quím.* **2002**, *47*, 371.
- (264) Jiménez, C. A.; Belmer, J. B. *Tetrahedron* **2005**, *61*, 3933.
- (265) Frantz, S.; Hartmann, H.; Doslik, N.; Wanner, M.; Kaim, W.; Kummerer, H.-J.; Denninger, G.; Barra, A.-L.; Duboc-Toia, C.; Fiedler, J.; Ciofini, I.; Urban, C.; Kaupp, M. *J. Am. Chem. Soc.* **2002**, *124*, 10563.
- (266) Kostova, I.; Trendafilova, N.; Mihaylov, T. *Chem. Phys.* **2005**, *314*, 73.
- (267) Parthasarathi, R.; Padmanabhan, J.; Elango, M.; Subramanian, V.; Chattaraj, P. K. *Chem. Phys. Lett.* **2004**, *394*, 225. For a correlation between the group philicity and pK_a values, see: Parthasarathi, R.; Padmanabhan, J.; Elango, M.; Chitra, K.; Subramanian, V.; Chattaraj, P. K. *J. Phys. Chem. A* **2006**, *110*, 6540. A comparison between Brønsted and Lewis acidities is also available in Gupta, K.; Roy, D. R.; Subramanian, V.; Chattaraj, P. K. *THEOCHEM* **2007**, *812*, 13.
- (268) Tsipis, C. A.; Karagiannis, E. E.; Kladou, P. F.; Tsipis, A. C. *J. Am. Chem. Soc.* **2004**, *126*, 12916. Tsipis, A. C.; Stalikas, A. V. *New J. Chem.* **2007**, *31*, 852.
- (269) Tsipis, A. C.; Tsipis, C. A. *J. Am. Chem. Soc.* **2003**, *125*, 1136.
- (270) Merino, G.; Mendez-Rojas, M. A.; Vela, A. *J. Am. Chem. Soc.* **2003**, *125*, 6026. Merino, G.; Mendez-Rojas, M. A.; Beltran, H. I.; Corminboeuf, C.; Heine, T.; Vela, A. *J. Am. Chem. Soc.* **2004**, *126*, 16160.
- (271) Yang, X.; Hajgato, B.; Yamada, M.; Ohno, K. *Chem. Lett.* **2005**, *34*, 506.
- (272) Yang, Y. J.; Su, Z. M. *Int. J. Quantum Chem.* **2005**, *103*, 54. See also: Elango, M.; Subramanian, V. *J. Phys. Chem. A* **2005**, *109*, 11870.
- (273) Elango, M.; Parthasarathi, R.; Narayanan, G. K.; Sabeelullah, A. M.; Sarkar, U.; Venkatasubramanian, N. S.; Subramanian, V.; Chattaraj, P. K. *J. Chem. Sci.* **2005**, *117*, 61. See also: Chattaraj, P. K.; Roy, D. R.; Elango, M.; Subramanian, V. *J. Phys. Chem. A* **2005**, *109*, 9590. In this paper, the aromaticity/antiaromaticity of various molecules, such as Al₄²⁻, Al₄⁴⁻, MAI₄⁻ [M = Li, Na, K, Cu], Li₃Al₄⁻, Li₄Al₄, and their sandwich complexes, is analyzed. Chattaraj, P. K.; Sarkar, U.; Roy, D. R. *J. Chem. Educ.* **2007**, *84*, 354. For connection with the Hammett constant, see: Galabov, B.; Ilieva, S.; Schaefer, H. F., III *J. Org. Chem.* **2006**, *71*, 6382. For aromaticity in polyacene analogues of Na₆, see: Khatua, S.; Roy, D. R.; Chattaraj, P. K.; Bhattacharjee, M. *Chem. Commun.* **2007**, *2*, 135. For that in BN⁻, CN⁻, AlN⁻, BO⁻, BS-acenes, see: Chattaraj, P. K.; Roy, D. R. *J. Phys. Chem. A* **2007**, *111*, 4684. For that in Al₇C⁻ and Al₇O⁻, see: Chattaraj, P. K.; Giri, S. *J. Phys. Chem. A* **2007**, *111*, 11116. For that in M₂Be₃ [M = Li, Na, Cu] as well as ZnBe₃, Cp₂Zn, [Zn₂Be₆]²⁻, Cp≡C₃H₅⁻ see: Roy, D. R.; Chattaraj, P. K. *J. Phys. Chem. A* **2008**, *112*, 1612. Giri, S.; Roy, D. R.; Duley, S.; Parthasarathi, R.; Elango, M.; Vijayraj, R.; Subramanian, V.; Islas, R.; Merino, G.; Chattaraj, P. K. *J. Comput. Chem.* **2010**, *31*, 1815. Chattaraj, P. K., Ed. *Aromaticity and Metal Clusters*; Taylor & Francis/CRC Press: Boca Raton, FL, 2010 (in press). Chakraborty, A.; Giri, S.; Chattaraj,

- P. K. *New J. Chem.* **2010**, *34*, 1936. Duley, S.; Chakraborty, A.; Giri, S.; Chattaraj, P. K. *J. Sulf. Chem.* **2010**, *31*, 231. Giri, S.; Chakraborty, A.; Chattaraj, P. K. *J. Mol. Model.* **2010**, DOI: 10.1007/s00894-010-0761-1. In most of these papers, a minimum electrophilicity principle has been shown to be valid. Bond-stretch isomerism in these clusters and the possibility of their use as hydrogen storage material have been studied.
- (274) Vianello, R.; Liebman, J. F.; Maksic, Z. B. *Chem.—Eur. J.* **2004**, *10*, 5751. For superbases, see: Chamorro, E.; Escobar, C. A.; Sienra, R.; Pérez, P. *J. Phys. Chem. A* **2005**, *109*, 10068.
- (275) Pérez, P. *J. Org. Chem.* **2004**, *69*, 5048.
- (276) Koltunov, K. Y.; Prakash, G. K. S.; Rasul, G.; Olah, G. A. *J. Org. Chem.* **2002**, *67*, 8943.
- (277) Esteves, P. M.; Ramirez-Solis, A.; Mota, C. J. A. *J. Am. Chem. Soc.* **2002**, *124*, 2672.
- (278) Ren, J.; Cramer, C. J.; Squires, R. R. *J. Am. Chem. Soc.* **1999**, *121*, 2633.
- (279) Klumpp, D. A.; Lau, S. J. *J. Org. Chem.* **1999**, *64*, 7309.
- (280) Schleyer, P. v. R.; Maerker, C.; Dransfeld, A.; Jiao, H.; Hommes, N. J. R. v. E. *J. Am. Chem. Soc.* **1996**, *118*, 6317.
- (281) (a) Khatua, S.; Roy, D. R.; Chattaraj, P. K.; Bhattacharjee, M. *Chem. Commun.* **2007**. (b) Khatua, S.; Roy, D. R.; Bhattacharjee, M.; Chattaraj, P. K. *J. Comp. Meth. Sci. Eng.* **2008**, *7*, 395 (Special Issue on Alkali Clusters). (c) Khatua, S.; Roy, D. R.; Bultinck, P.; Bhattacharjee, M.; Chattaraj, P. K. *Phys. Chem. Chem. Phys.* **2008**, *10*, 2461. (d) For the Li₆ system, see: Deb, D.; Giri, S.; Chattaraj, P. K.; Bhattacharjee, M. *J. Phys. Chem. A* **2010**, *114*, 10871.
- (282) Roy, D. R.; Chattaraj, P. K. *J. Phys. Chem. A* **2008**, *112*, 1612.
- (283) (a) Chattaraj, P. K.; Giri, S. *THEOCHEM* **2008**, *865*, 53. (b) Duley, S.; Goyal, P.; Giri, S.; Chattaraj, P. K. *Croat. Chem. Acta* **2009**, *82*, 193 (Special Issue on Professor Zvonimir Maksic's 70th Birthday). (c) Duley, S.; Giri, S.; Chakraborty, A.; Chattaraj, P. K. *J. Chem. Sci.* **2009**, *121*, 849 (S. K. Rangarajan Special Issue).
- (284) Chakraborty, A.; Giri, S.; Chattaraj, P. K. *THEOCHEM* **2009**, *913*, 70.
- (285) Resa, I.; Carmona, E.; Gutierrez-Puebla, E.; Monge, A. *Science* **2004**, *305*, 1136.
- (286) Chattaraj, P. K.; Roy, D. R.; Duley, S. *Chem. Phys. Lett.* **2008**, *460*, 382.
- (287) Roy, D. R.; Duley, S.; Chattaraj, P. K. *Proc. Ind. Natl. Sci. Acad. A* **2008**, *74*, 11.
- (288) Chattaraj, P. K.; Giri, S. *Int. J. Quantum Chem.* **2009**, *109*, 2373 (Special Issue on Professor Istvan Mayer's 65th birthday).
- (289) Chattaraj, P. K.; Duley, S.; Das, R. *Chem. Educ.* **2010** (in press).
- (290) Chang, C. M. *J. Phys. Chem. A* **2008**, *112*, 2482.
- (291) Fayet, G.; Jacquemin, D.; Wathelet, V.; Perpète, E. A.; Rotureau, P.; Adamo, C. *J. Mol. Graphics Modell.* **2010**, *28*, 465.
- (292) Flores-Holguín, N.; Aguilar-Elguézabal, A.; Rodríguez-Valdez, L. M.; Glossman-Mitnik, D. *THEOCHEM* **2008**, *854*, 81.
- (293) Núñez-Zarur, F.; Vivas-Reyes, R. *THEOCHEM* **2008**, *850*, 127.
- (294) Vivas-Reyes, R.; Núñez-Zarur, F.; Martínez, E. *Org. Elect.* **2008**, *9*, 625.
- (295) Liu, S.; Govind, N. *J. Phys. Chem. A* **2008**, *112*, 6690.
- (296) Norambuena, E.; Olea-Azar, C.; Delgadillo, A.; Barrera, M.; Loeb, B. *Chem. Phys. Lett.* **2009**, *359*, 92.
- (297) Tang, C.; Weihua, Z.; Deng, K. *THEOCHEM* **2009**, *909*, 43.
- (298) Uğur, İ.; De Vleeschouwer, F.; Tüzün, N.; Aviyente, V.; Geerlings, P.; Liu, S.; Ayers, P. W.; De Proft, F. *J. Phys. Chem. A* **2009**, *113*, 8704.
- (299) Kassaei, M. Z.; Shakib, F. A.; Momeni, M. R.; Ghambarian, M.; Musavi, S. M. *Tetrahedron* **2009**, *65*, 10093.
- (300) Labet, V.; Morell, C.; Douki, T.; Cadet, J.; Eriksson, L. A.; Grand, A. *J. Phys. Chem. A* **2010**, *114*, 1826.
- (301) Kassaei, M. Z.; Momeni, M. R.; Shakib, F. A.; Ghambarian, M. *J. Org. Chem.* **2010**, *695*, 760.
- (302) Vleeschouwer, F. D.; Speybroeck, V. V.; Waroquier, M.; Geerlings, P.; Proft, F. D. *J. Org. Chem.* **2008**, *73*, 9109.
- (303) Sánchez-Bojorge, N.-A.; Flores-Holguín, N.; Glossman-Mitnik, D.; Rodríguez-Valdez, L. M. *THEOCHEM* **2009**, *912*, 119.
- (304) Fayet, G.; Joubert, L.; Rotureau, P.; Adamo, C. *Chem. Phys. Lett.* **2009**, *467*, 407.
- (305) Fayet, G.; Rotureau, P.; Joubert, L.; Adamo, C. *J. Hazard. Mater.* **2009**, *171*, 845.
- (306) Ibrahim, A. A.; Abdalrazaq, E. A. *Am. J. Appl. Sci.* **2009**, *6*, 1385.
- (307) Vektariene, A.; Vektaris, G.; Svoboda, J. *ARKIVOC* **2009**, 311.
- (308) Campodónico, P. R.; Aizman, A.; Contreras, R. *Chem. Phys. Lett.* **2009**, *471*, 168.
- (309) Kassaei, M. Z.; Zandi, H.; Momeni, M. R.; Shakib, F. A.; Ghambarian, M. *THEOCHEM* **2009**, *913*, 16.
- (310) Tsipis, A. C. *Organometallics* **2010**, *29*, 354.
- (311) Vázquez-Lima, H.; Guadarrama, P.; Ramos, E.; Fomine, S. *J. Mol. Catal. A: Chem.* **2009**, *310*, 75.
- (312) Ortiz, E. V.; Lopez, M. B.; Castro, C. A. *J. Phys.: Conf. Ser.* **2009**, *167*, 012021.
- (313) Martínez, J.; Cruz, V.; Ramos, J.; Gutiérrez-Oliva, S.; Martínez-Salazar, J.; Toro-Labbe, A. *J. Phys. Chem. C* **2008**, *112*, 5023.
- (314) Noorizadeh, S.; Shakerzadeh, E. *J. Phys. Chem. A* **2008**, *112*, 2486.
- (315) Padmanabhan, J.; Parthasarathi, R.; Subramanian, V.; Chattaraj, P. K. *J. Phys. Chem. A* **2007**, *111*, 1358. An electrophilicity based charge transfer descriptor is prescribed in this paper.
- (316) Da Silva Miranda, F.; Signori, A. M.; Vicente, J.; de Souza, B.; Priebe, J. P.; Szpoganicz, B.; Goncalves, N. S.; Neves, A. *Tetrahedron* **2008**, *64*, 5410.
- (317) Sakurai, Y.; Shiozaki, H.; Yokoyama, M. *THEOCHEM* **2006**, *766*, 41.
- (318) Azzouzi, S.; El Messaoudi, M.; Esseffar, M.; Jalal, R.; Cano, F. H.; Aprea-Rojas, M. D. C.; Domingo, L. R. *J. Phys. Org. Chem.* **2005**, *18*, 522. See also: Zhang, X.; Du, H.; Wang, Z.; Wu, Y.-D.; Ding, K. *J. Org. Chem.* **2006**, *71*, 2862. Domingo, L. R.; Pérez, P.; Contreras, R. *Eur. J. Org. Chem.* **2006**, 498. Diev, V. V.; Kostikov, R. R.; Gleiter, R.; Molchanov, A. P. *J. Org. Chem.* **2006**, *71*, 4066.
- (319) García Ruano, J. L.; Fraile, A.; Gonzalez, G.; Martin, M. R.; Clemente, F. R.; Gordillo, R. *J. Org. Chem.* **2003**, *68*, 6522.
- (320) Merino, P.; Revuelta, J.; Tejero, T.; Chiacchio, U.; Rescifina, A.; Romeo, G. *Tetrahedron* **2003**, *59*, 3581.
- (321) Peréz, P.; Domingo, L. R.; Aurell, M. J.; Contreras, R. *Tetrahedron* **2003**, *59*, 3117.
- (322) Sáez, J. A.; Arnó, M.; Domingo, L. R. *Tetrahedron* **2003**, *59*, 9167. Benchouk, W.; Mekelleche, S. M. *THEOCHEM* **2008**, *852*, 46. Singh, L.; Ishar, M. P. S.; Elango, M.; Subramanian, V.; Gupta, V.; Kanwal, P. *J. Org. Chem.* **2008**, *73*, 2224. Domingo, L. R.; Sáez, J. A.; Zaragoza, R. J.; Arnó, M. *J. Org. Chem.* **2008**, *73*, 8791. Bentabed-Ababsa, G.; Derdour, A.; Roisnel, T.; Saéz, J. A.; Pérez, P.; Chamorro, E.; Domingo, L. R.; Mongin, F. *J. Org. Chem.* **2009**, *74*, 2120. Benchouk, W.; Mekelleche, S. M.; Aurell, M. J.; Domingo, L. R. *Tetrahedron* **2009**, *911*, 4644. Brasca, R.; Kneeteman, M. N.; Mancini, P. M. E.; Fabian, W. M. F. *THEOCHEM* **2009**, *911*, 124. Domingo, L. R.; Chamorro, E.; Pérez, P. *Eur. J. Org. Chem.* **2009**, 3036. Das, T. K.; Banerjee, M. *J. Phys. Org. Chem.* **2010**, *23*, 148. Acharjee, N.; Das, T. K.; Banerji, A.; Banerjee, M.; Prange, T. *J. Phys. Org. Chem.* **2010**, DOI 10.1002/poc.1690. Das, T. K.; Salampuria, S.; Banerjee, M. *THEOCHEM* **2010**, 959, 22. Jasiński, R.; Barański, A. *THEOCHEM* **2010**, 949, 8. Acharjee, N.; Banerji, A.; Prangé, T. *Monatsh Chem.* **2010**, DOI: 10.1007/s00706-010-0393-2. Bakavoli, M.; Moeinpour, F.; Davoodnia, A.; Morsali, A. *J. Mol. Struct.* **2010**, 969, 139. Monbaliu, J. C.; Dive, G.; Marchand-Brynaert, J.; Peeters, D. *THEOCHEM* **2010**, 959, 49. Soto-Delgado, J.; Domingo, L. R.; Contreras, R. *Org. Biomol. Chem.* **2010**, *8*, 3678. Rhyman, L.; Abdallah, H. H.; Jhaumeer-Laulloo, S.; Domingo, L. R.; Joule, J. A.; Ramasami, P. *Tetrahedron* **2010**, DOI: 10.1016/j.tet.2010.09.071. Nacereddine, A. K.; Yahia, W.; Bouacha, S.; Djerourou, A. *Tetrahedron Lett.* **2010**, *51*, 2617.
- (323) Domingo, L. R.; Picher, M. T. *Tetrahedron* **2004**, *60*, 5053. See also: Merino, P.; Tejero, T.; Chiacchio, U.; Romeo, G.; Rescifina, A. *Tetrahedron* **2007**, *63*, 1448.
- (324) Aurell, M. J.; Domingo, L. R.; Pérez, P.; Contreras, R. *Tetrahedron* **2004**, *60*, 11503. See also: Domingo, L. R.; Picher, M. T.; Arroyo, P.; Saez, J. A. *J. Org. Chem.* **2006**, *71*, 9319.
- (325) Domingo, L. R. *Eur. J. Org. Chem.* **2004**, 4788. For a recent review, see: Ess, D. H.; Jones, G. O.; Houk, K. N. *Adv. Synth. Catal.* **2006**, *348*, 2337. Alves, C. N.; Carneiro, A. S.; Andrés, J.; Domingo, L. R. *Tetrahedron* **2006**, *92*, 5502. Domingo, L. R.; Aurell, M. J.; Kneeteman, M. N.; Mancini, P. M. *THEOCHEM* **2008**, *853*, 68. Sousa, C. A. D.; Vale, M. L. C.; Rodríguez-Borges, J. E.; Garcia-Mera, X. *Tetrahedron Lett.* **2008**, *49*, 5777. Alajarín, M.; Cabrera, J.; Pastor, A.; Sánchez-Andrada, P.; Bautista, D. *J. Org. Chem.* **2008**, *73*, 963. Domingo, L. R.; Chamorro, E.; Pérez, P. *J. Phys. Chem. A* **2008**, *112*, 4046. Aghabozorg, H.; Moradi, S.; Fereyduni, E.; Khani, H.; Yaaghubi, E. *THEOCHEM* **2009**, *915*, 58. Domingo, L.; Sáez, J. *Org. Biomol. Chem.* **2009**, *7*, 3576. Soto-Delgado, J.; Domingo, L. R.; Contreras, R. *THEOCHEM* **2009**, *902*, 103. Jasiński, R.; Kwiatkowska, M.; Barański, A. *THEOCHEM* **2009**, *910*, 80. Domingo, L. R.; Tera Pichar, S.; Sáez, J. A. *J. Org. Chem.* **2009**, *74*, 2726. Sharma, P.; Kumar, A.; Sahu, V. *J. Phys. Chem. A* **2010**, *114*, 1032.
- (326) Domingo, L. R.; Jose Aurell, M.; Pérez, P.; Contreras, R. *J. Org. Chem.* **2003**, *68*, 3884.
- (327) Domingo, L. R.; Andrés, J. *J. Org. Chem.* **2003**, *68*, 8662.
- (328) Spino, C.; Rezaei, H.; Dory, Y. L. *J. Org. Chem.* **2004**, *69*, 757. See also ref 268. See also: Gomez-Bengoa, E.; Helm, M. D.; Plant, A.; Harry, J. P. A. *J. Am. Chem. Soc.* **2007**, *129*, 2691.
- (329) Domingo, L. R.; Zaragoza, R. J.; Williams, R. M. *J. Org. Chem.* **2003**, *68*, 2895.
- (330) Domingo, L. R. *Tetrahedron* **2002**, *58*, 3765.

- (331) Domingo, L. R.; Asensio, A.; Arroyo, P. *J. Phys. Org. Chem.* **2002**, *15*, 660.
- (332) Domingo, L. R.; Aurell, M. J.; Pérez, P.; Contreras, R. *Tetrahedron* **2002**, *58*, 4417. See also: Domingo, L. R.; Aurell, M. J.; Arno, M.; Saéz, J. A. *THEOCHEM* **2007**, *811*, 125.
- (333) Domingo, L. R.; Pérez, P.; Contreras, R. *Lett. Org. Chem.* **2005**, *2*, 68. Domingo, L. R.; Saéz, J. A.; Perez, P. *Chem. Phys. Lett.* **2007**, *438*, 341. Domingo, L. R.; Benchouk, W.; Mekelleche, S. M. *Tetrahedron* **2007**, *63*, 4464. Azzouzi, S.; Jalal, R.; El Messaoudi, M.; Domingo, L. R.; Esseffar, M.; Aurell, M. J. *J. Phys. Org. Chem.* **2007**, *20*, 245. Domingo, L. R.; Aurell, M. J.; Arno, M.; Saéz, J. A. *J. Org. Chem.* **2007**, *72*, 4220.
- (334) Domingo, L. R.; Aurell, M. J. *J. Org. Chem.* **2002**, *67*, 959. See also: Chemouri, H.; Benchouk, W.; Mekelleche, S. M. *J. Theor. Comp. Chem.* **2006**, *5*, 707. Domingo, L. R.; Picher, M. T.; Arroyo, P. *Eur. J. Org. Chem.* **2006**, 2570. Lecea, B.; Ayerbe, M.; Arrieta, A.; Cossio, F. P.; Branchadell, V.; Ortuno, R. M.; Baceiredo, A. J. *Org. Chem.* **2007**, *72*, 357.
- (335) Saéz, J. A.; Arno, M.; Domingo, L. R. *Org. Lett.* **2003**, *5*, 4117.
- (336) Corsaro, A.; Pistrà, V.; Rescifina, A.; Piperno, A.; Chiacchio, M. A.; Romeo, G. *Tetrahedron* **2004**, *60*, 6443.
- (337) Arnó, M.; Picher, M. T.; Domingo, L. R.; Andrés, J. *Chem.—Eur. J.* **2004**, *10*, 4742. See also: Domingo, L. R.; Sáez, J. A.; Palmucci, C.; Sepulveda-Arques, J.; González-Rosende, M. E. *Tetrahedron* **2006**, *62*, 10408. Domingo, L. R.; Arno, M.; Merino, P.; Tejero, T. *Eur. J. Org. Chem.* **2006**, 3464.
- (338) Domingo, L. R.; Arno, M.; Contreras, R.; Pérez, P. *J. Phys. Chem. A* **2002**, *106*, 952.
- (339) Benchouk, W.; Mekelleche, S. M. *THEOCHEM* **2008**, *1*, 862.
- (340) Lakhdar, S.; Goumont, R.; Berionni, G.; Boubaker, T.; Kurbatov, S.; Terrier, F. *Chem.—Eur. J.* **2007**, *13*, 8317. Lakhdar, S.; Terrier, F.; Vichard, D.; Berionni, G.; Guesmi, N. E.; Goumont, R.; Boubaker, T. *Chem.—Eur. J.* **2010**, *16*, 5681. Buncel, E.; Terrier, F. *Org. Biomol. Chem.* **2010**, *8*, 2285.
- (341) (a) Nenajdenko, V. G.; Moiseev, A. M.; Balenkova, E. S. *Russ. Chem. Bull.* **2004**, *53*, 2241. (b) Nenajdenko, V. G.; Korotchenko, V. N.; Shastin, A. V.; Balenkova, E. S. *Russ. Chem. Bull.* **2004**, *53*, 1034. The reactivity of the carbonyl group in cyclohexanone derivatives is analyzed in Nazari, F.; Zalia, F. R. *THEOCHEM* **2007**, *817*, 11.
- (342) Nenajdenko, V. G.; Korotchenko, V. N.; Shastin, A. V.; Tyurin, D. A.; Balenkova, E. S. *Russ. J. Org. Chem.* **2004**, *40*, 1750.
- (343) Araya-Maturana, R.; Heredia-Moya, J.; Escobar, C. A.; Pérez, P. *J. Chil. Chem. Soc.* **2004**, *49*, 35. See also: Rai, V.; Nambhothiri, I. N. N. *Eur. J. Org. Chem.* **2006**, 4693. Sobarzo-Sánchez, E.; Castedo, L.; De la Fuente, J. R. *Struct. Chem.* **2006**, *17*, 483.
- (344) Rivas, P.; Zapata-Torres, G.; Melin, J.; Contreras, R. *Tetrahedron* **2004**, *60*, 4189.
- (345) (a) Elango, M.; Parthasarathi, R.; Subramanian, V.; Sarkar, U.; Chattaraj, P. K. *THEOCHEM* **2005**, *723*, 43. (b) Krishnamoorthy, B. S.; Lalita, R.; Panchanatheswaran, K.; Chattaraj, P. K. *THEOCHEM* **2006**, *761*, 89. (c) Akin, F. A.; Ervin, K. M. *J. Phys. Chem. A* **2006**, *110*, 1342. (d) Sakurai, Y.; Shiozaki, H.; Yokoyama, M. *THEOCHEM* **2006**, *766*, 41. This paper highlights the power of the philicity concept. Philicity is also successfully used in analyzing the reactivity of furanones. Benmeddah, A.; Mekelleche, S. M.; Benchouk, W.; Mostefa-Kara, B.; Villemin, D. *THEOCHEM* **2007**, *821*, 42.
- (346) Domingo, L. R.; Pérez, P.; Contreras, R. *Tetrahedron* **2004**, *60*, 6585.
- (347) (a) Meneses, L.; Fuentealba, P.; Contreras, R. *Tetrahedron* **2005**, *61*, 831. (b) See also: Chattaraj, P. K.; Sarkar, U.; Elango, M.; Parthasarathi, R.; Subramanian, V. *Los Alamos Natl. Lab., Prepr. Arch., Chem. Phys.* **2005**, 1–38, arXiv:physics/0509089 for the state-of-the-art analysis. Generally, some outliers are to be removed to obtain a good correlation. Sometimes a linear combination of ω and $\omega_{\text{el}}^{\text{a}}$ provides a better correlation, albeit with a possibility of collinearity. (c) Roy, D. R.; Parthasarathi, R.; Padmanabhan, J.; Sarkar, U.; Subramanian, V.; Chattaraj, P. K. *J. Phys. Chem. A* **2006**, *110*, 1084. (d) Chattaraj, P. K.; Sarkar, U.; Roy, D. R.; Elango, M.; Parthasarathi, R.; Subramanian, V. *Ind. J. Chem.* **2006**, *45A*, 1099. (e) It has been shown that there is no correlation between ω and ΔG^{\ddagger} for the metathesis reactions: Fomine, S.; Ortega, J. V.; Tlenkopatchev, M. A. *J. Organomet. Chem.* **2006**, *691*, 3343. (f) Crawford, P.; Hu, P. *J. Phys. Chem. B* **2006**, *110*, 4157. See also: Pavez, P.; Herrera, B.; Castro, E. A.; Santos, J. G.; Toro-Labbe, A. *THEOCHEM* **2007**, *811*, 91. Zhong, A.; Rong, C.; Liu, S. *J. Phys. Chem. A* **2007**, *111*, 3132. Latelli, N.; Zeroual, S.; Ouddai, N.; Mokhtari, M.; Ciofini, I. *Chem. Phys. Lett.* **2008**, *461*, 16. For a handwaving argument, see: Bagaria, P.; Roy, R. K. *J. Phys. Chem. A* **2008**, *112*, 97. Bagaria, P.; Saha, S.; Murru, S.; Kavala, V.; Patel, B. K.; Roy, R. K. *Phys. Chem. Chem. Phys.* **2009**, *11*, 8306. Fuentealba, P.; David, J.; Guerra, D. *THEOCHEM* **2010**, *943*, 127.
- (348) Capriati, V.; Florio, S.; Luisi, R.; Rocchetti, M. T. *J. Org. Chem.* **2002**, *67*, 759. An analysis of the bond-forming ability of a metal is available in: Crawford, P.; Hu, P. *J. Chem. Phys.* **2007**, *126*, 194706.
- (349) Griveau, S.; Bedioui, F.; Adamo, C. *J. Phys. Chem. A* **2001**, *105*, 11304. For the reactivity of thiodiazoles, see: Glossman-Mitnik, D. *THEOCHEM* **2005**, *725*, 27.
- (350) Galabov, B.; Cheshmedzhieva, D.; Ilieva, S.; Hadjieva, B. *J. Phys. Chem. A* **2004**, *108*, 11457.
- (351) Leach, A. G.; Houk, K. N. *Org. Biomol. Chem.* **2003**, *1*, 1389.
- (352) Campodónico, P. R.; Fuentealba, P.; Castro, E. A.; Santos, J. G.; Contreras, R. *J. Org. Chem.* **2005**, *70*, 1754.
- (353) Mieusset, J. L.; Brinker, U. H. *J. Org. Chem.* **2008**, *73*, 1553. Campodónico, P. R.; Ormazábal-Toledo, R.; Aizman, A.; Contreras, R. *Chem. Phys. Lett.* **2010**, *498*, 221.
- (354) Campodónico, P. R.; Perez, C.; Aliaga, M.; Gazitua, M.; Contreras, R. *Chem. Phys. Lett.* **2008**, *447*, 375. Campodónico, P. R.; Ormazábal-Toledo, R.; Aizman, A.; Contreras, R. *Chem. Phys. Lett.* **2010**, *498*, 221.
- (355) Domingo, L. R.; Chamorro, E.; Perez, P. *J. Org. Chem.* **2008**, *73*, 4615.
- (356) (a) Pérez, P.; Toro-Labbé, A.; Contreras, R. *J. Am. Chem. Soc.* **2001**, *123*, 5527. (b) Meneses, L.; Fuentealba, P.; Contreras, R. *Chem. Phys. Lett.* **2006**, *433*, 54. See also: Pearson, R. G. *J. Am. Chem. Soc.* **1986**, *108*, 6109, where it has been shown that the χ values in gas and solution phases do not differ much whereas the corresponding η values become smaller on solvation.
- (357) Contreras, R.; Mendizábal, F.; Aizman, A. *Phys. Rev. A* **1994**, *49*, 4349.
- (358) Contreras, R.; Pérez, P.; Aizman, A. In *Solvent Effects and Chemical Reactivity*; Tapia, O., Bertrán, J., Eds.; Kluwer Academic Publishers: Dordrecht, The Netherlands, 1996; Vol. 17, p 81.
- (359) Pérez, P.; Contreras, R. *Chem. Phys. Lett.* **1996**, *260*, 236.
- (360) Chattaraj, P. K.; Pérez, P.; Zavallos, J.; Toro-Labbé, A. *THEOCHEM* **2002**, *580*, 171.
- (361) Chattaraj, P. K.; Pérez, P.; Zavallos, J.; Toro-Labbé, A. *J. Phys. Chem. A* **2001**, *105*, 4272.
- (362) Aparicio, F.; Contreras, R.; Galván, M.; Cedillo, A. *J. Phys. Chem. A* **2003**, *107*, 10098. (a) Hydration of Mg^{+2} is presented in Adrian-Scotto, M.; Mallet, G.; Vasilescu, D. *THEOCHEM* **2005**, *728*, 231.
- (363) Cimino, P.; Improta, R.; Bifulco, G.; Riccio, R.; Gomez-Paloma, L.; Barone, V. *J. Org. Chem.* **2004**, *69*, 2816.
- (364) Ciofini, I.; Hazebrucq, S.; Joubert, L.; Adamo, C. *Theor. Chem. Acc.* **2004**, *111*, 188.
- (365) Padmanabhan, J.; Parthasarathi, R.; Sarkar, U.; Subramanian, V.; Chattaraj, P. K. *Chem. Phys. Lett.* **2004**, *383*, 122.
- (366) Joubert, L.; Guillemoles, J.-F.; Adamo, C. *Chem. Phys. Lett.* **2003**, *371*, 378.
- (367) Caro, C. A.; Zagal, J. H.; Bedioui, F.; Adamo, C.; Cardenas-Jiron, G. I. *J. Phys. Chem. A* **2004**, *108*, 6045. Kesharwani, M. K.; Ganguly, B. *Croat. Chem. Acta.* **2009**, *82*, 291. Jaramillo, P.; Pérez, P.; Fuentealba, P.; Canuto, S.; Coutinho, K. *J. Phys. Chem. B* **2009**, *113*, 4314.
- (368) Parthasarathi, R.; Subramanian, V.; Chattaraj, P. K. *Chem. Phys. Lett.* **2003**, *382*, 48.
- (369) Ayers, P. W.; Parr, R. G. *J. Am. Chem. Soc.* **2001**, *123*, 2007.
- (370) Ayers, P. W.; Anderson, J. S. M.; Bartolotti, L. J. *Int. J. Quantum Chem.* **2005**, *101*, 520. (a) This paper has a bearing in the new electrophilicity definition provided by Gázquez et al.^{63c} See also: Ayers, P. W.; Melin, J. *Theor. Chem. Acc.* **2007**, *117*, 371.
- (371) Fuentealba, P.; Cedillo, A. *J. Chem. Phys.* **1999**, *110*, 9807.
- (372) Ciofini, I.; Bedioui, F.; Zagal, J. H.; Adamo, C. *Chem. Phys. Lett.* **2003**, *376*, 690.
- (373) Miller, J. A.; Miller, E. C. Ultimate Chemical Carcinogens as Reactive Mutagenic Electrophiles. In *Origins of Human Cancer*; Hiatt, H. H., Watson, J. D., Winsten, J. A., Eds.; Cold Spring Harbor Laboratory Press: Cold Spring Harbor, NY, 1977; pp 605–627.
- (374) (a) Ashby, J.; Tennant, R. W. *Mutat. Res.* **1991**, *257*, 229. (b) Rong, C.; Lian, S.; Yin, D.; Zhong, A.; Zhang, R.; Liu, S. *Chem. Phys. Lett.* **2007**, *434*, 149.
- (375) Rosenkranz, H. S.; Klopman, G.; Zhang, Y. P.; Graham, C.; Karol, M. H. *Environ. Health Perspect.* **1999**, *107*, 129.
- (376) Roberts, D. W.; York, M.; Basketter, D. A. *Contact Dermatitis* **1999**, *41*, 14.
- (377) Roberts, D. W.; Patlewicz, G. *SAR QSAR Environ. Res.* **2002**, *13*, 145.
- (378) Dinkova-Kostova, A. T.; Massiah, M. A.; Bozak, R. E.; Hicks, R. J.; Talalay, P. *Proc. Natl. Acad. Sci. U.S.A.* **2001**, *98*, 3404.
- (379) Cronin, M. T. D.; Netzeva, T. I.; Dearden, J. C.; Edwards, R.; Worgan, A. D. *P. Chem. Res. Toxicol.* **2004**, *17*, 545.
- (380) Nakamura, Y.; Kumagai, T.; Yoshida, C.; Naito, Y.; Miyamoto, M.; Ohigashi, H.; Osawa, T.; Uchida, K. *Biochemistry* **2003**, *42*, 4300.

- (381) (a) Banks, T. M.; Bonin, A. M.; Glover, S. A.; Prakash, A. S. *Org. Biomol. Chem.* **2003**, *1*, 2238. (b) Cimino, P.; Bifulco, G.; Riccio, R.; Gomez-Paloma, L.; Barone, V. *Org. Biomol. Chem.* **2006**, *4*, 1242.
- (382) (a) Mendoza-Wilson, A. M.; Glossman-Mitnik, D. *THEOCHEM* **2005**, *716*, 67. (b) Mendoza-Wilson, A. M.; Glossman-Mitnik, D. *THEOCHEM* **2006**, *761*, 97.
- (383) (a) Laufer, R. S.; Dmitrienko, G. I. *J. Am. Chem. Soc.* **2002**, *124*, 1854. (b) Pasha, F. A.; Srivastava, H. K.; Beg, Y.; Singh, P. P. *Am. J. Immun.* **2006**, *2*, 23. For the analysis of antioxidant properties using some of these descriptors, see: Reis, M.; Lobato, B.; Lameira, J.; Santos, A. S.; Alves, C. N. *Eur. J. Med. Chem.* **2007**, *42*, 440. Calgarotto, A. K.; Miotto, S.; Honorio, K. M.; Da Silva, A. B. F.; Marangoni, S.; Silva, J. L.; Comar, M.; Oliveira, K. M. T.; Da Silva, S. L. *THEOCHEM* **2007**, *808*, 25. And for that of CDK2 inhibitors, see: Alzate-Morales, J. H.; Tiznado, W.; Santos, J. C.; Cardenas, C.; Contreras, R. *J. Phys. Chem. B* **2007**, *111*, 3293. (f) They are also shown to be useful in understanding the selective cancer cells membrane lysis associated with specific fragments of amino acids derived from PNC-27: Barrientos-Salcedo, C.; Arenas-Aranda, D.; Salamanca-Gomez, F.; Ortiz-Muniz, R.; Soriano-Correa, C. *J. Phys. Chem. A* **2007**, *111*, 4362.
- (384) Rétey, J. *Biochim. Biophys. Acta* **2003**, *1647*, 179.
- (385) Svensson, R.; Pamedyite, V.; Juodaityte, J.; Makuška, R.; Morgenstern, R. *Toxicology* **2001**, *168*, 251.
- (386) Ramirez-Arizmendi, L. E.; Heidbrink, J. L.; Guler, L. P.; Kenttamaa, H. I. *J. Am. Chem. Soc.* **2003**, *125*, 2272.
- (387) Tapia, R. A.; Salas, C.; Morello, A.; Maya, J. D.; Toro-Labbé, A. *Bioorg. Med. Chem.* **2004**, *12*, 2451.
- (388) Håkansson, K. *Int. J. Biol. Macromol.* **2002**, *30*, 273.
- (389) Roos, G.; Messens, J.; Loverix, S.; Wyns, L.; Geerlings, P. *J. Phys. Chem. B* **2004**, *108*, 17216.
- (390) Wang, L. H.; Yang, X. Y.; Zhang, X.; Mihalic, K.; Fan, Y.-X.; Xiao, W.; Howard, O. M. Z.; Appella, E.; Maynard, A. T.; Farrar, W. L. *Nature Med.* **2004**, *10*, 40.
- (391) Parthasarathi, R.; Padmanabhan, J.; Subramanian, V.; Maiti, B.; Chattaraj, P. K. *J. Phys. Chem. A* **2003**, *107*, 10346. See also: Padmanabhan, J.; Parthasarathi, R.; Subramanian, V.; Chattaraj, P. K. *J. Phys. Chem. A* **2005**, *109*, 11043. (b) Padmanabhan, J.; Parthasarathi, R.; Subramanian, V.; Chattaraj, P. K. *J. Phys. Chem. A* **2006**, *110*, 2739. (c) Padmanabhan, J.; Parthasarathi, R.; Subramanian, V.; Chattaraj, P. K. *Chem. Res. Toxicol.* **2006**, *19*, 356.
- (392) Parthasarathi, R.; Padmanabhan, J.; Subramanian, V.; Maiti, B.; Chattaraj, P. K. *Curr. Sci.* **2004**, *86*, 535.
- (393) Arulmozhiraja, S.; Selvin, P. C.; Fujii, T. *J. Phys. Chem. A* **2002**, *106*, 1765.
- (394) Arulmozhiraja, S.; Fujii, T.; Sato, G. *Mol. Phys.* **2002**, *100*, 423.
- (395) (a) Parthasarathi, R.; Subramanian, V.; Roy, D. R.; Chattaraj, P. K. *Bioorg. Med. Chem.* **2004**, *12*, 5533. (b) Roy, D. R.; Pal, N.; Mitra, A.; Bultinck, P.; Parthasarathi, R.; Subramanian, V.; Chattaraj, P. K. *Eur. J. Med. Chem.* **2007**, *42*, 1365.
- (396) Arulmozhiraja, S.; Morita, M. *Chem. Res. Toxicol.* **2004**, *17*, 348. See also: Gu, C.; Jiang, X.; Ju, X.; Yu, G.; Bian, Y. *Chemosphere* **2007**, *67*, 1325, for a better correlation.
- (397) (a) Roy, D. R.; Parthasarathi, R.; Maiti, B.; Subramanian, V.; Chattaraj, P. K. *Bioorg. Med. Chem.* **2005**, *13*, 3405. (b) Roy, D. R.; Sarkar, U.; Chattaraj, P. K.; Mitra, A.; Padmanabhan, J.; Parthasarathi, R.; Subramanian, V.; Vandamme, S.; Bultinck, P. *Mol. Diversity* **2006**, *10*, 119. (c) Chattaraj, P. K.; Roy, D. R.; Giri, S.; Mukherjee, S.; Subramanian, V.; Parthasarathi, R.; Bultinck, P.; Van Damme, S. *J. Chem. Sci.* **2007**, *119*, 475 (Theor. Chem. Special Issue) (d) Giri, S.; Roy, D. R.; Bultinck, P.; Subramanian, V.; Chattaraj, P. K. *Los Alamos Natl. Lab., Prepr. Arch., Chem. Phys.* **2006**, 1–35, arXiv: physics/0611204. (e) Padmanabhan, J.; Parthasarathi, R.; Subramanian, V.; Chattaraj, P. K. *QSAR Comb. Sci.* **2007**, *26*, 227. (f) Lameira, J.; Medeiros, I. G.; Reis, M.; Santos, A. S.; Alves, C. N. *Bioorg. Med. Chem.* **2006**, *14*, 7105. (g) Glossman-Mitnik, D. *J. Mol. Graphics Modell.* **2006**, *25*, 455. (h) Boiani, M.; Boiani, L.; Denicola, A.; Torres de Ortiz, S.; Serna, E.; Vera de Bilbao, N.; Sanabria, L.; Yaluff, G.; Nakayama, H.; Rojas de Arias, A.; Vega, C.; Rolan, M.; Gómez-Barrio, A.; Cerecetto, H.; González, M. *J. Med. Chem.* **2006**, *49*, 3215. (i) Brasil, D. S. B.; Moreira, R. Y. O.; Muller, A. H.; Alves, C. N. *Int. J. Quantum Chem.* **2006**, *106*, 2706. (j) Soriano-Correa, C.; Ruiz, J. F. S.; Raya, A.; Esquivel, R. O. *Int. J. Quantum Chem.* **2007**, *107*, 628. (k) Taskin, T.; Sevin, F. *THEOCHEM* **2007**, *803*, 61. (l) Toropov, A. A.; Benfenati, E. *Bioorg. Med. Chem.* **2006**, *14*, 3923. (m) Roohi, H.; Nowroozi, A.; Ebrahimi, A.; Makiabadi, B. *J. Mol. Struct. (THEOCHEM)* **2010**, *952*, 36. (n) Gómez-Carrillo, S. C.; Ortega, J.; Bolcatto, P. G. *J. Phys.: Conf. Ser.* **2009**, *167*, 012062. (o) Anantharaj, R.; Banerjee, T. *Fluid Phase Equilib.* **2010**, *293*, 22. (p) Boughdiri, S.; Essalah, K. *J. Mol. Struct. (THEOCHEM)* **2010**, *958*, 21.
- (398) (a) Sarkar, U.; Roy, D. R.; Chattaraj, P. K.; Parthasarathi, R.; Padmanabhan, J.; Subramanian, V. *J. Chem. Sci.* **2005**, *117*, 599.
- (b) Roy, D. R.; Parthasarathi, R.; Subramanian, V.; Chattaraj, P. K. *QSAR Comb. Sci.* **2006**, *25*, 114. (c) Sarkar, U.; Padmanabhan, J.; Parthasarathi, R.; Subramanian, V.; Chattaraj, P. K. *THEOCHEM* **2006**, *758*, 119. (d) Parthasarathi, R.; Elango, M.; Padmanabhan, J.; Subramanian, V.; Roy, D. R.; Sarkar, U.; Chattaraj, P. K. *Ind. J. Chem.* **2006**, *45A*, 111. (e) Padmanabhan, J.; Parthasarathi, R.; Subramanian, V.; Chattaraj, P. K. *Bioorg. Med. Chem.* **2006**, *14*, 1021. (f) For a similar study on chloroanisolles, see: Padmanabhan, J.; Parthasarathi, R.; Subramanian, V.; Chattaraj, P. K. *THEOCHEM* **2006**, *774*, 49.
- (399) Echeverria, C.; Santibañez, J. F.; Donoso-Tauda, O.; Escobar, C. A.; Ramirez-Tagle, R. *Int. J. Mol. Sci.* **2009**, *10*, 221.
- (400) Sarmah, P.; Deka, R. C. *J. Comput.-Aided Mol. Des.* **2009**, *23*, 343. For other related studies, see: da Costa, E. B.; Trsic, M. *J. Mol. Graphics Modell.* **2010**, *28*, 657. Payán-Gómez, S. A.; Flores-Holguín, N.; Pérez-Miranda, A.; Piñón-Miramontes, M.; Glossman-Mitnik, D. *Chem. Central J.* **2010**, *4*, 12. Mineva, T.; Russo, N. *THEOCHEM* **2010**, *943*, 71. Sarmah, P.; Deka, R. C. *J. Mol. Model.* **2010**, *16*, 411. Saikia, N.; Deka, R. C. *Chem. Phys. Lett.* **2010**, DOI: 10.1016/j.cplett.2010.09.075. Saracoglu, M.; Kandemirli, F.; Kovalishyn, V.; Arslan, T.; Ebenso, E. E. *J. Biomed. Biotech.* **2010**, DOI: 10.1155/2010/693031. Bensassona, R. V.; Zoeteb, V.; Berthier, G.; Talalay, P.; Dinkova-Kostova, A. T. *Chem.-Biol. Interact.* **2010**, *186*, 118. Wondrousch, D.; Böhme, A.; Thaens, D.; Ost, N.; Schüttürmann, G. *J. Phys. Chem. Lett.* **2010**, *1*, 1605.
- (401) Sarmah, P.; Deka, R. C. *Int. J. Quantum Chem.* **2008**, *108*, 1400. Hazarika, K. K.; Baruah, N. C.; Deka, R. C. *Struct. Chem.* **2009**, *20*, 1079.
- (402) Srivastava, H. K.; Pasha, F. A.; Mishra, S. K.; Singh, P. P. *Med. Chem. Res.* **2009**, *18*, 455.
- (403) LoPachin, R. M.; Geohagen, B. C.; Gavin, T. *Toxicol. Sci.* **2009**, *107*, 171.
- (404) Zhao, Y. Y.; Tao, F. M.; Zeng, E. Y. *Chemosphere* **2008**, *73*, 86.
- (405) Ishihara, M.; Fujisawa, S. *Chemosphere* **2008**, *70*, 1898.
- (406) Alonso, M.; Casado, S.; Miranda, C.; Tarazona, J. V.; Navas, J. M.; Herradón, B. *Chem. Res. Toxicol.* **2008**, *21*, 643.
- (407) Feng, X. T.; Yu, J. G.; Lei, M.; Fang, W. H.; Liu, S. J. *Phys. Chem. B* **2009**, *113*, 13381.
- (408) Bian, H.; Chen, J.; Cai, X.; Liu, P.; Wang, Y.; Huang, L.; Qiao, X.; Hao, C. *Water Res.* **2009**, *43*, 2566.
- (409) Huang, Y.; Zhong, A.; Rong, C.; Xiao, X.; Liu, S. J. *Phys. Chem. A* **2008**, *112*, 305. For a study on NCN-pincer carbene complex, see: Das Adhikary, S.; Samanta, T.; Roymahapatra, G.; Loiseau, F.; Jouvenot, D.; Giri, S.; Chattaraj, P. K.; Dinda, J. *New J. Chem.* **2010**, *34*, 1974.
- (410) Librando, V.; Alparone, A. *J. Hazard. Mater.* **2009**, *161*, 1338.
- (411) Pasha, F. A.; Muddassar, M.; Cho, S. J.; Ahmad, K.; Beg, Y. *Eur. J. Med. Chem.* **2008**, *43*, 2361.
- (412) Pasha, F. A.; Cho, S. J.; Beg, Y.; Tripathi, Y. B. *Med. Chem. Res.* **2007**, *16*, 408.
- (413) Barrientos-Salcedo, C.; Arenas-Aranda, D.; Salamanca-Gómez, F.; Ortiz-Muñiz, R.; Soriano-Correa, C. *J. Phys. Chem. A* **2007**, *111*, 4362.
- (414) Eroglu, E.; Türkmen, H. *J. Mol. Graphics Modell.* **2007**, *26*, 701.
- (415) Enoch, S. J.; Cronin, M. T. D.; Schultz, T. W.; Madden, J. C. *Chem. Res. Toxicol.* **2008**, *21*, 513.
- (416) LoPachin, R. M.; Gavin, T.; Geohagen, B. C.; Das, S. *Toxicol. Sci.* **2007**, *98*, 561.
- (417) Roy, D. R.; Giri, S.; Chattaraj, P. K. *Mol. Diversity* **2009**, *13*, 551.
- (418) Campodonico, P. R.; Contreras, R. *Bioorg. Med. Chem.* **2008**, *16*, 3184.
- (419) Pasha, F. A.; Chung, H. W.; Kang, S. B.; Cho, S. J. *Int. J. Quantum Chem.* **2008**, *108*, 391.
- (420) Pasha, F. A.; Muddassar, M.; Lee, C.; Cho, S. J. *Environ. Toxicol. Pharmacol.* **2008**, *26*, 128.
- (421) Soriano-Correa, C.; Raya, A.; Esquivel, R. O. *Int. J. Quantum Chem.* **2008**, *108*, 1369.
- (422) Gupta, K.; Giri, S.; Chattaraj, P. K. *New J. Chem.* **2008**, *32*, 1945.
- (423) Giri, S.; Roy, D. R.; Van Damme, S.; Bultinck, P.; Subramanian, V.; Chattaraj, P. K. *QSAR Comb. Sci.* **2008**, *27*, 208.
- (424) Towler, C. S.; Li, T.; Wikstrom, H.; Remick, D. M.; Sanchez-Felix, M. V.; Taylor, L. S. *Mol. Pharmaceutics* **2008**, *5*, 946.
- (425) Da Silva Miranda, F.; Signori, A. M.; Vicente, J.; de Souza, B.; Priebe, J. P.; Szpoganicz, B.; Goncalves, N. S.; Neves, A. *Tetrahedron* **2008**, *64*, 5410.
- (426) Parthasarathi, R.; Elango, M.; Subramanian, V.; Chattaraj, P. K. *Theor. Chem. Acc.* **2005**, *113*, 257.
- (427) Cadet, J.; Grand, A.; Morell, C.; Letelier, J. R.; Moncada, J. L.; Toro-Labbé, A. *J. Phys. Chem. A* **2003**, *107*, 5334.
- (428) Zevallos, J.; Letelier, J. R.; Toro-Labbé, A. *J. Phys. Chem. A* **2004**, *108*, 10186.

- (429) (a) Herrera, B.; Toro-Labbé, A. *J. Phys. Chem. A* **2004**, *108*, 1830.
(b) Noorizadeh, S.; Maihami, H. *THEOCHEM* **2006**, 763, 133. (c) Noorizadeh, S.; Shakerzadeh, E. *Chem. Phys. Lett.* **2010**, *484*, 363.
- (430) Morell, C.; Labet, V.; Grand, A.; Chermette, H. *Phys. Chem. Chem. Phys.* **2009**, *11*, 3414.
- (431) Chaquin, P. *Chem. Phys. Lett.* **2008**, *458*, 231.
- (432) Chattaraj, P. K.; Gutierrez-Oliva, S.; Jaque, P.; Toro-Labbé, A. *Mol. Phys.* **2003**, *101*, 2841.
- (433) (a) Chattaraj, P. K.; Roy, D. R. *J. Phys. Chem. A* **2005**, *109*, 3771.
(b) For a Hammond postulate in terms of reactivity descriptors, see: Chattaraj, P. K.; Roy, D. R. *J. Phys. Chem. A* **2006**, *110*, 11401.
- (434) Chattaraj, P. K.; Sengupta, S. *Phys. Lett. A* **1993**, *181*, 225.
- (435) Sengupta, S.; Chattaraj, P. K. *Phys. Lett. A* **1996**, *215*, 119.
- (436) Runge, E.; Gross, E. K. U. *Phys. Rev. Lett.* **1984**, *52*, 997.
- (437) Dhara, A. K.; Ghosh, S. K. *Phys. Rev. A* **1987**, *35*, 442.
- (438) Deb, B. M.; Chattaraj, P. K. *Phys. Rev. A* **1989**, *39*, 1696.
- (439) Deb, B. M.; Chattaraj, P. K.; Mishra, S. *Phys. Rev. A* **1991**, *43*, 1248.
- (440) Chattaraj, P. K. *Int. J. Quantum Chem.* **1992**, *41*, 845.
- (441) Dey, B. K.; Deb, B. M. *Int. J. Quantum Chem.* **1995**, *56*, 707.
- (442) Aizman, A.; Contreras, R.; Pérez, P. *Tetrahedron* **2005**, *61*, 889.
- (443) Domingo, L. R.; Pérez, P.; Contreras, R. *J. Org. Chem.* **2003**, *68*, 6060.
- (444) Lalevee, J.; Allonas, X.; Fouassier, J.-P. *J. Phys. Chem. A* **2004**, *108*, 4326.
- (445) Lu, H.-F.; Sun, Y.-C. *Surf. Sci. Lett.* **2001**, *494*, L787.
- (446) Arnaut, L. G.; Pais, A. A. C. C.; Formosinho, S. J.; Barroso, M. *J. Am. Chem. Soc.* **2003**, *125*, 5236. Barroso, M.; Chattopadhyay, N.; Klymchenko, A. S.; Demchenko, A. P.; Arnaut, L. G.; Formosinho, S. J. *J. Phys. Chem. A* **2006**, *110*, 13419. Arnaut, L. G.; Formosinho, S. J.; Barroso, M. *J. Mol. Struct.* **2006**, *786*, 207. Barroso, M.; Arnaut, L. G.; Formosinho, S. J. *J. Phys. Chem. A* **2007**, *111*, 591.
- (447) Barroso, M.; Arnaut, L. G.; Formosinho, S. J. *Chem. Phys. Chem.* **2005**, *6*, 363. See also: Barroso, M.; Pereira, J. C.; Pais, A. A. C. C.; Arnaut, L. G.; Formosinho, S. J. *Mol. Phys.* **2006**, *104*, 731. Arnaut, L. G.; Formosinho, S. J. *Chem.—Eur. J.* **2007**, *13*, 8018.
- (448) Barroso, M.; Arnaut, L. G.; Formosinho, S. J. *J. Photochem. Photobiol., A* **2002**, *154*, 13.
- (449) Galván, M.; Vela, A.; Gázquez, J. L. *J. Phys. Chem.* **1988**, *92*, 6470.
- (450) Galván, M.; Vargas, R. *J. Phys. Chem.* **1992**, *96*, 1625.
- (451) Vargas, R.; Galván, M. *J. Phys. Chem.* **1996**, *100*, 14651.
- (452) Vargas, R.; Galván, M.; Vela, A. *J. Phys. Chem. A* **1998**, *102*, 3134.
- (453) Vargas, R.; Cedillo, A.; Garza, J.; Galván, M. In *Reviews of Modern Quantum Chemistry, A Celebration to the Contributions of R. G. Parr*; Sen, K. D., Eds.; World Scientific: Singapore, 2002; p 936.
- (454) Ghanty, T. K.; Ghosh, S. K. *J. Am. Chem. Soc.* **1994**, *116*, 3943.
- (455) Garza, J.; Vargas, R.; Cedillo, A.; Galván, M.; Chattaraj, P. K. *Theor. Chem. Acc.* **2006**, *115*, 257.
- (456) Pérez, P.; Andres, J.; Safont, V. S.; Tapia, O.; Contreras, R. *J. Phys. Chem. A* **2002**, *106*, 5353.
- (457) Melin, J.; Aparicio, F.; Galván, M.; Fuentealba, P.; Contreras, R. *J. Phys. Chem. A* **2003**, *107*, 3831.
- (458) Olah, J.; De Proft, F.; Veszpremi, T.; Geerlings, P. *J. Phys. Chem. A* **2004**, *108*, 490.
- (459) Olah, J.; De Proft, F.; Veszpremi, T.; Geerlings, P. *J. Phys. Chem. A* **2005**, *109*, 1608.
- (460) De Proft, F.; Fias, S.; Van Alsenoy, C.; Geerlings, P. *J. Phys. Chem. A* **2005**, *109*, 6335. See also: Chamorro, E.; Pérez, P. *J. Chem. Phys.* **2005**, *123*, 114107. Chamorro, E.; Pérez, P.; De Proft, F.; Geerlings, P. *J. Chem. Phys.* **2006**, *124*, 044105. Chamorro, E.; Santos, J. C.; Escobar, C. A.; Perez, P. *Chem. Phys. Lett.* **2006**, *431*, 210. Oláh, J.; De Proft, F.; Veszpremi, T.; Geerlings, P. *THEOCHEM* **2006**, *771*, 135. Pintér, B.; De Proft, F.; Van Speybroeck, V.; Hemelsoet, K.; Waroquier, M.; Chamorro, E.; Veszpremi, T.; Geerlings, P. *J. Org. Chem.* **2007**, *72*, 348. Rincón, E.; Pérez, P.; Chamorro, E. *Chem. Phys. Lett.* **2007**, *448*, 273. Guerra, D.; Andrés, J.; Chamorro, E.; Pérez, P. *Theor. Chem. Acc.* **2007**, *118*, 325. Chamorro, E.; Pérez, P.; Duque, M.; De Proft, F.; Geerlings, P. *J. Chem. Phys.* **2008**, *129*, 064117. De Proft, F.; Chamorro, E.; Pérez, P.; Duque, M.; Vleeschouwer, F. D.; Geerlings, P. Spin-polarized reactivity indices from density functional theory and applications. In *Chemical Modelling: Applications and Theory*; Springborg, M., Ed.; RSC Publishing: Cambridge, U.K., 2009; Vol. 6, p 63. De Vleeschouwer, F.; De Proft, F.; Geerlings, P. *THEOCHEM* **2010**, *943*, 94. Pérez, P.; Chamorro, E. *THEOCHEM* **2010**, *943*, 110.

CR100149P